

THE LARYNGOSCOPE.

VOL. XXVI. ST. LOUIS, APRIL, 1916.

No. 4.

To the Readers of THE LARYNGOSCOPE:

Many of the foreign general and special medical journals continue to be issued at irregular intervals, with very meagre contents, and some of those that had temporarily suspended publication seem not yet to have revived. In those that continue either steady or sporadic publication there has, in the past year, sprung up a considerable literature relating to otolaryngologic surgery from bullet, shrapnel and other wounds sustained in war. In the compilation of the present Index Medicus we have tried to have the bibliography in this field of surgery as complete as circumstances would permit.

The amount of American oto-laryngologic literature is as large as ever and hence it has compelled us, because of lack of space, to curtail the subject-matter in our abstract department.

We again invite our readers to send us any suggestions or recommendations tending towards the improvement of the Index Medicus.

Very respectfully,

PHILIP FRANK,
Assistant Editor.

OUTLINE OF INDEX-MEDICUS.

I. NOSE AND NASO-PHARYNX.

- | | |
|--------------|--------------------------------|
| 1 Septum. | 5 Neoplasms (Nasal). |
| 2 Turbinals. | 6 Neoplasms (Naso-pharyngeal). |
| 3 Adenoids. | 7 General. |
| 4 Neuroses. | |

II. MOUTH AND PHARYNX.

- | | |
|------------|------------|
| 1 Palate. | 4 Tongue. |
| 2 Tonsils. | 5 General. |
| 3 Uvula. | |

III. ACCESSORY SINUSES.

- | | |
|----------------------|---------------------|
| 1 Frontal Sinus. | 4 Maxillary Antrum. |
| 2 Sphenoid Cells. | 5 General. |
| 3 Ethmoid Labyrinth. | |

IV. LARYNX. BRONCHI. TRACHEA. ESOPHAGUS.

- | | |
|-----------------------------|--------------|
| 1 Epiglottis. | 5 Esophagus. |
| 2 Voice and Speech Defects. | 6 Asthma. |
| 3 Larynx. | 7 Endoscopy. |
| 4 Trachea and Bronchi. | |

V. DIPHTHERIA AND THYROID GLAND.

- | | |
|---------------|------------------|
| 1 Diphtheria. | 2 Thyroid Gland. |
|---------------|------------------|

VI. EAR.

- | | |
|--------------------|-----------------------|
| 1 External Ear. | 5 Labyrinth. |
| 2 Tympanic Cavity. | 6 Defects of Hearing. |
| 3 Ossicles. | 7 General. |
| 4 Eustachian Tube. | |

VII. MASTOID AND INTRA-CRANIAL COMPLICATIONS.

- | | |
|-----------------------------------------------------|---------------|
| 1 Mastoid. | 4 Meningitis. |
| 2 Lateral Sinus. | 5 General. |
| 3 Intra-cranial Complications
and Brain Abscess. | |

VIII. NEW INSTRUMENTS.

- | | |
|--------------------------|-----------------------------|
| 1 Ear. | 5 Larynx. Trachea. Bronchi. |
| 2 Nose and Naso-pharynx. | Esophagus. |
| 3 Mouth and Pharynx. | 6 General. |
| 4 Accessory Sinuses. | |

1915

Index-Medicus and Digest of Oto-Laryngology.

NOTE.—All titles marked with a * are abstracted under their respective numbers in the second section. All articles marked with a † have appeared as original papers in THE LARYNGOSCOPE. All articles marked with a § have been abstracted in THE LARYNGOSCOPE.

I. NOSE AND NASO-PHARYNX.

Septum.

- 1 BALLMANN, ERICH. Relations between the septum, ethmoid, etc. (Welche Beziehungen bestehen zwischen den Verbiegungen der Nasenscheidewand, den Nasenmuscheln und dem Siebbein?) *Inaug. Dissert.*, Univ. Giessen, March, 1915.
- 2 CARTER, WM. WESLEY. Two cases of depressed nasal deformity resulting from the submucous operation. *Medical Record*, May 15, 1915.
- 3 CHABERT, DR. Paraffin in nasal septum operations. (Décollement de la muqueuse par la paraffine dans les opérations sur la cloison nasale.) *Rev. de Laryngol., d'Otol. et de Rhinol.*, p. 270, Nos. 42-43, Oct. 31, 1915.
- 4 DEWEY, M. The etiology and treatment of some types of deflected nasal septum. *Internat. Jour. Orthodontia*, p. 547, Vol. 1, No. 11, Nov., 1915.
- 5 DIEBOLD, F. Perforation of nasal septum. *Corresp. Bl. f. Schweizer Aerzte*, Jan. 23, 1915.
- 6 EASTMAN, W. R. Operative perforation of nasal septum. *Milit. Surg.*, June, 1915.
- 7 FISHER, L. One thousand submucous resections of the nasal septum. *N. Y. Med. Jour.*, May 22, 1915.
- 8 GARRAGHAN, E. F. Deformities of nasal septum and their correction. *Ill. Med. Jour.*, June, 1915.
- 9 GRANT, J. D. Case of bent nose straightened by submucous rotation of the cartilaginous septum. *Proc. Roy. Soc. Med., Laryngol. Sec.*, p. 102, Vol. 8, No. 8, June, 1915.

- 10 HETT, G. S. Bullet wound through face with perforation of antra and septum; operation. *Proc. Laryngol. Section, Royal Soc. Med.*, Vol. 3, No. 6, April, 1915.
- 11 HILL, W. Angioma of the nasal septum. *Proc. Laryngol. Section, Royal Soc. Med.*, Vol. 8, No. 6, April, 1915.
- *12 LEWIS, F. O. Normal nasal septum and the pathology of deflections. *N. Y. Med. Jour.*, p. 736, April 10, 1915.
- *13 LEWIS, F. O. Transplantation of cartilage in the correction of septal deformities. *Ann. of Otol., Rhinol. and Laryngol.*, p. 488, Sept., 1915.
- 14 MARSHALL, G. M. Some methods and devices in correction of nasal septum that make for success. *Pa. Med. Jour.*, March, 1915.
- 15 MILLIGAN, W. A note on the after-treatment of cases of submucous resection of the nasal septum without the use of splints. *Proc. Royal Soc. Med., Laryngological Section*, Vol. 8, No. 5, March, 1915.
- 16 PRENN, J. Submucous resection as a means of obliterating superficial blood-vessels. *Bost. Med. and Surg. Jour.*, Vol. 172, No. 4, 1915.
- 17 ROTH, J. B. Nasal septum. *Northwest Med.*, Vol. 7, No. 7, July, 1915.
- 18 SHORE, F. E. V. Submucous resection of nasal septum. *Iowa State Med. Soc. Jour.*, March, 1915.
- 19 SMITH, CONRAD. Submucous resection of the nasal septum. *Jour. Ophthalmol., Otol. and Laryngol.*, p. 1018, Dec., 1915.
- *20 TOWNSEND, I. A practical method of correcting septal deformities. *Jour. Ophthal., Otol. and Laryngol.*, Feb. and June, 1915.
- 21 WARBRICK, J. C. Perforation of the nasal septum. *Buffalo Med. Jour.*, May, 1915, and *Jour. Ophthal. and Oto-Laryngol.*, June, 1915.
- †22 WARREN, E. L. Plastic operation for dislocated columnar cartilage of the nose. *THE LARYNGOSCOPE*, p. 80, Feb., 1915.
- 23 WARREN, E. L. The submucous operation. *St. Paul Med. Jour.*, Aug., 1915.

Turbinals.

- *24 CAVANAUGH, J. A. The inferior turbinate. *Ann. Otol., Rhinol. and Laryngol.*, p. 621, Sept., 1915.
- 25 COWGILL, N. M. The lower turbinate. *West. Med. Rev.*, Nov., 1915.
- 26 FREER, O. T. Inferior turbinate—Its flap resection to reduce it when obstructive. *Mich. State Med. Soc. Jour.*, Jan., 1915.
- 27 GRAHAM, C. I. Case of tuberculous inferior turbinate. *Proc. Laryngol. Section, Royal Soc. Med.*, Vol. 3, No. 6, April, 1915.
- 28 HARDOUR, LE MAR. Adenoids and otitis in infants. (Les végétations adénoïdes et les otites chez les nourrissons.) *Le Jour. de Med. et de Chir.*, May, 1915.
- *29 LEALE, GIUSEPPE. Inferior turbinate operations and the rational cure of hypertrophic rhinitis. (Sugli inconvenienti delle operazioni demolitrici sui turbinati inferiori e sui metodi per la cura razionale della rinite ipertrofica.) *Boll. d. Mal. dell'Orecchio*, p. 201, Vol. 33, No. 9, Sept., 1915.
- 30 REYNOLDS, C. W. Indications for and limitations of turbinectomy. *Kentucky Med. Jour.*, Oct., 1915.

- 31 TOB, H. Malignant papillomatous growth of left inferior turbinate and ethmoidal region. *Proc. Royal Soc. Med., Laryngol. Section*, Vol. 9, No. 2, Dec., 1915.
- 32 WEITH, M. S. A new conservative turbinate operation. *Jour. Iowa State Med. Soc.*, Jan. 15, 1915.

Adenoids.

- 33 BENJAMINS, C. E. Gritting the teeth a sign of adenoid vegetations. *Nederl. Tijdsch. v. Geneesk.*, Vol. 2, No. 2, July 10, 1915.
- 34 BENJAMINS, C. E. Teeth-grinding and adenoids. (Zähneknirschen und adenoiden vegetationen.) *Zeitsch. f. Ohrenheilk.*, p. 151, Vol. 73, No. 12, 1915.
- 35 BENJAMINS, C. E. Comparative observations on adenoids in the tropics and in Holland. (Vergleichende Notizen über adenoiden vegetationen in den Tropen und in Holland.) *Zeitsch. f. Ohrenheilk.*, p. 157, Vol. 73, No. 12, 1915.
- 36 BRADY, W. Unsuccessful adenoid operations. *N. Y. Med. Jour.*, July 17, 1915.
- 37 CHARLES, C. Post-operative care of tonsil and adenoid cases. *The Trained Nurse*, Feb., 1915.
- 38 GRAZIANI, V. Psychic disturbances following removal of adenoids. (Scomparsa di disturbi psichici successiva ad ablazione di vegetazioni adenoidi.) *Boll. delle malat. dell'orecchio*, p. 25, Vol. 33, No. 2, February, 1915.
- 39 MINCHEW, B. H. Adenoid tissue—Its relation to middle ear; consequences. *Ga. Med. Assn. Jour.*, March, 1915.
- 40 MOSSE, P. Adenoids and acetone vomit. (Vegetaciones adenoides y vomitos acetónicos.) *Arch. de rhinolog., laringolog. y otolog.*, p. 43, Vol. 26, No. 190, Jan., 1915.
- *41 PAGE, J. D. Adenoids in general practice. (Les végétations adénoïdes dans la pratique courante.) *Bulletin Medical*, Nov., 1915.
- 42 VICENTE, S. G. Etiology of adenoids. *Rev. de Med. y cir. pract.*, July 14, 1915.
- 43 WALDRON, C. W. Present status of adenoid and tonsil operations. *Dent. Cosmos*, June, 1915.

Neuroses.

- 44 BEATTIE, R. Headaches from pressure in the nose. *Det. Med. Jour.*, June, 1915.
- 45 BURCH, E. J. Curing hay-fever patients. *Med. Cent.*, June, 1915.
- †46 COOKE, R. A. The treatment of hay fever by active immunization. *THE LARYNGOSCOPE*, p. 108, Feb., 1915.
- 47 EMMERICH, R., and LOEW, O. Therapy of hay fever. (Weitere Mitteilungen über erfolgreiche Behandlungen des Heufiebers.) *Muench. med. Wchnschr.*, p. 41, Jan. 12, 1915.
- 48 GOODALE, J. L. Pollen therapy in hay fever. *Bost. Med. and Surg. Jour.*, July 8, 1915.
- *49 GOODALE, J. L. Pollen therapy in hay fever. *Ann. Otol., Rhinol. and Laryngol.*, p. 269, Vol. 24, No. 2, June, 1915.

- 50 HAYS, H. Experiments with autoserum in the treatment of hay fever. The relation of anaphylaxis and eosinophilia to hay fever. A general survey of new methods of treatment. *Ann. Otol., Rhinol. and Laryngol.*, p. 287, Vol. 24, No. 2, June, 1915.
- 51 ISKOWITZ, H. Multi-pollen vaccines in the treatment of hay fever. *Med. Rec.*, p. 270, Aug. 14, 1915.
- §52 KELLOGG, K. E. Sodium bicarbonate in hay fever. *N. Y. Med. Jour.*, p. 393, Aug. 21, 1915. *THE LARYNGOSCOPE*, p. 738, Oct., 1915.
- *53 MANNING, E. T. Hay fever—Its treatment by injections of a solution of ragweed pollen. *Jour. A. M. A.*, p. 655, Feb. 20, 1915.
- 54 MOLL, A. M. A case of hallucination of smell. *Medical Jour. of South Africa*, p. 4, Aug., 1915.
- 55 OPPENHEIMER, S., and GOTTLIEB, M. J. Active immunization of hay fever; a preliminary report. *N. Y. Med. Jour.*, p. 229, Feb. 6, 1915.
- *56 OPPENHEIMER, S., and GOTTLIEB, M. J. Pollinosis (hay fever); consideration of its treatment by active immunization. *N. Y. State Jour. Med.*, May, 1915; *Bull. Med. and Surg. Faculty of Md.*, July, 1915; *Maryland Med. Jour.*, Sept., 1915; and *N. J. Med. Soc. Jour.*, Sept., 1915.
- 57 RICE, G. B. Hay fever and its treatment. *Jour. Ophthalm., Otol. and Laryngol.*, p. 683, Aug., 1915.
- §58 SCHMIDT, K. Treatment of hay fever by roentgen rays. (Ein Versuch den Heuschnupfen durch Roentgenstrahlen zu beeinflussen.) *Munch. med. Woch.*, No. 23, June 8, 1915; *THE LARYNGOSCOPE*, p. 589, Aug., 1915.
- 59 SLUDER, G. The sympathetic syndrome (undescribed) of sphenopalatine or nasal ganglion neurosis, together with a consideration of previously described symptoms and treatment. *Lan. Clin.*, p. 457, April 24, 1915.
- 60 SORMANI, B. P. Vaccination against hay fever. *Nederl. Tijdschr. v. Geneesk.*, No. 23, Dec. 4, 1915.
- 61 WATTERS, W. H. Vaccine in hay fever. *Jour. Ophthalm., Otol. and Laryngol.*, p. 687, Aug., 1915.
- 62 WOOD, G. B. Treatment of hay fever. *Progressive Med.*, Vol. 17, No. 1, 1915.
- 63 WOOD, F. M. Hay fever and asthma. *Chicago Med. Recorder*, Aug., 1915.

Neoplasms (Nasal).

- 64 CALDERA, C. Diagnosis of malignant tumors; remarks on case of fibromyxoma of the nose. (La diagnosi clinica dei tumori maligni; considerazioni sopra un caso di fibromixoma del naso.) *Arch. ital. di Otol., Rinol. e Laringol.*, p. 490, Vol. 26, Fasc. 6, Dec., 1915.
- 65 CANFIELD, R. B. Cancer of the nose. *Jour. Mich. State Med. Soc.*, July, 1915.
- 66 CLERC, L. Hemorrhagic benign tumors of the nose. (Varieta di tumori benigni emorragici delle cavita nasali.) *Boll. delle malat. dell'orecchio*, p. 129, Vol. 33, No. 6, June, 1915.

- 67 DAVIS, E. D. Post-operative specimen of a pituitary cyst opened by the Killian-Hirsch operation. *Proc. Roy. Soc. Med., Otol. Sect.*, 57, No. 8, 1914.
- 68 FREUDENTHAL, W. Cancer of the upper air tract, with special reference to its treatment with radium. *N. Y. Med. Jour.*, July 3, 1915.
- 69 GAREL and PISTRE. Nasal angioma in a child three years old. (Volumineux angiome naso-génien à évolution rapide et fatale chez une enfant de trois ans.) *Rev. de Laryngol., d'Otol. e de Rhinol.*, p. 257, No. 42-43, Oct. 31, 1915.
- 70 MALAN, A. Adenocarcinoma of the nose. (Ipertrofia ghiandolari-adenomi-adenocarcinomi del naso.) *Arch. ital. di otol., rinol. e laringol.*, p. 69, Vol. 26, No. 1, Jan., 1915.
- 71 NATALE, P. Hemorrhagic polypoid tumor of the nasal cavity. (Sopra un tumore polipoide sanguinante delle cavità nasale.) *Arch. ital. di otol., rinol. e laringol.*, p. 265, Vol. 26, No. 4, Aug., 1915.
- 72 SPOHN, G. W. Nasal polypl. *Ind. State Med. Ass'n Jour.*, June, 1915.

Neoplasms (Naso-pharyngeal).

- *73 BASILE, G. Changes in the pituitary in a case of lymphosarcoma of the nasopharynx. (Histologische und funktionelle Veränderungen der zentralen Hypophyse des Menschen in einem Falle von Lymphosarkom des Nasenrachens.) *Zeitschr. f. Laryngol.*, p. 659, Bd. 7, H. 6, Sept., 1915; and *Riv. ital. Neuropatol., Psichiat. e Elettroterap.*, Vol. 8, No. 2, 1915.
- 74 BUTLER, R. History of a tumor of the pharynx eventually terminating in sarcoma. *Trans. Amer. Laryngol. Ass'n*, p. 275, 1915.
- 75 DABNEY, V. True myxoma of the rhinopharynx; report of two cases. *Trans. Amer. Laryngol. Ass'n*, p. 271, 1915; and *Ann. Otol., Rhinol. and Laryngol.*, p. 530, Sept., 1915.
- 76 DAVIS, E. D. A nasopharyngeal fibroma enucleated by a curved dissector and the finger. *Proc. Roy. Soc. Med., Laryngological Section*, Vol. 8, No. 5, March, 1915.
- 77 DELAVAN, D. B. The effects of radioactivity upon nasopharyngeal fibroma. *Med. Rec.*, June 26, 1915.
- 78 DIGHTON, A. Sarcoma of nose and nasopharynx. *Brit. Jour. of Surg.*, Jan., 1915.
- 79 GRAZIANI, V. Congenital polypus of the pharynx. *Arch. ital. di Laryngol.*, p. 1, Jan. 5, 1915.
- 80 GUTHRIE, T. The operative treatment of nasopharyngeal fibroma. *Jour. Laryngol., Rhinol. and Otol.*, p. 427, Vol. 30, No. 11, Nov., 1915.
- 81 LEVY, R. Nasopharyngeal polypus. *Colo. Med.*, March, 1915.
- 82 MOORE, J. L. I. Fibroma of nasopharynx. *Proc. Laryngol. Section, Royal Soc. Med.*, Vol. 8, No. 6, April, 1915.
- 83 MOORE, J. L. I. A nasopharyngeal polypus (choanal polypus) originating in the right maxillary antrum. *Proc. Royal Soc. Med., Laryngol. Section*, Vol. 9, No. 2, Dec., 1915.
- 84 MOORE, I. Nasopharyngeal fibroma removed through the mouth without any preliminary operation. *Proc. Roy. Soc. Med., Laryngol. Sect.*, p. 103, Vol. 8, No. 8, June, 1915.

- †85 ORTON, H. B. Spindle-cell sarcoma of the nasopharynx. *THE LARYNGOSCOPE*, p. 709, Oct., 1915.
- 86 PIERCE, NORVAL H. Choanal fibroma. *Ill. Med. Jour.*, May, 1915; *Ann. Otol., Rhinol. and Laryngol.*, p. 322, Vol. 24, No. 2, June, 1915.
- 87 SCHMITTEN, H. Tumors of the naso-pharynx. (Tumoren des Nasenrachenraumes.) *Inaug. Dissert. Bonn*, 1915.
- †88 SMITH, HARMON. Case of naso-pharyngeal sarcoma and two cases of naso-pharyngeal fibromata. *THE LARYNGOSCOPE*, p. 224, April, 1915.
- 89 SPICER, F. New growth in post-nasal space. *Proc. Roy. Soc. Med., Laryngol. Sect.*, p. 107, Vol. 8, No. 8, June, 1915.
- 90 SZMURLO, J. So-called naso-pharyngeal polypi and their treatment. *Ztschr. f. Laryngol.*, p. 473, Bd. 7, Heft. 4, 1915.

General.

- 91 ALBANNIS. Radium and mesothorium in cancer of the upper respiratory tract. *Bruns. Beitr.*, 92, 1915.
- 92 ARKIN, L. Case of autotransplantation of bone for nasal deformity due to syphilis. *Boston Med. and Surg. Jour.*, No. 18, May, 1915.
- 93 BARKER, C. B. Nasal obstruction in children. *Oklahoma State Med. Ass'n Jour.*, Vol. 2, No. 9, Feb., 1915.
- †94 BATROFF, W. C. Collapse of the alae nasi, its etiology and treatment. *THE LARYNGOSCOPE*, p. 72, Feb., 1915.
- 95 BECK, J. C. Local versus general anesthesia in nose and throat operations. *Ann. Otol., Rhinol. and Laryngol.*, p. 597, Sept., 1915.
- 96 BECKMAN, E. H. Correction of depressed fractures of the nose by transplant of cartilage. *Surg., Gyn. and Obst.*, p. 694, Dec., 1915.
- 97 BLAIR, E. G. Dactylocostal (osseous and cartilaginous) rhinoplasty. *Med. Herald*, p. 164, Vol. 34, May, 1915.
- §98 BLEGVAD, N. R. Treatment of ozena by injection of paraffin. *Ugeskr. f. Leger*, Jan. 28, 1915; *THE LARYNGOSCOPE*, p. 588, Aug., 1915.
- *99 BRUNNER, J., and JAKUBOWSKI, C. Treatment of scleroma of the upper air tract with auto-vaccines. *Arch. f. Laryngol.*, p. 282, Bd. 29, Heft. 2, 1915.
- †100 BRYANT, W. S. Transitional epithelium in the rhino-pharynx. *THE LARYNGOSCOPE*, p. 346, June, 1915.
- †101 CARTER, W. W. Cases of nasal deformity corrected by bone transplantation. *THE LARYNGOSCOPE*, p. 321, June, 1915.
- †102 CECIL, R. L. Streptococcus viridans in its relation to infections of upper respiratory tract. *THE LARYNGOSCOPE*, p. 97, Feb., 1915.
- 103 CERRY, G. Hygiene and therapy of catarrh in posterior naso-pharynx. *Amer. Med. Jour.*, Feb., 1915.
- 104 CHAPPELL, W. F. Effects of protein extracts, from fruits and pollen, on the upper air tract. *Trans. Amer. Laryngol. Ass'n*, p. 37, 1915.
- 105 CIAMPOLINI, ARNOLFO. Fracture of the nose. (Valutazione di danno lavorativo in caso di frattura delle ossa nasali.) *Boll. delle mal. dell'Orecchio, della Gola e del Naso*, Vol. 33, No. 7, July, 1915.
- 106 CLARK, J. S. Some experiences with the intra-nasal partial resection of the tear sac. *Jour. Ophthal. and Oto-Laryngol.*, p. 71, March, 1915; and *Ophthalmology*, April, 1915.

- 107 CLARK, J. S. West intranasal partial resection of tear sac for dacryocystitis, dacryostenosis, phlegmon or epiphora. *Ill. Med. Jour.*, Jan., 1915; *Ophthalmology*, No. 3, April, 1915.
- †108 COCKS, GERHARD H. Experimental studies of the effect of various atmospheric conditions upon the upper respiratory tract. *THE LARYNGOSCOPE*, p. 603, Sept., 1915.
- 109 COLEMAN, J. Nasal tuberculosis. *Med. Rec.*, p. 147, Jan. 23, 1915.
- 110 CORSON, G. R. S. Internal medication in the treatment of "catarrh" of the upper respiratory tract. *Penn. Med. Jour.*, p. 31, Oct., 1915.
- 111 DELAVAN, G. B. The neglect of injuries to the nose from football. *Med. Rec.*, p. 781, Nov. 6, 1915.
- 112 DEWEY, M. Cause of failure of some rhinological operations. *Jour. Ophth., Otol. and Laryngol.*, p. 291, April, 1915.
- 113 DIGHTON, A. Technique of analgesia in intranasal surgery. *Brit. Med. Jour.*, p. 13, July 3, 1915.
- 114 EITNER, E. Correction of saddle nose. (Sattelnasenkorrekturen.) *Deut. med. Woch.*, July 29, 1915.
- 115 FERGUSON, WM. Atrophic rhinitis. *N. Y. Med. Jour.*, p. 1091, Nov. 27, 1915.
- 116 FETTEROLF, G. Hemorrhage from nose and throat. *Penn. Med. Jour.*, Vol. 18, No. 10, July, 1915.
- 117 FREE, J. E. Nosebleed in measles. *Med. Fortnightly*, July 15, 1915.
- 118 FRITZSCHE, ARTHUR. Malformations of the nose and their correction. (Nasendeformationen und ihre Korrekturen.) *Inaug. Dissert.*, Würzburg, July, 1915.
- 119 FRÜHWALD. Gunshot wounds in relation to the upper air passages. *Berl. klin. Woch.*, p. 142, No. 6, 1915.
- 120 GERBER, PROF. Gunshot wounds of the upper air passages. (Über Schussverletzungen der oberen Luftwege und benachbarter Teile.) *Arch. f. Laryngol. u. Rhinol.*, p. 331, Vol. 29, No. 3, 1915.
- 121 GIFFORD, H. Method of destroying lacrimal sac in chronic dacryocystitis. *Ophthal. Record*, Jan., 1915.
- †122 GLOGAU, O. A case of dacryocysto-rhinostomy. *THE LARYNGOSCOPE*, p. 28, Jan., 1915.
- 123 GOUX, L. J. Rhinitis—acute and chronic. *Mich. State Med. Soc. Jour.*, Jan., 1915.
- §124 GRAHAM, C. I. Tuberculosis of the nasal fossae. *Proc. Roy. Soc. Med., Laryngol. Sect.*, Vol. 8, No. 5, March, 1915; *THE LARYNGOSCOPE*, p. 56, Jan., 1916.
- 125 GRAYSON, C. P. Syphilis of the nose and throat. *Penn. Med. Jour.*, p. 100, Nov., 1915.
- 126 GRIFFIN, E. H. How to blow the nose. *Med. Record*, p. 1007, Dec., 1915.
- 127 GUGGENHEIM, L. K. Cocco-bacillus fetidus ozenae Perez vaccine in treatment of ozena. *Interstate Med. Jour.*, p. 129, Feb., 1915.
- †128 GUNDELACH, C. A. Posterior nasal operation by means of the nasopharyngoscope. *THE LARYNGOSCOPE*, p. 83, Feb., 1915.
- 129 GUTTMAN, J. Radium in rhinoscleroma. *N. Y. Med. Jour.*, p. 711, Oct. 2, 1915.
- 130 HAGERMANN, J. A. The correlation between olfactory and genital functions in the human female. *Med. Record*, p. 1091, Dec. 25, 1915.

- †131 HANGAR, F. M. An intra-nasal operation with a guide for the cure of dacryocystitis. *THE LARYNGOSCOPE*, p. 23, Jan., 1915.
- 132 HARRY, P. A. Hygiene of the nose and throat. *Can. Pract. and Rev.*, p. 340, July, 1915.
- 133 HARTMANN, A. Inhalation of ammonium chlorid in catarrhal affections. *Deut. mediz. Woch.*, Vol. 41, No. 30, July 22, 1915.
- 134 HEKTOEN, L., and RAPPAPORT, B. The use of kaolin to remove bacteria from the nose and throat. *Jour. A. M. A.*, June 12, 1915.
- 135 HINNEN, G. A. Case of epistaxis of syphilitic origin. *Lan.-Clin.*, Vol. 114, No. 5, July 31, 1915.
- 136 HOLLAENDER, E. Rhinoplasty from the skin of the sternum. *Urologic and Cutaneous Review*, Tech. Supp., 1914.
- *137 HORN, HENRY. Etiology and treatment of ozena. *Jour. A. M. A.*, p. 788, Aug. 28, 1915.
- 138 HOWE, A. C. Relation of nasal symptoms to endamebic infections of the nose and mouth. *Med. Record*, p. 604, Oct. 9, 1915.
- *139 HOWELL, K. Complement-fixation in acute rhinitis. *Jour. Infec. Dis.*, No. 3, May, 1915.
- 140 INGRAM, L. C. Progress of intranasal surgery. *Florida Med. Ass'n Jour.*, Nov., 1915.
- †141 ISRAEL, SIDNEY. Texas screw-worm infection of the entire nose and throat, including the accessory sinuses. *THE LARYNGOSCOPE*, p. 657, Sept., 1915.
- 142 JASTER, C. O. Nasal obstruction and its treatment. *Ohio State Jour. Med.*, Dec., 1915.
- *143 JOHNSTON, R. H. Total rhinoplasty; a case report. *Am. Jour. of Surg.*, p. 149, April, 1915.
- 144 KAFEMANN, R. Gunshot wounds of the upper respiratory tract. *Deut. med. Woch.*, No. 17, 1915.
- 145 KAHN, ALFRED. Notes on a case of naso-lachrymal disease. *Interstate Med. Jour.*, p. 901, Vol. 22, No. 9, Sept., 1915.
- 146 KASSEL, K. A case of rhinitis vasomotoria produced by the spool-worm. (Ein Fall von Rhinitis vasomotoria verursacht durch Spulwurm.) *Ztschr. f. Laryngol. u. Rhinol.*, Bd. 7, H. 5, April, 1915.
- 147 KLENK, CHAS. L. The treatment of ozena and asthma with autogenous vaccines. *Interstate Med. Jour.*, p. 1209, Dec., 1915.
- 148 KÖNIG, F. Restoration of nose. *Beitr. z. klin. Chir.*, Vol. 94, No. 3, 1915.
- 149 KREBS, G. Gunshot wounds of the face involving the nasal cavity. (Bemerkungen zu den Gesichtsschüssen mit Beteiligung der Nasenhöhle.) *Munch. mediz. Woch.*, p. 1203, No. 35, Aug. 31, 1915.
- 150 KRUSE. Cause of rhinitis. (Erreger von Husten und Schnupfen.) *Deut. med. Woch.*, No. 5, 1915.
- 151 LUBMAN, M. Epistaxis. *Medical Times*, p. 389, Dec., 1915.
- 152 LUZZATI, DR. Early nasal syphilis. (Sopra un caso di terziarismo nasale precoce.) *Arch. ital. di otol., rinol. e laringol.*, p. 29, Vol. 26, No. 1, Jan., 1915.
- 153 MARINHO, J. Naso-pharyngitis as a cause of protracted fever in infants. *Brazil-Medico*, No. 9, March 1, 1915.
- *154 MAYER, E. The early recognition of cancer of the upper air passages. *American Jour. Surg.*, July, 1915.

- 155 MAZZA, S. Vaccine therapy of ozena. *Wien. klin. Woch.*, No. 18, May 6, 1915.
- 156 MAZZETTI, L. Meningococci in the noses of horses. (Il meningococco nel muco nasale dei cavalli durante un epidemia di meningite cerebro-spinale.) *Rif. Med.*, May 29, 1915.
- 157 MCGOWAN, J. P. On the Perez bacillus and the alleged relationship of ozena to a certain condition in the dog and more especially to distemper. *Jour. of Laryngol.*, p. 57, Feb., 1915.
- 158 MCINTOSH, J., and BULLOCK, W. E. Recognition and isolation of meningococcus in naso-pharynx of cerebrospinal fever patients. *Lancet*, Nov. 27, 1915.
- 159 MENZEL, K. M. Occupational diseases of the mucous membrane of the upper air passages in bakers. (Berufliche Erkrankungen an der Schleimhaut der oberen Luftwege der Bäcker.) *Arch. f. Laryngol. u. Rhinol.*, p. 394, Vol. 29, No. 3, 1915.
- 160 MILLIGAN, W. Photographs of a case of chronic lymphangitis of the tissues covering the nose; hints regarding treatment. *Proc. Roy. Soc. Med., Laryngological Sections*, Vol. 8, No. 5, March, 1915.
- *161 MILLIGAN, W. Rhinophyma; its etiology, pathology and treatment. *Lancet*, Sept. 18, 1915.
- 162 MORESTIN, H. Extraction of ball fixed in the nasal fossae. (Extraction d'une balle fixée dans la cloison osseuse des fosses nasales.) *Soc. de chir. de Paris*, Feb. 10, 1915; *Presse Med.*, Feb. 18, 1915.
- †163 MOSHER, H. P. An operation for draining the lacrimal sac and the nasal duct into the unciform fossa. *THE LARYNGOSCOPE*, p. 739, Nov., 1915.
- †164 MURRAY, W. R., and LARSON, W. P. The bacillus of Perez as an etiological factor in ozena. *THE LARYNGOSCOPE*, p. 763, Nov., 1915.
- 165 O'CONNELL, G. A. Intranasal obstructions. *South. Med. Jour.*, p. 531, Vol. 8, June, 1915; *Jour. Ophthal. and Oto-Laryngol.*, p. 316, Vol. 9, No. 10, Oct., 1915.
- 166 O'MALLEY, J. F. Two cases of intra-nasal adhesions. *Proc. Roy. Soc. Med., Laryngol. Sect.*, p. 129, Vol. 8, No. 8, June, 1915.
- 167 ODELL, A. Bacteriology and bacterio-therapy of upper air passages. *Mich. State Med. Soc. Jour.*, Jan., 1915.
- 168 ONODI, A. Eye affections of nasal origin. (Über die Lehre von den Augenleiden nasalen Ursprungs.) *Arch. f. Laryngol. u. Rhinol.*, p. 430, Vol. 29, No. 3, 1915.
- 169 O'REILLY, C. A. Intra-nasal treatment of dysmenorrhea. *Amer. Jour. Obst. and Dis. Women and Children*, Oct., 1915.
- 170 PAGET, OWEN. Nasal props and an operation for the incapable nose. *Med. Rev. Reviews*, p. 532, Sept., 1915.
- 171 PEGLER, L. H. Case of synechiae and contraction of the vestibules. *Proc. Laryngol. Section Royal Soc. Med.*, Vol. 3, No. 6, April, 1915.
- *172 PEGLER, L. H. Headache associated with intra-nasal disorders. *Lancet*, Feb. 27, 1915.
- 173 PERKINS, C. E. Epistaxis in arteriosclerosis. *N. Y. Med. Jour.*, May 15, 1915.
- 174 PFISTER, F. Plea for corrective and cosmetic surgery of the nose. *Wisconsin Med. Jour.*, June, 1915.

- 175 PILCHER, J. S. The absorption of drugs from the nasal mucosa of dogs. *Jour. A. M. A.*, page 232, July 17, 1915.
- 176 POLLOCK, H. L. Nerve-blocking; injection of the spheno-palatine ganglion in some of the commoner diseases of the nose. *Ill. Med. Jour.*, May, 1915.
- 177 PONT, A. Improved technic for making artificial nose. *Lyon Med.*, Vol. 46, Nos. 32-35, 1915.
- 178 PRINCE, A. E. Operation for direct drainage of lachrymal sac into middle meatus from the standpoint of the ophthalmologist. *Ophthal. Rec.*, Vol. 24, No. 8, Aug., 1915.
- 179 PULLEINE, R. Notes on the etiology and treatment of catarrh of the nose and pharynx. *Med. Jour. Australia*, Sept. 25, 1915.
- 180 QUICKEL, T. C. Obstructions of nose and throat. *S. Carolina Med. Ass'n Jour.*, Jan., 1915.
- 181 RETHI, L. Relations between ozena and disease of the nasal accessory sinuses. *Wiener klin. Woch.*, Nov. 7, 1915.
- 182 ROE, J. O. Nasal deformity as a cause of nasal obstruction corrected by the subcutaneous method. *Trans. Amer. Laryngol. Ass'n*, p. 285, 1915.
- 183 ROSENTHAL, H. Treatment of rhinitis. *Deut. med. Woch.*, No. 13, 1915.
- 184 ROY, J. N. Ozena among different races. (L'ozène chez les différentes races de la terre.) *Ann. des mal. de l'oreille*, p. 733, Vol. 40, No. 8, 1914-1915; *Rev. de Laryngol.*, p. 238, No. 40-41, Oct. 15, 1915.
- 185 SEIDEL, O. Gunshot wounds of the nose. *Corresp. Blatt d. allg. arztl. Ver. von Thüringen*, April 3, 1915.
- 186 STEINTHAL, Restoration of nose by Italian method. *Beitr. z. klin. Chir.*, Vol. 94, No. 2, 1915.
- 187 STENGER, LUDWIG. Nasal hemorrhage with special reference to habitual epistaxis. (Blutungen aus der Nase mit besonderer Berücksichtigung der habituellen Epistaxis.) *Inaug. Dissert.*, Würzburg, June, 1915.
- 188 STOUT, G. C. Headaches due to non-suppurative intra-nasal conditions. *Trans. Amer. Laryngol. Ass'n*, p. 206, 1915.
- 189 TEETS, C. E. Euonymin in the treatment of diseased conditions of the nose and throat. *Jour. Ophthal., Otol. and Laryngol.*, June, 1915.
- †190 THOMASSON, WM. J. Congenital bony occlusion of the right nasal choana. *THE LARYNGOSCOPE*, p. 221, April, 1915.
- 191 TOP, H. Case of congenital fistula of the nose. *Proc. Royal Soc. Med., Laryngol. Section*, Vol. 9, No. 2, Dec., 1915.
- 192 TORRIGIANI, C. A. Hypophyseal syndrome of inflammation of the adnexal glands of the nasal fossae. (Sindrome ipofisaria da infiammazione delle glandole annesse alle fosse nasali.) *Arch. ital. di otol., rinol. e laringol.*, p. 283, Vol. 26, No. 4, Aug., 1915.
- 193 TORRINI, U. L. Endonasal tuberculosis. (Contributo all conoscenza della tubercolosi endonasale. Granuloma tubercolare del cornetto medio.) *Arch. ital. di otol., rinol. e laringol.*, p. 14, Vol. 26, No. 1, Jan., 1915.
- 194 TOUSEY, S. X-ray prevention of nasal diseases. *N. Y. Med. Jour.*, p. 506, March 13, 1915.

- 195 TUNNICLIFF, R. Bacteriology of rhinitis with special reference to anærobic organism (bacillus rhinitis). *Jour. Infec. Dis.*, No. 3, May, 1915.
- 196 VITTO-MASSEL, R. Treatment of rhinoscleroma with autogenous vaccines. (Breve nota sulla cura del rinoscleroma con l'autovaccino.) *Arch. ital. di Laringol.*, p. 159, Vol. 35, No. 4, Oct., 1915.
- 197 WEIH, E. P. Novocain as used in simpler nose and throat operations. *Iowa St. Med. Soc. Jour.*, June, 1915.
- 198 WEINSTEIN, J. Clinical report of the successful use of emetine for the control of hemorrhage following naso-pharyngeal operations. *Med. Record*, Jan. 16, 1915.
- 199 WEISS, M. J. Chronic rhinitis; its treatment. *N. J. Med. Soc. Jour.*, Dec., 1915.
- 200 WIENBERG, S. W. Epistaxis and its treatment. *Therapeutic Gazette*, Nov., 1915.
- 201 WILBUR, E. P. Intra-nasal and pharyngeal infections in their relation to eye and ear. *Mich. State Med. Soc. Jour.*, Jan., 1915.
- 202 WILSON, WM. Technique of analgesia in intra-nasal surgery. *Brit. Med. Jour.*, p. 1083, June 26, 1915.
- 203 WINSLOW, J. R. Obstruction of the posterior nasal orifices (choanae). *Maryland Med. Jour.*, p. 295, Dec., 1915.
- 204 YOUNG, E. W. General systemic disturbances and catarrhal conditions of the nose and throat. *Jour. Oph., Oto-Laryngol.*, May, 1915.
- *205 ZANGE, J. Gonorrheal infection of the upper respiratory passages in adults. (Gonorrhoeische Infektion der oberen Luftwege beim Erwachsenen.) *Zeitschr. f. Ohrenheilk.*, p. 165, Bd. 73, H. 3, Nov., 1915.
- 206 ZBINDEN, T. Brief review of the pathology and bacteriology of naso-pharyngeal infections with some suggestions for effective vaccine therapy. *Jour. of Ophth., Otol. and Laryngol.*, p. 202, March, 1915.

II. MOUTH AND PHARYNX.

Palate.

- 207 BADGEROW, G. Tumor of palate. *Proc. Royal Soc. Med., Laryngol. Section*, Vol. 9, No. 2, Dec., 1915.
- 208 BERRY, J. Surgery of cleft palate. *Surg., Gynecol. and Obstetr.*, p. 85, Jan., 1915.
- 209 BJÖRLING, E. Primary syphilitic chancre of the hard palate. *Dermatolog. Zeitschr.*, Vol. 12, No. 4, April, 1915.
- §210 BLAKEWAY, H. Operative treatment of cleft palate. *Lancet*, p. 533, March 13, 1915; *THE LARYNGOSCOPE*, p. 866, Dec., 1915.
- 211 BLAKEWAY, H., LANE, A., JAMES, W., and KELLOCK, T. H. Effects of cleft-palate operations on the dental arch. *Practitioner*, Aug., 1915.
- *212 BROPHY, T. W. Late results of cleft-palate operations. *Surg., Gynecol. and Obstetr.*, p. 98, Jan., 1915.
- 213 BROWN, G. V. I. Principles which govern the ultimate results of harellip and cleft palate. *Surg., Gynecol. and Obstetr.*, p. 87, Jan., 1915.

- 214 EASTMAN, J. R. Factors of safety in cleft-palate surgery. *Surg., Gynecol. and Obstetr.*, p. 91, Jan., 1915.
- 215 EASTMAN, JOSEPH R. The surgical anatomy of cleft palate, *Jour. A. M. A.*, p. 915, Sept. 11, 1915.
- 216 EASTMAN, J. R. Principles of harelip and cleft-palate surgery. *Lancet-Clinic*, Nov. 28, 1915, and *Louisville Monthly Jour. Med. and Surg.*, p. 204, Dec., 1915.
- §217 EMERSON, L. Operations for clefts of the hard and soft palate. *Jour. A. M. A.*, p. 301, Jan. 23, 1915; *THE LARYNGOSCOPE*, p. 866, Dec., 1915.
- 218 FRÜHWALD, R. Syphilis of the nose and palate. (Zwei Fälle von ungewöhnlich schwerer Syphilis der Nase und des Gaumens.) *Dermat. Woch.*, No. 1, 1915.
- 219 GOYDER, F. W. On the treatment of cleft palate. *Surg., Gynecol. and Obstetr.*, p. 95, Jan., 1915.
- 220 HYBBINETTE, S. Treatment of cleft palate. (Zur Frage von der Behandlung angeb. Gaumendefekte.) *Nord. mediz. Archiv.*, Stockholm, Vol. 47, No. 3, 1915.
- 221 IVY, R. H. Endothelioma of the soft palate. *N. Y. Med. Jour.*, p. 193, Jan. 30, 1915.
- 222 MACKENTY, J. E. Operative treatment of cleft palate. *Ann. Otol., Rhinol. and Laryngol.*, March, 1915.
- 223 MORONE, G. Mixed tumors of the palate. *Policlinico*, Sept. 15, 1915, Oct. 10, 1915, and Nov. 21, 1915.
- 224 MORONE, G. Tumors of the palate. *Policlinico*, No. 12, Dec., 1915.
- †225 ROY, DUNBAR. Partial paralysis of the soft palate following removal of tonsil and adenoids. *THE LARYNGOSCOPE*, p. 361, June, 1915.
- 226 SMITH, E. T. Cleft palate and harelip. *Med. Century*, Feb., 1915.
- 227 SMITH, E. T. A bloodless method of enucleating tonsils. *Trans. Amer. Oto. Soc.*, p. 489, Vol. 8, Part 3, 1915.
- 228 THOMPSON, G. S. Nasal flap and modified Langenbeck operation for cleft palate. *Med. Jour. of Australia*, May 22, 1915, and *Lancet*, June 19, 1915.

Tonsils.

- 229 ADAMS, CHARLES FRANKLIN. Tonsilloadenectomy and the control of the hemorrhage. *New York Medical Journal*, May 1, 1915.
- 230 AGNEW, F. F. Hemorrhage following removal of the faucial tonsil. *Ann. Otol.*, p. 44, March, 1915.
- 231 ALEXANDER, G. J. Acute leukemia as first manifested in the tonsils. *Jour. Ophthal., Otol. and Laryngol.*, p. 780, Sept., 1915.
- *232 ANDERSON, H. B. Appendicitis as a sequela of tonsillitis. *Amer. Jour. Med. Sc.*, p. 541, Oct., 1915.
- 233 ASHCRAFT, L. T. The tonsil from the standpoint of the urologist. *Jour. Ophthal., Otol. and Laryngol.*, July, 1915.
- 234 BALFOUR, DONALD C. Tonsillectomy in children. *Annals of Surgery* March, 1915.
- 235 BALFOUR, D. C. Tonsillectomy in children from standpoint of a general surgeon. *Ann. of Surg.*, p. 257, March, 1915.
- 236 BARNES, H. A. Radical treatment of peritonsillar abscess by tonsillectomy during the acute stage of the disease. *Bost. Med. and Surg. Jour.*, Dec. 30, 1915.

- †237 BLUM, SANFORD. Theory of tonsillectomy; results in pediatrics. *THE LARYNGOSCOPE*, p. 661, Sept., 1915.
- *238 BLUM, S. The proper position of tonsillectomy in pediatrics. *Arch. Pediatric*, Nov., 1915.
- *239 BLUM, S. Tonsils as excretory organs for cervical glands. *Arch. Pediatrics*, Nov., 1915.
- 240 BOND, J. W. Encapsulated tumor removed from region of left tonsil, soft palate, etc. *Proc. Royal Soc. Med., Laryngol. Section*, Vol. 9, No. 2, Dec., 1915.
- 241 BRAWLEY, F. Focal infection with special reference to the tonsil. *Ill. Med. Jour.*, Sept., 1915.
- 242 BRAY, H. A. Milk-borne epidemic of tonsillitis in tuberculous patients. *Jour. A. M. A.*, p. 1127, April 3, 1915.
- 243 BULATNIKOW, T. J. Regio-latero-pharyngea. Topography of its arteries in relation to question of dangerous hemorrhage in operative procedures in the tonsillar regions. *Arch. f. Laryngol.*, p. 225, Bd. 29, Heft 2, 1915.
- 244 BURACK, S. M. Tonsil enucleation. (Zur Frage von der Enukleation der Tonsillen.) *Zeitschr. f. Laryngol.*, p. 673, Vol. 7, H. 6, Sept., 1915.
- 245 CARMODY, T. E., and FINNOFF, W. C. Death from hemorrhage following peritonsillar abscess. *Colo. Med.*, Vol. 12, No. 9, Sept., 1915.
- †246 CARMODY, T. E. The histopathology of the faucial tonsil. *THE LARYNGOSCOPE*, p. 272, May, 1915.
- 247 CASTELLANI, A. Mycotic acute and subacute tonsillitis. (Tonsilliti acute e subacute di origine iformicetica.) *Pediatria*, Sept., 1915.
- 248 CHARLES, C. Post-operative care of tonsils and adenoid cases. *The Trained Nurse*, Feb., 1915.
- 249 CLAY, J. V. F. The faucial tonsils—atomy. *Jour. Ophthal., Otol. and Laryngol.*, July, 1915.
- 250 CLAY, J. F. V., and PALEN, G. J. The faucial tonsil—focal infection. *Jour. Ophthal., Otol. and Laryngol.*, July, 1915.
- 251 CLEVINGER, W. F. Tonsil and adenoid pathology. *Dent. Summary*, Oct., 1915.
- 252 COMBY, J. Opening retropharyngeal and tonsillar abscesses without the use of the knife. *Bull. et Mem. Soc. med. d. hop. de Paris*, p. 823, Vol. 31, 1915.
- 253 DANCEY, A. B. Indications for removal of faucial tonsils. *Tenn. State Med. Ass'n*, Jan., 1915.
- 254 DENMAN, I. O., and MCKESSON, E. I. Tonsillectomy: the forward inclined sitting posture under nitrous oxid-oxygen anesthesia, and other features. *Amer. Jour. Surg.*, Vol. 29, May, 1915.
- 255 DILLINGHAM, W. B. Chronic infection of the tonsils. *Med. Summary*, June, 1915.
- 256 DORSEY, R. T. Infected tonsils. *Ga. Med. Ass'n Jour.*, Aug., 1915.
- 257 ECCLES, R. G. The tonsil and the struggle for existence. *Med. Rec.*, July 10, 1915.
- *258 FARMACHIDIS, C. B., and VATTUONE, A. Effect of tonsil extract on the blood picture. (Influenza dell'estratto di tonsille palatine sulla crasi sanguigna.) *Policlinico*, No. 3, March, 1915.

- 259 FARMACHIDIS, G. B. Tonsil extract useful in glycosuria and diabetes. (L'opoterapia delle glicosurie semplici e del diabete mellito, mediante la somministrazione di estratto di tonsille palatine.) *Rif. med.*, Vol. 31, No. 29, July 17, 1915.
- 260 FERRER, J. A. The surgical tonsil. *Med. Century*, April, 1915.
- 261 FISHER, A. G. Sigmoid tortuosity of the internal carotid artery and its relation to the tonsil and pharynx. *Lancet*, July 17, 1915.
- 262 FLETCHER, J. R. The standard tonsillectomy. *Ann. Otol., Rhinol. and Laryngol.*, p. 591, Sept., 1915.
- *263 GARDINER, H. The tonsils and chronic cervical adenitis. *Lancet*, Oct. 2, 1915.
- 264 GILMER, T. L. The significance to the dentist of oral infections in their relation to systemic disorders. *Dental Review*, p. 1029, Oct., 1915.
- 265 GRAHAM, C. I. Epithelioma of the left tonsil and adjacent part of tongue after operation (specimen shown). *Proc. Roy. Soc. Med., Laryngological Section*, Vol. 8, No. 5, March, 1915.
- 266 GRIFFIN, E. H. Quinsy or peritonsillar abscess. *Med. Rec.*, p. 224, Feb. 6, 1915.
- 267 GÜTTICH, A. So-called capsule of palatal tonsil. *Ztschr. f. Laryngol.*, p. 497, Bd. 7, Heft 4, 1915.
- 268 GUTHRIE, D. Practical hints on removal of tonsils and adenoids. *Practitioner*, Nov., 1915.
- 269 HALSTED, T. H. The tonsil in its relation to rheumatic infections. *N. Y. State Jour. Med.*, Nov., 1915.
- 270 HARMON, G. O. Concerning the tonsils. *Med. Summary*, Jan., 1915.
- 271 HARRISON, W. G. Tonsillectomy during acute endocarditis; report of cases. *South. Med. Jour.*, Jan., 1915.
- 272 HARTLEY, A. Anesthesia in tonsil operations. *Jour. Ophthalm., Otol. and Laryngol.*, July, 1915.
- 273 HASELTINE, B. Tonsil surgery and voice function. *Jour. Ophthalm., Otol. and Laryngol.*, July, 1915.
- *274 HERRMAN, C. Tonsillar manifestations in the early diagnosis of measles. *Amer. Jour. Dis. Child.*, Oct., 1915.
- 275 HESTER, J. H. Complete and incomplete tonsillectomy. *Kentucky Med. Jour.*, Sept., 1915.
- 276 HIGGINBOTHAM, T. L. Can mortality of tonsillitis be reduced? *South-west Jour. Med. and Surg.*, Dec., 1915.
- *277 HORGAN, J. B. Tonsillectomy by blunt dissection under local anesthesia. *Jour. Laryngol., Rhinol. and Otol.*, p. 482, Dec., 1915.
- 278 HORNE, W. J. Epithelioma of the tonsil. *Proc. Roy. Soc. Med., Laryngological Section*, Vol. 8, No. 5, March, 1915.
- 279 JERVEY, J. W. Vascular ligation in the tonsillar fossa. *South. Med. Jour.*, May, 1915.
- 280 JERVEY, J. W. The control of hemorrhage in tonsil enucleation. *Jour. So. Carolina Med. Ass'n*, Oct., 1915.
- 281 JOHNSON, O. H. Tonsillectomy. *Florida Med. Ass'n Jour.*, Nov., 1915.
- 282 JOHNSTON, ANNA. Tonsillitis and its sequelae. *Hahneman Month.*, Nov., 1915.
- 283 JOHNSTON, W. Relation of tonsils to rheumatism. *Northwest Med.*, Oct., 1915.

- 284 KNACK, A. V. Tonsillar abscess in diphtheria. *Zeitschr. f. Hyg. und Infektionskrankh.*, p. 163, Vol. 80, 1915.
- †285 LA MOTTE, W. O. A case of abnormal bone in the throat. *THE LARYNGOSCOPE*, p. 776, Nov., 1915.
- 286 LEOPOLD, R. S. The faucial tonsils—pathology and bacteriology. *Jour. Ophthal., Otol. and Laryngol.*, July, 1915.
- 287 LESCHKE, E. Pneumococcus tonsillitis. *Muench. med. Wchnschr.*, p. 2433, Dec. 29, 1914.
- †288 MAKUEN, G. HUDSON. The surgical anatomy of the so-called capsule of the faucial tonsil. *THE LARYNGOSCOPE*, p. 685, Oct., 1915.
- *289 MARQUIS, G. P. The post-operative antiseptic treatment of the tonsillar fossae. *Jour. A. M. A.*, July 10, 1915.
- 290 MARTIN, H. H. Sluder tonsil operation. *Ga. Med. Ass'n Jour.*, Jan., 1915.
- 291 McLEARY, J. R. Tonsillectomy and its effect upon the voice. *Clinique*, p. 123, March, 1915.
- 292 McMURRAY, J. B. Primary carcinoma of tonsil; report of cases. *Penn. Med. Jour.*, Vol. 18, No. 10, July, 1915.
- 293 McREYNOLDS, J. O. Present status of tonsil surgery. *Southwest Jour. Med. and Surg.*, Oct., 1915.
- 294 MIDDLETON, A. B. Round-worm within the body of a tonsil. *Jour. A. M. A.*, p. 659, Feb. 20, 1915.
- 295 MUHLY, E. G. The faucial tonsils—histology. *Jour. Ophthal., Otol. and Laryngol.*, July, 1915.
- 296 MURPHY, F. G. Circumcision of the tonsil. *Jour. Iowa State Med. Soc.*, p. 328, Vol. 5, No. 8, Aug. 15, 1915.
- 297 MURRAY, G. D. Study of 848 cases of tonsillectomy. *Penn. Med. Jour.*, Vol. 18, No. 11, Aug., 1915.
- 298 MYERS, D. W. The faucial tonsil—treatment and materia medica. *Jour. Ophthal., Otol. and Laryngol.*, July, 1915.
- 299 OFTEDAL, S. Some clinical manifestations of focal infections. *Jour. A. M. A.*, p. 1911, Nov. 27, 1915.
- 300 PACKARD, F. R. Indications for the removal of tonsils and adenoids in children. *Therapeutic Gazette*, Feb. 15, 1915.
- 301 PALEN, G. J. The tonsil operation. *Jour. Ophthal., Otol. and Laryngol.*, July, 1915.
- 302 PARKER. The tonsils in relation to systemic diseases. *Jour.-Lan.*, April 1, 1915.
- 303 PARSONS, J. G. Lingual tonsillitis. *Jour. Ophthal. and Oto-Laryngol.*, p. 253, Vol. 9, No. 8, Aug., 1915.
- 304 PÄSSLER, H. Chronic infections in tonsils and nasal sinuses. (Die chronischen Infektionen im Bereiche der Mundhöhle und der Krieg.) *Therap. d. Gegenwart*, Oct., 1915.
- 305 PATTEE, J. J. Anatomic, clinical and surgical consideration of tonsil. *N. Mex. Med. Jour.*, March, 1915.
- 306 PURCELL, C. E. Diseased tonsils. What shall we do with them? *Kentucky Med. Jour.*, Dec., 1915.
- 307 PYBUS, FREDERICK C. Some infections of the tonsils. *Brit. Med. Jour.*, May 22, 1915.

- 308 RICHARDS, G. L. Relation of the tonsils, adenoids and other throat conditions to tuberculous cervical adenitis. *Bost. Med. and Surg. Jour.*, 172, 1915.
- †309 RICHARDSON, C. W. Tonsillectomy in the adult. Are we justified in doing so many indiscriminate tonsillectomies for remote infections? *THE LARYNGOSCOPE*, p. 293, May, 1915.
- 310 ROBINSON, J. M. Mutilation of tonsil. *Kans. Med. Soc. Jour.*, Jan., 1915.
- *311 ROMAN, D. The relation of the tonsil to the thyroid gland. *Jour. Ophthal., Otol. and Laryngol.*, July, 1915.
- 312 ROWLAND, W. D. Tonsillar infection. *Jour. of Ophth., Otol. and Laryngol.*, p. 213, March, 1915.
- 313 RYLAND, A. Acute ulcerative tonsillitis associated with virulent pyorrhea alveolaris. *Jour. Laryngol., Rhinol. and Otol.*, p. 344, Vol. 30, No. 9, Sept., 1915.
- 314 SAMUELS, S. M. Tonsillectomy and illustrations of operation with a special knife. *Northwest Med.*, Oct., 1915.
- 315 SANGER, F. D. Some phases of the tonsil question. *South. Med. Jour.*, Vol. 8, No. 8, Aug., 1915.
- 316 SAVAGE, M. M. Systemic infections for which the tonsil is held responsible, and control of hemorrhage during tonsillectomy. *Id. Med. Jour.*, p. 27, Feb., 1915.
- †317 SCHOOLMAN, N. Report of a case showing the bi-polar origin of the faucial tonsil. *THE LARYNGOSCOPE*, p. 338, June, 1915.
- 318 SMITH, C. H. Tonsillectomy. *N. Y. Med. Jour.*, p. 353, Feb. 20, 1915.
- *319 SMITH, C. M. Primary syphilis of the tonsil. *Jour. Cutaneous Dis.*, Oct., 1915.
- 320 SMITH, T. and BROWN, J. H. Study of streptococci isolated from certain presumably milk-borne epidemics of tonsillitis occurring in Massachusetts in 1913 and 1914. *Jour. of Med. Research*, Jan., 1915.
- 321 STATLER, W. K. Tonsillitis. *Jour. Mo. State Med. Ass'n*, p. 114, March, 1915.
- 322 STREET, R. H. Tonsillotomy versus tonsillectomy. *Jour. Ophth. Otol. and Laryngol.*, p. 140, Feb., 1915.
- 323 STRIETMANN, W. H. Endocarditis with streptococcemia. Infection from tonsils. *Cal. State Jour. of Med.*, March, 1915.
- §324 SYME, W. S. Abnormally long styloid process causing throat symptoms; removal. *Jour. Laryngol., Rhinol. and Otol.* Vol. 30, No. 8, p. 303, Aug., 1915. *THE LARYNGOSCOPE*, p. 776, No., 1915.
- 325 TRAUTMANN, G. Capsule and neighboring fascia of the tonsil (Über die Kapsel und die benachbarten Faszien der Tonsille). *Zeitschr. f. Laryngol.*, p. 691, Vol. 7, H. 6, Sept., 1915.
- 326 TURNER, A. LOGAN. Fibro-sarcoma of the oro-pharynx. *Edinburgh Med. Jour.*, p. 261, Oct., 1915.
- 327 TURNER, A. LOGGAN. Retropharyngeal abscess. *Edinburgh, Med. Jour.*, p. 261, Oct., 1915.
- 328 URBANTSCHITSCH, E. Tonsillitis keratosa punctata. *Wiener. Klin. Woch.*, No. 11, 1915.

- †329 VANDERHOOF, D. A. Abscess following tonsillectomy under local anesthesia. *THE LARYNGOSCOPE*, p. 20, Jan., 1915.
- 330 VARNER, ANNA D. Medical treatment of hypertrophied tonsils. *Hahneman Month.*, Nov., 1915.
- 331 VOORHEES, I. W. Tonsil operations with especial reference to singers. *Am. Medicine*, p. 168, March, 1915.
- 332 WEAVER, J. S. Some factors in the removal of tonsils. *Jour. Mo. State Med. Assn.*, July, 1915.
- 333 WIDMAN, F. H. The faucial tonsils—embryology and physiology. *Jour. Ophthal., Otol. and Laryngol.*, July, 1915.
- 334 WILLIAMS, C. Plea for conservatism in nasal and tonsillar operations. *Jour.-Lancet*, Aug. 15, 1915.
- 335 WILLIAMS, W. R. The tonsil from the standpoint of the internist. *Jour. Ophthal., Otol. and Laryngol.*, July, 1915.
- 336 WILLIS, B. C. Importance of the faucial tonsil as portal of entry in tuberculous cervical adenitis; twenty-nine cases. *South. Med. Jour.*, Dec., 1915.
- *337 WINGRAVE, W. Spirochetal ulceration of tonsils in soldiers. *Lancet*, July 24, 1915.
- *338 WINGRAVE, W. Vincent's angina. A review of the present position. *Jour. Laryngol., Rhinol., and Otol.*, p. 472, Dec., 1915.
- 339 WISHART, D. J. GIBB. The tonsil operation; an inquiry into the actual results obtained in a series of cases. *Ann. Otol., Rhinol. and Laryngol.*, p. 641, Sept., 1915.
- 340 YEARSLEY, M. Abnormal styloid process causing tonsil irritation. *Jour. of Laryngol.*, p. 116, March, 1915.
- *341 YOUNG, J. H. Tonsillectomy as a therapeutic measure in the treatment of chorea and endocarditis. *Bost. Med. and Surg. Jour.*, Sept. 2, 1915.

Uvula.

- †342 GOLDSTEIN, M. A. Angioma of the uvula. *THE LARYNGOSCOPE*, p. 90, Feb., 1915.
- †343 STOUT, P. S. Tumors of the uvula with report of a case of papilloma of uvula. *THE LARYNGOSCOPE*, p. 88, Feb., 1915.
- 344 TATAR, K. Case of scleroma of the uvula, palate and nasal cavities. *Orvosi Hetilap.*, No. 31, 1915.

Tongue.

- †345 ARROWSMITH, H. Cavernous angioma of the tongue. *THE LARYNGOSCOPE*, p. 94, Feb., 1915.
- 346 BLANCARD, W. Tuberculosis of the tongue. *Jour. Cutaneous Dis.*, Oct., 1915.
- 347 BURKE, C. B. Thyroglossal cysts. Report of a case. *Med. Rec.*, June 19, 1915.
- 348 CISNEROS, E. Epithelioma of the tongue. *Semana Medica, Buenos Aires*, Oct. 28, 1915.
- §349 COUGHLIN, W. T. Sarcoma of the tongue. A study of the published cases with reports of two new cases. *Jour. A. M. A.*, p. 291, Jan. 23, 1915. *THE LARYNGOSCOPE*, p. 700, Oct., 1915.

- 350 CUMBERBATCH, E. P. Malignant growths treated by diathermy: (1) Carcinoma of tongue; (2) Carcinomatous ulcer of cheek. *Proc. Roy. Soc. Med., Electro-therap. Section*, p. 53, April, 1915.
- 351 TERRAND, L. Scrap of shell in tongue without any reaction. *Paris Medical*, Nov. 13, 1915.
- 352 GAUCHER, BIZARD and BRALEZ. Gumma of the tongue in a child 9 years old (Gomme de la langue survenue à l'âge de neuf ans chez une syphilitique héréditaire, présentant de l'écartement des incisives médianes supérieures). *Ann. des mal. vénériennes*, May, 1915.
- 353 GILMORE, A. A case of chancre of the tongue. *N. Y. State Jour. Med.*, Aug., 1915.
- 354 GOODHART, J. Nervous furring of tongue and disturbed taste. *Lancet*, Sept. 25, 1915.
- 355 HILL, WM. Malignant tumor at base of tongue almost dispersed by radium. *Proc. Roy. Soc. Med. Laryngol. Sect.*, p. 101, Vol. 8, June, 1915.
- 356 LOW, W. S. Epithelioma of the pharynx and tongue. *Proc. Roy. Soc. Med., Laryngological Section*, Vol. 8, No. 5, March, 1915.
- 357 LOW, W. S. Epithelioma of pharynx, tongue and cervical glands in male. *Proc. Laryngol. Section, Royal Soc. Med.*, Vol. 111, No. 6, April, 1915.
- 358 LYDSTON, G. F. Pre-cancerous lesions and transition types of malignant disease of the tongue and their relation to syphilis with remarks on early diagnosis and operation. *Am. Jour. of Surg.*, p. 33, Feb., 1915.
- 359 MAUNSELL, R. C. B. Cancer of the tongue and floor of the mouth treated by radium emanations. *Med. Press and Cir.*, May 12, 1915.
- *360 MORESTIN. Paralysis of the tongue from traumatic division of hypoglossal nerves. *Soc. de Chirurg. de Paris*, June 23, 1915.
- 361 NEUHOF, H. Acute non-epidemic inflammation of sublingual glands in children; an affection of unknown etiology. *Amer. Jour. Dis. Child.*, Vol. 10, No. 2, Aug., 1915.
- 362 POTLENGER, F. M. Healed tuberculosis of the tongue, larynx and lungs. *N. Y. Med. Jour.*, p. 22, Jan. 2, 1915.
- †363 RAYNOR, F. C. Parenchymatous glossitis following submucous resection of the septum. *THE LARYNGOSCOPE*, p. 227, April, 1915.
- *364 SIERRA, D. L. R. F. Leukoplakia of the tongue and its treatment by radium (Leucoplasia de la lengua y su tratamiento por el radium). *Revista Espanola de Urologia y Dermatologia*, p. 447, Aug., 1915.
- 365 SUMMERS, B. E. Acute glossitis. *Old Dom. Jour., Med. and Surg.*, Aug., 1915.

General.

- §366 ABBE, R. Cancer of the mouth. *N. Y. Med. Jour.*, July 3, 1915. *THE LARYNGOSCOPE*, p. 559, Aug., 1915.
- 367 ACHARD and WELTER. Pustular dermatitis propagated from the hands to the lips and mouth (Dermatite pustuleuse des mains propagée aux lèvres et à la bouche). *Soc. Med. des Hopitaux*, Feb. 19, 1915, and *Presse Med.*, March 4, 1915.

- 368 ALLEN, O. H. Report of twenty-eight cases of pyorrhea alveolaris treated with emetin hydrochlorid. *U. S. Nav. Med. Bull.*, Vol. 9, No. 3, July, 1915.
- *369 ALMKVIST, J. Pathogenesis, treatment and prophylaxis of mercurial stomatitis. *Hygiea*, Vol. 77, No. 16, 1915.
- 370 ALMKVIST, J. Salvarsan in the treatment of mercurial stomatitis. *Nord. Med. Arch.*, No. 1-4, 1915.
- *371 BAER, T. Leucoplakia; its pathogenesis and treatment with salvarsan. *Dermatologische Zeitschrift* Bd, 22, H. 3, March, 1915.
- 372 BARNES, G. E. Pyorrhea alveolaris; what it is and how prevented. *Med. Rev. Reviews*, p. 715, Dec., 1915.
- 373 BARRETT, T. M. Foreign bodies in the throat. *Jour. A. M. A.*, p. 1026, Sept. 18, 1915.
- 374 BASS, C. C. Relation of endamebae to alveolar pyorrhea. *Ind. State Med. Ass'n, Jour.*, Oct., 1915.
- 375 BASS, C. C. and JOHNS, F. M. Emetin in pyorrhea alveolaris. *N. O. Med. and Surg. Jour.*, No. 8, Feb., 1915.
- 376 BENJAMINS, C. E. Scrap of bone impacted in the pharynx. *Nederl. Tijdschr. v. Geneesk.*, Dec., 11, 1915.
- 377 BERGERON, J. Z. Vincent's angina. *Ill. Med. Jour.*, Feb., 1915.
- 378 BIGGS, H. M. Milk-borne septic sore throat; a new health problem. *Medical Record*, p. 945, Dec. 4, 1915.
- 379 BLAIR, V. P. Dental disorders and periodontal infections; their relation to neighboring organs. *Surg., Gyn. and Obst.*, April, 1914.
- 380 BLAIR, V. P. Importance of early diagnosis of malignant tumors of the month. *Dental Review*, June, 1915.
- 381 BRAV, A. Orbitopalpebral emphysema caused by perforation of a dental canal. *Jour. A. M. A.*, p. 1275, Oct. 9, 1915.
- 382 BRAXTON, H. H. Pyorrhea dentalis. *Arizona Med. Jour.*, Vol. 3, No. 8, Sept. 1915.
- 383 BRUNELLE, G. C. and GINSBERG, G. Endamoeba buccalis and alveolar dental pyorrhea. *N. Y. Med. Jour.*, p. 554, Sept. 11, 1915.
- 384 BURKHART, H. Affection of facial portion of skull in late hereditary lues with special reference to the maxilla. *Arch. f. Laryngol.*, Bd. 29, Heft 2, p. 205, 1915.
- §385 CALDERA, C. Treatment of Vincent's angina. *Arch. ital. di Otol., Rinol., Laringol.*, Vol. 25, April, 1915. *THE LARYNGOSCOPE*, p. 573, Aug., 1915.
- *386 CARMODY, T. E. Oral tuberculosis. *Ann. Otol., Rhinol. and Laryngol.*, p. 193, Vol. 24, No. 2, June, 1915.
- 387 CAVANAUGH, J. A. The topography of the pharynx. *Jour. Ophthal., Oto-Laryngol.*, No. 4, April, 1915.
- 388 CHRISTELLER, E. Hypophysis of the pharyngeal vault in normal and pathological conditions. *Virchow's Arch.*, 128, p. 185, 1915.
- 389 CLEMM, W. N. Fatal alveolar pyorrhea. *Deut. Med. Woch.*, No. 34, Aug. 19, 1915.
- 390 COLE, H. P. Hare-lip surgery; essentials in production of scarless incisions. *South. Med. Jour.*, Vol. 8, No. 9, Sept., 1915.
- 391 COLLISCHOUN, DR. A throat pastille. *Med. Klin.*, July 25, 1915.

- 392 COX, CHAS. N. Report of two cases imbedded in the wall of the pharynx. *Ann. Otol., Rhinol. and Laryngol.*, March, 1915.
- 393 DABNEY, S. G. Throat coughs. *Louisville Monthly Jour., Med. and Surg.*, p. 193, Dec., 1915.
- 394 DA MATTA, A. A. Leishmaniasis of the mouth and throat. *Brazil-Medico*, No. 10, March 8, 1915.
- 395 DAVIS, G. S. Plastic operation for buyo cheek cancer in its early stages and a further report of buyo cheek cancer cases. *Surg., Gyn. and Obst.*, p. 48, July, 1915.
- *396 DE SANCTIS, A. G. Vincent's angina. *N. Y. Med. Jour.*, p. 953, Nov. 6, 1915.
- 397 DEY, W. P. Syphilis of the oral cavity. *Jour. Florida Med. Assn.*, Feb., 1915.
- 398 FABRY, D. Chancre of the lip. *Med. Klin.*, No. 2, 1915.
- 399 FAHM. Some cases of serious sepsis in connection with sore throat. *Correspond. Blatt f. Schweizer Arzte*, Sept. 4, 1915.
- 400 FANTHAM, H. B. Spirocheta bronchialis Castellani, 1907; spirochetes of human mouth. *Ann. Trop. Med. and Parasitol.*, Vol. 9, No. 3, July, 1915.
- 401 FEDERSPIEL, M. N. The surgical correction of certain types of oral deformities. *Internat. Jour. Orthodon.*, p. 595, Dec., 1915.
- 402 FERGUSON, R. T. Stone in Wharton's duct. *Jour. So. Carolina Med. Ass'n*, Oct., 1915.
- 403 FISK, E. L. Mouth hygiene. *N. Y. Med. Jour.*, p. 273, Aug. 7, 1915.
- 404 FOSSIER, A. E. Pyorrhea alveolaris as a cause of systemic disturbance. *N. Y. Med. Jour.*, p. 286, Aug. 7, 1915.
- 405 FOSTER, H. A. Vincent's angina. *Jour. of Ophth., Otol. and Laryngol.*, p. 240, March, 1915.
- 406 FREUNDLICH, D. B. The diabetic type of pyorrhea alveolaris. The necessity for its early recognition by the medical profession. *Med. Rec.*, p. 943, June 5, 1915.
- 407 GATCH, W. D. Cancerous and precancerous lesions of the mouth and lips. *Dent. Summary*, Feb., 1915.
- 408 GIFFORD, A. C. The cause and effect of oral deformities. *Internat. Jour. Orthodon.*, p. 606, Dec., 1915.
- 409 GOODALL, A. On an outbreak of septic pharyngitis. *Edin. Med. Jour.*, p. 276, April, 1915.
- 410 GOPPERT, F. Treatment of aphthous stomatitis. *Therap. Monatschrift*, Sept., 1915.
- 411 GOSLINE, H. I. Results of the treatment of pyorrhea alveolaris and allied conditions with emetine hydrochlorid at the Danvers State Hospital. *Bost. Med. and Surg. Jour.*, July 22, 1915.
- 412 GRABER, C. M. Threshold stimulation of the chorda tympani nerve in relation to salivary secretion and vasodilatation. *Amer. Jour. Physiol.*, No. 3, Feb., 1915.
- 413 HALL, G. C. Streptococcic sore throat. *Pediatrics*, Nov., 1915.
- 414 HALLARMAN, H. Pharyngeal stenosis. *N. Y. Med. Jour.*, p. 1282, Dec. 25, 1915.

- 415 HARMER, W. D. Pharyngeal fistula in an old man after operation on abscess in the neck. *Proc. Roy. Soc. Med., Laryngol. Sect.*, p. 108, Vol. 8, No. 8, June, 1915.
- 416 HARPER, J. Vincent's angina. *Glasgow Med. Jour.*, p. 34, July, 1915.
- †417 HASKIN, W. H. The responsibility of the physician in oral infections. *THE LARYNGOSCOPE*, p. 23, April, 1915.
- 418 HEAD, J. Vaccines in relation to mouth infection. *Med. Rec.*, Jan. 16, 1915.
- 419 HEAD, J. A study of blood counts in relation to vaccine therapy. Auto-inoculation in the treatment of mouth infections. *Med. Rec.*, p. 179, July 31, 1915.
- *420 HECKER, F. A study of 250 stained blood films in pyorrhea alveolaris. *Amer. Jour. Med. Sc.*, p. 889, June, 1915.
- 421 HEIMAN, H. Parapharyngeal abscess as distinguished from retropharyngeal and peritonsillar abscesses. *Amer. Jour. Dis., Child.*, Vol. 10, No. 2, Aug., 1915.
- 422 HELLER, I. M. Acute infectious inflammations of the throat. *N. Y. Med. Jour.*, p. 406, Aug. 21, 1915.
- 423 HOXIE, G. H. Pyorrhea due to organisms other than the amebas. *Jour. A. M. A.*, p. 1908, Nov. 27, 1915.
- 424 IBERSHOFF, A. E. Carious teeth as a factor in ocular disease. *Jour. Ophthal. and Oto-Laryngol.*, May, 1915.
- 425 IBERSHOFF, A. E. Ocular disease of dental origin. *Jour. Ophthal. and Oto-Laryngol.*, p. 209, Vol. 9, No. 7, July, 1915.
- 426 INDEMANS, J. W. M. Congenital affection resembling Mikulicz's disease (Hypertrophie congenita glandularum cum lymphomate colli congenito). *Berl. Klin. Woch.*, No. 12, March 22, 1915.
- 427 JACQUES, P. A new treatment in cicatricial stenoses of the pharynx (Un procédé nouveau de cure des stenoses cicatricielles du pharynx). *Rev. de Laryngol., d'otol. et de Rhinol.*, p. 193, Nos. 36-37, Sept. 15, 1915.
- 428 JONES, H. B. Tuberculous ulcer of the mouth. *Proc. Roy. Soc. Med., Laryngological Section*, Vol. 8, No. 5, March, 1915.
- 429 KAESS. Bilateral nuclear paralysis of hypoglossal nerve (Bilaterale nucleäre hypoglossuslähmung und parese beider arme durch Unfall). *Med. Klin.*, No. 7, 1915.
- 430 KANTORWICZ, DR. Anesthesia of the second branch of the trigeminus in the pterygopalatine fossa. *Deut. Monatsh. f. Zahnheilk.*, Bd. 33, p. 128, 1915.
- 431 KING, J. J. The Connellan-King diplococci infections of the throat, with especial reference to rheumatism. *Medical Recrd.*, p. 958, Dec. 4, 1915.
- †432 KING, J. J. Preliminary report on the Connellan-King diplococcus infections of the throat. *THE LARYNGOSCOPE*, p. 299, April, 1915.
- 433 KINGSBURY, B. F. Development of the human pharynx. *Amer. Jour. Anatomy*, p. 329, Vol. 18, No. 3, Nov. 15, 1915.
- 434 KINYOUN, J. J. Uhlenhuth's method for sputum examination for tubercle bacilli. *Amer. Jour., Public Health*, Sept., 1915.

- 435 KOERNER, O. Paralysis of the buccal branches of the facial nerve following trauma to the nerve trunk. (Isolierte Lähmung der Mundäste des Nervus facialis infolge einer Schädigung des Nervenstammes innerhalb des Schläfenbeins.) *Arch. f. Ohrenheilk.*, Bd. 72, H. 4, May, 1915.
- 436 KRUMWIEDE, C. and VALENTINE, E. Bacteriologic study of epidemic of sore throat. *Jour. Med. Research*, Nov., 1915.
- 437 KUMMER, E. The descending branch of the hypoglossal nerve. *Rev. Med. de la Suisse Rom.*, Vol. 34, No. 7, July, 1915.
- 438 LADD, W. E. Modification of technic for hare-lip operation. *Boston Med. and Surg. Jour.*, Jan. 14, 1915.
- 439 LANDOUZY, L. White patches on the buccal mucosa as a sign of syphilis. *Bull. de l'acad. de Med.*, May 11, 1915.
- 440 LANDOUZY, L. White spots or "smokers' plaques" on the buccal mucosa as a sign of syphilis (Les taches blanches de la Muqueuse jugale et commissurale dites "plaques de fumeur" une fonction de syphilis). *Le Jour. de Med. et de Chir. de Montreal*, p. 298, Aug., 1915.
- 441 LANGE, S. Epithelioma of the lip cured by a single X-ray treatment. *Lan.-Chn.*, p. 461, April 24, 1915.
- 442 LANGE, S. Sarcoma of the upper jaw symptomatically cured by Roentgen ray. *Jour.-Lancet*, Nov. 6, 1915.
- 443 LEVINSTEIN, OSWALD. Pathology and therapy of pharyngitis lateralis (Pathologie und Therapie der Pharyngitis lateralis). *Arch. f. Laryngol. u. Rhinol.*, p. 403, Vol. 29, No. 3, 1915.
- 444 LEVY, ROBERT. Leukoplakia buccalis. *Jour. Ia. State Med. Soc.*, Vol. 5, p. 169, May 15, 1915.
- †445 LEVY, ROBERT. Leucoplakia buccalis et linguæ. *THE LARYNGOSCOPE*, p. 529, Aug., 1915.
- 446 LYNCH, K. M. Endamebiasis of the mouth. *Amer. Jour. Trop. Dis. and Prevent. Med.*, Oct., 1915.
- 447 LYNCH, K. M. An ameba in suppurative and hyperplastic osteoperiostitis of the inferior maxilla. *Jour. A. M. A.*, p. 2077, Dec. 11, 1915.
- 448 LYONS, RANDOLPH. Mode of action and use of emetine in entamebiasis. *Amer. Jour. Med. Sc.*, p. 97, July, 1915.
- 449 MACKENZIE, ALICE V. A large calculus in Wharton's duct. *Jour. Ophthal., Otol. and Laryngol.*, p. 881, Oct., 1915.
- 450 MACMILLAN, H. W. The use of conductive anesthesia in intra-oral operations. *Amer. Jour. Surg., Anesthesia Supp.*, p. 130, Oct., 1915.
- 451 MAKUEN, G. HUDSON. An aching throat. *Ann. Otol. Rhinol. and Laryngol.*, March, 1915.
- 452 MARINE, D. Epithelioma of pharyngeal mucosa in fowl. *Cleveland Med. Jour.*, Feb., 1915.
- *453 MASSEI, F. Vincent's angina (Qualche considerazione sulla cure odierna dell angina di Vincent). *Arch. ital. di Laringol.* p. 155, Vol. 35, No. 4, Oct., 1915.
- 454 MAYO, C. H. Mouth infection as a source of systemic disease. *Amer. Dent. Jour.*, March-April, 1915.
- 455 MCCURDY, J. R. Advanced epithelioma of the lower lip; report of a case. *Radium*, p. 64, Vol. 5, 1915.

- 456 MEANS, J. W. Precancerous conditions of the face and mouth. *Dent. Summary*, Dec., 1915.
- 457 MELLON, R. R. The early diagnosis of streptococcic sore throat and the effect of vaccines in its treatment. *Medical Record*, May 15, 1915.
- 458 MELLON, R. R. Method of early diagnosis of streptococcus sore throat. *Jour. Ophth. Otol. and Laryngol.*, p. 291, April, 1915.
- 459 MERRITT, A. H. Mouth infections: their cause, treatment and systemic effect. *N. Y. Med. Jour.*, p. 279, Aug. 7, 1915.
- 460 MERZ, MAX. A rare form of retropharyngeal connective tissue tumor (Über eine seltene retropharyngeal gelegene Bindegewebsgeschwulst mit Einschlüssen von Ganglionzellen). *Arch. f. Ohrenheilk.*, Bd. 72, H. 4, May, 1915.
- 461 MINGAZZINI, G. Unilateral glosso-pharyngo—laringo-cervical paralysis (Contributo clinico allo studio della paralysis glosso-pharingo-laringo-cervicalls unilateralis). *Arch. ital. di Otolog.*, p. 353, Vol. 26, No. 5, Oct., 1915.
- 462 MORESTIN, H. Treatment of deformities following fracture with loss of substance of the anterior arch of lower jaw. *Bull. et Mem. Soc. de chir. de Paris*, p. 1314, Vol. 41, 1915.
- 463 MUNGER, C. E. Diseases of the salivary ducts. *Ann. Otol. Rhinol. and Laryngol.*, March, 1915.
- 464 NEW, G. B. Cystic odontomas. *Jour. A. M. A.*, p. 34, Jan. 2, 1915.
- 465 NEWCOMB, W. H. The throat in relation to rheumatism. *Amer. Med.*, p. 494, June, 1915.
- 466 NEWCOMB, W. H. Formaldehyde as a bactericide and antiseptic in the treatment of certain infectious conditions and inflammatory affections of the throat. *Amer. Medicine*, Sept., 1915.
- 467 NOGUERAS, DR. Treatment of noma (La terapeutica del noma). *Los Prog. de la Clin.*, p. 61, Vol. 5, 1915.
- 468 PARKER, E. F. Membranous lesions of throat. *S. Carolina Med. Assn. Jour.*, Jan., 1915.
- 469 PELLIZARI, C. Epithelioma of the mouth cured by radium. *Lo Sperimentale*, April 30, 1915.
- 470 PERRY. Dental malocclusion. *Amer. Jour. Clin. Med.*, May, 1915.
- 471 PUSATERI, S. Verruca spinosa of the pharynx. *La Pratica Otorino-laringoiatrica*, No. 4, June, 1915.
- *472 REICHE, F. Rare forms and complications of Plaut-Vincent pharyngeal and mouth infections (Seltene Verlaufsformen und Komplikationen der Plaut-Vincent'schen Rachen—und Mundentzündungen). *Munch. Med. Woch.*, p. 219, No. 7, Feb. 16, 1915.
- 473 REINWALD, P. Hemorrhage after operations in the mouth. *Ergebn. d. ges. Zahnheilk.*, p. 263, Vol. 4, 1915.
- 474 ROLLESTON, J. D. Congenital heart disease and ulcerative sore throat. *Brit. Jour., Child. Dis.*, Sept., 1915.
- 475 SANDY, WM. and MCGAFFIN, C. G. Pneumococcic infection of the throat. *Med. Rec.*, p. 433, Sept. 11, 1915.
- 476 SCHREINER, K. Sarcoma of lateral pharyngeal wall with sudden death from suffocation. *Arch. f. Ohrenh.*, Bd. 96, Heft 3-4, p. 204, 1915.

- 477 SMITH, A. J. and BARRETT, M. T. A review of the relations of oral endamebae to pyorrhea and of the use of emetin in the treatment of amebic pyorrhea. *Dental Review*, p. 1034, Oct., 1915.
- 478 SMITH, J. M. Case of Vincent's angina. *Ga. Med. Assn. Jour.*, March, 1915.
- 479 SMITH, T. S. Emetine hydrochloride in treatment of periodontal diseases. *Jour. A. M. A.*, May 8, 1915.
- 480 SMITH, A. J. and BARRETT, M. T. The parasite of oral endamebiasis, endameba gingivalis (Gros). *Jour. Parasitology*, June, 1915.
- 481 SMITH, A. J. and BARRETT, M. T. Comparison of endameba gingivalis (Gros) and endameba histolytica Schaudinn. *Jour. Parasitology*, Dec., 1915.
- 482 STARK, H. H. Twenty-five cases of Vincent's angina successfully treated with sodium perborate—special report of three cases. *Ann. Otol. Rhinol. and Laryngol.*, March, 1915.
- 483 STARR, N. Anginas of the exanthemata. *Jour. of Ophth. Otol. and Laryngol.*, p. 237, March, 1915.
- 484 STERN, H. Perversities of internal secretions in their bearing on oral pathology. *Arch. of Diag.*, Vol. 8, No. 3, July, 1915.
- 485 STEWART, F. E. Pyorrhea alveolaris. *N. Y. Med. Jour.* p. 289, Aug. 7, 1915.
- 486 STEWART, F. E. Recent discoveries concerning the etiology and treatment of pyorrhea alveolaris. *Med. Herald*, p. 210, June, 1915.
- 487 STEWART, F. E. Pyorrhea alveolaris; its treatment with bacterin and emetin hydrochloride. *Med. Press*, Sept. 22, 1915.
- *488 STICKER, A. Fifteen cases of cancer of the mouth successfully treated with radium (Fünfzehn Fälle von Mundhöhlenkrebs mit Radium günstig behandelt). *Berl. Klin. Woch.*, p. 1040, Vol. 52, 1915.
- 489 SYNNOTT, M. J. Vincent's angina. *N. Y. Jour.*, p. 454, March 6, 1915.
- 490 TALBOT, E. S. Interstitial gingivitis and pyorrhea alveolaris. *Ill. Med. Jour.*, Dec., 1915.
- 491 TEICHE, F. End results and complications of Vincent's angina. *Munch. Med. Woch.*, p. 219, Vol. 62, 1915.
- 492 ULRICH, H. L. The blind dental abscess. *Jour. A. M. A.*, p. 1619, Nov. 6, 1915.
- 493 VITTO-MASSEI, R. A case of labio-glosso-laryngeal paralysis. *Archiv. ital. di Laringol.*, Vol. 35, No. 3, 1915.
- 494 WALKO, K. Primary smallpox in pharynx and respiratory tract. *Prag. Med. Woch.*, Bd. 40, p. 125, 1915.
- 495 WALKER, F. B. Mouth infections. *Mich. State. Med. Soc. Jour.*, June, 1915.
- 496 WHITE, S. M. Mouth infections as related to systemic disease. *N. Y. State Jour. Med.*, Dec., 1915.
- 497 WILLIAMS, ANNA W., VON SHOLLY, ANNA I., ROSENBERG, CAROLINE, and MANN, ALICE, G. Amebic mouth infections. *Jour. A. M. A.*, Dec. 11, 1915.
- 498 WRIGHT, B. L. and WHITE, P. The treatment of pyorrhea alveolaris and its secondary systemic infections by deep muscular injections of mercury. *Med. Rec.*, p. 424, March 13, 1915.

III. ACCESSORY SINUSES.

Frontal Sinus.

- 499 BANISTER, J. M. Affections of the frontal sinus. *West. Med. Rev.*, Dec., 1915.
- 500 DRISKILL, T. F. The antrum of Highmore. *Dent. Summary*, Oct., 1915.
- †501 FREER, O. T. The frontal sinus: opening it through the nose for chronic suppuration. *THE LARYNGOSCOPE*, p. 803, Dec., 1915.
- 502 GOETS, J. S. Symposium on affections of the frontal sinus. *West. Med. Rev.*, Dec., 1915.
- 503 GOOD, R. H. Acute frontal sinusitis. *Ill. Med. Jour.*, Dec., 1915.
- 504 HALLE, DR. Winckler's "Therapy of frontal sinus disease" (Zu Winckler's Arbeit: "Über Therapie der Stirnhöhlen-erkrankungen."). *Arch. f. Laryngol. u. Rhinol.*, p. 466, Vol. 29, No. 3, 1915.
- †505 IMPERATORI, C. JOHNSTONE. Report of a case of frontal sinusitis, sphenoiditis, influenzal meningitis; death, necropsy. *THE LARYNGOSCOPE*, p. 580, Aug., 1915.
- *506 LOTHROP, H. A. Frontal sinus suppuration with results of new operative procedure. *Jour. A. M. A.*, July 10, 1915.
- 507 MACFARLAN, D. The rationale of a new treatment for the acute frontal sinusitis. *Jour. Ophthal., Otol. and Laryngol.*, p. 777, Sept., 1915.
- 508 MOORE, J. L. I. Case of chronic suppuration of left frontal, ethmoidal and maxillary sinuses. *Proc. Royal Soc. Med., Laryngol. Section*, Vol. 9, No. 2, Dec. 1915.
- 509 ONODI, L. Cerebral wall of frontal sinus and its practical significance. *Arch. f. Ohrenh.*, Bd. 98, Heft 1, p. 33, 1915.
- 510 PEERY, T. E. Case of frontal and maxillary sinusitis. *W. Va. Med. Jour.*, Feb., 1915.
- 511 PEGLER, L. H. Empyema of frontal sinus. *Proc. Royal Soc. Med., Laryngol. Section*, Vol. 9, No. 2, Dec., 1915.
- 512 POSEY, W. C. Report of unusually large mucocoele of frontal and ethmoidal cells. Operation and cure. *Ophthal. Rec.*, March, 1915.
- 513 POTTS, J. B. The frontal sinus; ocular relations. *West. Med. Rev.*, Dec., 1915.
- 514 PRATT, E. L. Trichinosis simulating frontal sinusitis. A report of three cases. *Jour. A. M. A.*, p. 1277, Oct. 9, 1915.
- 515 RICHARDSON, C. W. Two cases of frontal sinusitis and the Killian operation. *Wash. Med. Ann.*, March, 1915.
- †516 SKILLEEN, ROSS H. The external operation of the frontal sinus. *THE LARYNGOSCOPE*, p. 212, April, 1915, and *Jour. Laryngol., Rhinol. and Otol.*, p. 329, Vol. 30, No. 9, Sept., 1915.
- 517 STIMSON, G. W. Empyema of frontal and ethmoidal sinuses. *J. A. M. A.*, p. 418, July 31, 1915.
- *518 THOMAS, J. B. Tuberculosis of the frontal sinus. Report of two cases. *Jour. A. M. A.*, p. 308, July 24, 1915.
- *519 WILLIAMS, P. W. Pernasal operation for frontal sinus suppuration by anterior route. *Lancet*, Feb. 20, 1915, and *Bristol Med.-Chir. Jour.*, p. 24, 1915.

Sphenoid Cells.

- 520 BOILER, W. F. Diagnosis of sphenoidal sinus empyema. *Iowa State Med. Soc. Jour.*, Vol. 5, No. 7, July, 1915.
- §521 BRADBURN, A. A. Case of bilateral optic neuritis due to sphenoid sinusitis. *Brit. Med. Jour.*, p. 109, Jan. 16, 1915. *THE LARYNGOSCOPE*, p. 197, March, 1916.
- 522 DUNN, J. S. Ganglioneuroma of the sphenomaxillary fossa. *Glasg. Med. Jour.*, p. 98, Aug., 1915.
- §523 EMBELTON, D. and PETERS, E. A. Cerebro-spinal fever and the sphenoidal sinus. *Lancet*, p. 1078, May 22, 1915. *THE LARYNGOSCOPE*, p. 710, Oct. 1915.
- †524 GRAYSON, C. P. Exploratory opening of the sphenoid sinus. *THE LARYNGOSCOPE*, p. 65, Feb., 1915.
- 525 HEINDELL, A. Removal of bullet from the sphenoid bone. *Wiener. Klin. Woch.*, No. 21, 1915.
- 526 MEYER, A. W. Sinister unrecorded anomalies of the sphenoid. *Ann. Otol., Rhinol. and Laryngol.*, p. 257, Vol. 24, No. 2, June, 1915.
- *527 MINK, P. J. Pathology and therapy of recessus sphenomaxillaris. *Arch. f. Laryngol.*, Bd. 29, Heft 2, p. 165, 1915.
- 528 PRENTISS, H. J. Some variations in sphenoidal sinus as found in anatomic laboratory of S. U. I. *Iowa State Med. Soc. Jour.*, Vol. 5, No. 7, July, 1915.
- 529 SLUDER, G. Hyperplastic sphenoiditis: and its clinical relations to the second, third, fourth, fifth, sixth and vidian nerves and nasal ganglion. *Trans. Amer. Laryngol. Ass'n*, p. 215, 1915.
- †530 STEIN, O. J. Report of a case of hypophyseal growth operated through the nose and sphenoid. *THE LARYNGOSCOPE*, p. 159, March, 1915.

Ethmoid Labyrinth.

- 531 DAVIS, J. L. Ethmoiditis; its varied effects and their probable prevention, or when fully established, their possible cure. *Penn. Med. Jour.*, Dec., 1915.
- 532 MACNAB, J. C. G. Case of monocular diplopia of three years' standing cured by exenteration of posterior ethmoids and opening of sphenoid. *Jour. Laryngol., Rhinol. and Otol.*, p. 336. Vol. 30, No. 9, Sept., 1915.
- 533 MALAN, A. Fibroma of the ethmoid. *Arch. ital. di Otol., Rhinol. e Laringol.*, Vol. 26, April, 1915.
- 534 MCCULLAGH, S. Treatment of ethmoiditis. *N. Y. Med. Jour.*, July 24, 1915.
- 535 MINER, S. G. Suppurative ethmoiditis. *Mich. State Med. Soc. Jour.*, Nov., 1915.
- 536 MOORE, F. B. A review of Mosher's intranasal surgery of the ethmoidal labyrinth. *Memphis Med. Monthly*, Sept., 1915.
- 537 MOSHER, H. P. Observations upon the intranasal exenteration of the ethmoidal labyrinth in pan-sinusitis. *Trans. Amer. Laryngol. Ass'n*, p. 147, 1915.

- 538 PRADA, ENRIQUE. Osteoma of ethmoid labyrinth and maxillary sinus complicated with nasal polypi. *Los Progresos de la Clinica*, Dec., 1915.
- 539 RHESE, DR. Bacteriology of the diseased ethmoid (Die Bakteriologie des kranken Siebbeins). *Passows' Beitrage*, p. 235, Vol. 8, No. 4, Aug. 12, 1915.
- 540 ROBERTSON, C. M. Ethmoidal exenteration. *Ann. Otol., Rhinol. and Laryngol.*, p. 319, Vol. 24, No. 2, June, 1915.
- †541 SMITH, HARMON. Blindness incidental to external ethmoidal operation. *THE LARYNGOSCOPE*, p. 216, April, 1915.

Maxillary Antrum.

- 542 ALAGNA, G. Traumatic maxillary sinusitis (Contributo alla causistica delle sinusiti mascellari traumatiche). *Boll. delle mal. dell' Orecchio, della Gola e del Naso*, No. 4, April, 1915.
- 543 BICHATON, DR. Two cases of maxillary sinusitis (Deux cas de sinusite maxillaire). *Rev. de laryngol., d'otol. et de rhinol.*, p. 199, No. 36-37, Sept. 15, 1915.
- 544 BOURQUET, M. J. The internal wall of the maxillary sinus (La paroi interne du sinus maxillaire, les fontanelles nasales). *Ann. des mal. de l'oreille*, p. 652, Vol. 40, No. 7, 1914-1915.
- 545 CASTLE, C. H. Empyema of the maxillary sinus treated with phylacogen. *Lancet-Clinic*, May 22, 1915.
- 546 FINZI, N. S. and HETT, G. S. Inflammatory disease of the maxillary antrum—its diagnosis and treatment. *Practitioner*, Vol. 95, No. 1, July, 1915.
- §547 FINZI, N. S. and HETT, G. S. Roentgenography of maxillary antrum. *Arch. Radiol. and Electrother.*, Vol. 20, No. 2, July, 1915. *THE LARYNGOSCOPE*, p. 713, Oct., 1915.
- †548 GOLDSTEIN, M. A. Lipoma of the maxillary antrum. *THE LARYNGOSCOPE*, p. 142, March, 1915.
- 549 KYLE, J. J. A brief reference to the surgery of the antrum of Highmore. *Medical Times*, Aug., 1915.
- 550 LOMBARD, BLECHMANN, G. and BLOCH, A. Septicemia in the course of fronto-maxillary sinusitis (Septicémie avec déterminations sereuses multiples au cours d'une sinusite fronto-maxillaire). *Ann. des mal. de l'oreille* p. 667, Vol. 40, No. 7, 1914-1915.
- 551 LOW, W. STUART. Case of recurrent columnar-celled carcinoma of the antrum. *Proc. Roy. Soc. Med., Laryngol. Sec.*, p. 106, Vol. 8, No. 8, June, 1915.
- *552 MACNAB, J. C. G. A modification of Skillern's preturbinal operation on the maxillary sinus. *Jour. Laryngol., Rhinol. and Otol.*, p. 333, Vol. 30, No. 9, Sept., 1915.
- 553 MARQUIS, G. P. The operation of choice in maxillary sinus disease. *Ann. Otol. Rhinol. and Laryngol.*, March, 1915.
- *554 McNAUGHT, H. Intransal antrum operations with report of ninety cases. *Jour. A. M. A.*, p. 872, Sept. 4, 1915.
- 555 MILLIGAN, W. Inoperable angiofibroma; maxillary antral sarcoma. *Proc. Roy. Soc. Med., Laryngol. Section*, No. 8, 1915.

- 556 MOORE, I. Growth in maxillary sinus extending into nasopharynx. *Proc. Roy. Soc. Med., Laryngol. Sect.*, p. 104, Vol. 8, No. 8, June, 1915.
- 557 MORESTIN, H. Extraction of a ball from the maxillo-pharyngeal space (Extraction par la bouche d'une balle arrêtée dans l'espace maxillo-pharyngien). *Soc. de chir. de Paris*, Feb. 10, 1915; *Presse Med.*, Feb. 18, 1915.
- 558 MURPHY, J. W. Diseases of the maxillary antrum. *Texas Dent. Jour.*, p. 31, 1915.
- 559 ROBBINS, E. D. Chronic maxillary sinusitis; its remote effects as applied to dentistry. *Dent. Summary*, May, 1915.
- 560 SCHLESINGER. Radical, endo-nasal operation of the maxillary sinus, (Sturmann's operation). *Ztschr. f. Laryngol.*, Bd. 7, Heft 4, p. 461, 1915.
- 561 SOLENBERGER, A. R. Subperiosteal abscess of maxillary sinus. *Colo. Med.*, Vol. 12, No. 9, Sept., 1915.
- 562 SYKES, E. M. Case of double chronic maxillary sinusitis complicated with unilateral ethmoiditis. *Texas State Jour. Med.*, Oct., 1915.
- §563 WHALE, H. L. Perithelioma of maxillary antrum. *Proc. Laryngol. Section, Royal Soc. Med.*, Vol. 8, No. 6, April 1915. *THE LARYNGOSCOPE*, p. 195, Mar., 1916.

General.

- 564 BECK, J. C. Ultimate results of operations for chronic sinus disease, chronic tonsillar, and tonsillar and adenoid disease and chronic diseases of the middle ear. *Jour. Ophth. and Oto-Laryngol.*
- 565 BEEBE, H. M. Skiagraphic diagnosis of nasal accessory sinuses. *Jour. Ophth., Otol. and Laryngol.*, p. 319, April, 1915.
- 566 BEEBE, H. M. Sinus roentgenography. *Jour. Ophthal., Otol. and Laryngol.*, p. 962, Nov., 1915.
- 567 BERRY, H. M. Roentgenography in diagnosis of diseases of the accessory nasal sinuses. *Arch. of Roentgen Ray*, London, No. 12, May, 1915.
- *568 BLISS, M. A. Importance of the paranasal sinuses in the explanation of pain in the face, head, neck and shoulders. *Ann. Jour. Med. Sci.*, p. 230, Feb., 1915.
- 569 BRADFELD, J. A. L. Sinusitis. *Wis. Med. Jour.*, April, 1915.
- 570 BROPHY, J. A. Three cases of monocular blindness due to sinus obstruction with recovery of vision. *Ophthalmic Rec.*, Vol. 24, No. 7, July, 1915.
- 571 CAVANAUGH, J. A. Non-suppurative sinus disease in relation to the eye. *Ill. Med. Jour.*, Sept., 1915.
- 572 CLAY, J. V. F. Influence of nasal accessory sinus disease upon the ear. *Hahnemannian Monthly*, Sept., 1915.
- 573 COFFIN, L. A. New, non-operative treatment of disease of the accessory sinuses of the nose. *Med. Record*, p. 556, April 3, 1915.
- †574 COFFIN, L. A. Non-operative treatment of the accessory sinuses. *THE LARYNGOSCOPE*, p. 832, Dec., 1915.

- 575 DAVIES, D. L. Affections of the eye and orbit in diseases of the nasal accessory sinuses. *Clin. Jour.*, p. 185 and 197, June 9 and 16, 1915.
- 576 DENKER, A. The surgical treatment of suppuration of the accessory cavities after injuries in war (Die chirurgische Behandlung der Nebenhöhlenentzündungen nach Kriegsverletzungen). *Munch. med. Woch.*, No. 24, p. 821, June, 1915.
- *577 DIEBOLD, F. Diagnosis and treatment of suppuration of the nasal accessory cavities (Nasenhöhlenentzündungen). *Correspond. Blatt. f. Schweizer Aerzte*, Aug. 14, 1915.
- 578 DOWLING, J. I. Accessory nasal sinus disease a possible cause of suicide. *Jour. Ophthalmol., Otol., and Laryngol.*, p. 1006, Dec., 1915.
- 579 FERRERI, G. Lesions of the nose and accessory sinuses in war. *Policlinico*, Oct. 10, 1915.
- 580 FOOTE, E. C. Accessory sinus disease. *West. Med. Rev.*, Nov., 1915.
- 581 GOGARTY, O. St. J. Latent empyemata of the nasal accessory sinuses. *Jour. of Laryngol.*, p. 9, Jan., 1915.
- 582 GROVE, W. E. Nasal sinus infection in relation to general disease. *Wis. Med. Jour.*, April, 1915.
- 583 HASELTINE, B. Obscure sinus disease in relation to general health. *Jour. Ophth., Otol. and Laryngol.*, p. 13, Jan., 1915.
- 584 JOHNSON, G. B. Trifacial neuralgia from nasal and accessory sinus disease. *Penn. Med. Jour.*, Vol. 18, p. 448, 1915.
- 585 KYLE, J. J. Nasal sinus suppuration. *West. Med. Times*, July, 1915.
- 586 LEWIS, J. D. Empyema of the nasal accessory sinuses. *St. Paul Med. Jour.*, Feb., 1915.
- *587 MACNAB, J. C. G. Three cases showing the shoulder-arm-hand syndrome of paranasal sinus disease cured by operation. *Jour. Laryngol., Rhinol. and Otol.*, p. 334, Vol. 30, No. 9, Sept., 1915.
- 588 MANGES, W. F. Roentgen ray examination of the accessory sinuses. *Penn. Med. Jour.*, April, 1915.
- 589 MAYBAUM, J. L. Diagnosis and conservative treatment of inflammation of the accessory sinuses of the nose. *Med. Rec.*, June 19, 1915.
- 590 McDUGALL, J. C. Empyema of the accessory sinuses. *Atlanta Jour. Rec. Med.*, June, 1915.
- 591 MINK, P. J. The function of the nasal accessory cavities (Über die Funktion der Nebenhöhlen der Nase). *Arch. f. Laryngol. u. Rhinol.*, p. 453, Vol. 29, No. 3, 1915.
- 592 MURRAY, W. R. Nasal accessory sinuses as foci of infection. *Jour. Lancet*, Oct. 15, 1915.
- 593 ONODI, A. Treatment of diseases of nasal accessory sinuses in infants (Die Nasennebenhöhlenerkrankungen in den ersten Lebensjahren). *Jahrbuch f. Kinderheilk.*, Bd. 81, H. 2, Feb., 1915.
- 594 PERRY, T. E. An unusual case of sinusitis. *West Va. Med. Jour.*, Feb., 1915.

- 595 REEDER, J. E. Eye symptoms and their value in chronic sinusitis. *Iowa State Med. Soc. Jour.*, Nov., 1915.
- 596 SCHENCK, C. P. Accessory sinusitis. *Texas State Jour. Med.*, Oct., 1915.
- 597 SEIDEL, O. Injuries and diseases of the nose and accessory cavities in war and their treatment (Über Verletzungen und Erkrankungen der Nase und ihrer Nebenhöhlen im Kriege und ihre Behandlung). *Munch. med. Woch.*, No. 24, p. 825, June, 1915.
- 598 SHAMBAUGH, G. E. Two cases of chronic pan-sinusitis associated with systemic infection. *Trans.-Amer. Laryngol. Ass'n*, p. 179, 1915.
- 599 SMITH, F. N. Acute and chronic sinusitis of nasal and pharyngeal origin. *Mich. State Med. Soc. Jour.*, Jan., 1915.
- 600 SMITH, HARMON. Empyema of the nasal accessory cavities. *N. Y. Med. Jour.*, June 26, 1915.
- *601 STARK, H. H. Sudden blindness due to suppuration of the nasal accessory sinuses with report of three cases. *Jour. A. M. A.*, p. 1513, Oct. 30, 1915.
- †602 SULLIVAN, J. J. The examination of 106 cases of nasal accessory sinusitis, showing the value of routine examination and certain internasal operative procedures. *THE LARYNGOSCOPE*, p. 667, Sept., 1915.
- 603 SYME, W. S. Treatment of nasal accessory sinus disease. *Practitioner*, p. 789, June, 1915.
- 604 TILLEY, H. Aspergillosis of nasal accessory sinuses. *Jour. of Laryngol.*, p. 145, April, 1915.
- *605 UFFENORDE, W. The treatment of the various forms of inflammation of the mucous membrane of the nasal accessory sinuses (Die verschiedenen Entzündungsformen der Nasennebenhöhlenschleimhaut und ihre Behandlung). *Arch. f. Ohrenheilk.*, Bd. 72, H. 4, May, 1915.
- *606 WALDRON, C. W. Roentgenology of the accessory nasal sinuses with special reference to sinusitis in children. *Interstate Med. Jour.*, p. 1031, Vol. 22, No. 10, Oct., 1915.
- 607 WATERS, C. A. and WALDRON, C. W. Modification of the occipito-frontal position in roentgenology of the nasal accessory sinuses. *Amer. Jour. Roentgenology*, Feb., 1915.
- 608 WHELCHER, H. C. Inflammatory affections of nasal mucosa and accessory sinuses in children. *Ga. Med. Ass'n Jour.*, March, 1915.
- †609 WILSON, J. C. The etiology of pansinusitis. *THE LARYNGOSCOPE*, p. 823, Dec., 1915.
- 610 WINSLOW, J. R. A report of some unusual cases of diseases of the nasal accessory sinuses. *Trans. Amer. Laryngol. Ass'n*, p. 184, 1915, and *Ann. Otol., Rhinol. and Laryngol.*, p. 501, Sept., 1915.

IV. LARYNX. BRONCHI. TRACHEA. ESOPHAGUS.

Epiglottis.

- 611 PUSATERI, SANTI. Cysts of the epiglottis (Contributo allo studio delle cisti dell'epiglottide). *Arch. ital. di Laringol.*, April, 1915.

- 612 PESCHARDT, C. Incarceration of the epiglottis. *Ugesk. f. Leger*, March 4, 1915.

Voice and Speech Defects.

- 613 BELL, M. Dictionary of English sounds. *Volta Rev.*, Vol. 17, No. 8, August, 1915.
- 614 BELL, A. M. Cure of stammering and other impediments of speech. *Volta, Review*, p. 494, Dec., 1915.
- 615 BELL, A. M. Phonetic syllabication. *Volta Rev.*, p. 97, March, 1915.
- 616 BELL, A. M. Principles of speech and dictionary of sounds. *Volta Rev.*, Jan.-March, 1915.
- 617 BENEDICT, A. L. Speech-elementary sounds. *Medical Record*, May 15, 1915.
- 618 BINGHAM, K. T. Speech is as natural to the deaf child as to the hearing child. *Volta Rev.*, p. 25, Jan., 1915.
- 619 BOTEX, R. Pharyngeal voice (De la voix pharyngée). *Ann. des mal. de l'oreille*, p. 675, Vol. 49, No. 7, 1914-1915.
- 620 BUELL, E. M. Easy and natural speech. *Am. Ann. of Deaf*, p. 129, March, 1915.
- 621 CORLAT, I. H. Stammering as psychoneurosis. *Jour. Abnormal Psychol.*, Feb.-March, 1915.
- 622 DRUMMOND, W. B. Idioglossia. *Brit. Med. Jour.*, April 17, 1915.
- 623 FERRERI, G. and BILANCIONI, G. Laryngeal and respiratory graphic curves as indices of the glottologic varieties of Italian dialects (La grafica laringea e respiratoria come indice della varietà glottologica dei dialetti italiani). *Arch. ital. di Otol., Rinol. e Laringol.*, Vol. 26, No. 1, p. 1, January, 1915.
- 624 FISCHER, H. The loud and ringing element of speech. (Das lautliche und Klangliche sprach element.) *Stimme*, p. 82, Jan., 1915.
- 625 FRÖSCHELS, E. Nystagmus and stuttering. *Monatschr. f. Ohrenheilk.*, Bd. 46, p. 161, 1915.
- 626 FRÖSCHELS, E. Central mechanism of speech (Über den zentralen Mechanismus der Sprache). *Deut. Zeitschr. f. Nervenheilk.*, Bd. 54, H. 1, 1915.
- 627 GUTZMANN, A. Care of the speech of children. *Volta Rev.*, p. 121, Jan., 1915.
- 628 HEAD, H. Haughlings Jackson on aphasia and kindred affections of speech. *Brain*, July, 1915.
- 629 KEITH, J. R. Vaso-dilators in the treatment of hysterical aphonia. *Brit. Med. Jour.*, p. 847, May 15, 1915.
- †630 MAKUEN, G. HUDSON. The Voice—The relation of the lymphoid tissue in the upper respiratory tract to the voice. *THE LARYNGOSCOPE*, p. 46, Jan., 1915.
- 631 MAKUEN, G. HUDSON. The relation of the lymphoid tissue in the upper respiratory tract to the voice. *Therapeutic Gazette*, Feb. 15, 1915.
- *632 MAKUEN, G. HUDSON. A study of one thousand cases of stammering with special reference to the etiology and treatment of the affection. *Volta Review*, July, 1915.

- *633 MAKUEN, G. HUDSON. The psychology of stammering. *New York Med. Jour.*, July 17, 1915.
- 634 MAKUEN, G. HUDSON. Report of three unusual cases of stammering. *New Albany, Med. Herald*, Oct., 1915.
- 635 MARAGE. Treatment of mutism following war injuries. *Compt. rend. de l'Acad. des Sciences*, No. 29, Nov. 15, 1915.
- 636 McLOONE, J. J. Defects of the singing voice due to nasal and accessory sinus disease. *Jour. A. M. A.*, p. 310, July 24, 1915.
- 637 McLEARY, J. R. Tonsillectomy and its effect upon the voice. *Clinique*, p. 123, March, 1915.
- 638 MOTT, F. W. The psychic mechanism of the voice in relation to the emotions. *Brit. Med. Jour.*, p. 845, Dec. 11, 1915.
- †639 NATIER, MARCEL. Hysterical mutism caused by shell explosion. *THE LARYNGOSCOPE*, p. 584, Aug., 1915.
- 640 O'MALLEY, J. F. Functional aphonia. *Proc. Roy. Soc. Med., Laryngol. Sect.*, p. 116, Vol. 8, No. 8, June, 1915.
- 641 PROCTOR, A. P. Three cases of concussion aphasia; treatment by general anesthesia. *Lancet*, Oct. 30, 1915.
- 642 REED, F. A. Speech impediments and their correction in the public schools of Detroit. *Volta Rev.*, p. 7, Jan., 1915.
- 643 REED, F. A. Speech is purely mechanical. *Volta Rev.*, p. 59, Feb., 1915.
- 644 REED, R. A clinical study of stammering. *Lancet-Clinic*, Oct., 1915.
- 645 RETHI, L. Amount of air used in singing. *Wiener med. Woch.*, Bd. 28, p. 598, 1915.
- 646 RICE, B. M. More about speech-reading and the experience system. *Volta Rev.*, p. 65, Feb., 1915.
- 647 ROTHE, K. K. Elementary classes for speech defects (Das erste Jahr in der Sonder-Elementarklasse für sprachkranke Kinder). *Monatschr. f. Ohrenheilk.*, p. 168, Bd. 46, 1915.
- 648 SCHMITZ, A. The vocal teacher and functional disturbances of the voice (Der Gesanglehrer und die funktionellen Stimmstörungen). *Die Stimme*, Vol. 9, H. 9, p. 209, June, 1915.
- 649 SCHOLZ, W. Aphonia and mutism from fright neurosis (Funktionelle Sprachlähmung im Felde). *Medizin Klinik*, Dec. 26, 1915.
- 650 SICARD, J. A. Simulating deaf-mutism in war (psycho-néuroses de guerre; simulateurs sourds-muets). *Paris Medical*, p. 423, Oct. 23, 1915.
- 651 SILVA, GIULIO. The pneumograph in singing instruction (La pneumografia nell' insegnamento del canto). *Arch. ital. di otol., rinol. e laringol.*, p. 310, Vol. 26, No. 4, Aug., 1915.
- 652 SOLOMON, M. Remarks on D Corliots' paper, "Stammering as a Psychoneurosis." *Jour. Abnorm. Psychol.*, June-July, 1915.
- 653 SPENCER, B. B. How to develop spontaneous language. *Volta Rev.*, p. 60, Feb., 1915.
- 654 SWIFT, W. B. Voice sign in chorea—technic of elicitation. *Am. Jour. Dis. of Children*, Feb., 1915.
- 655 SWIFT, W. B. Elimination of voice defects following adenoid and tonsil operations. *Bost. Med. and Surg. Jour.*, Nov. 18, 1915.

- 656 SWIFT, W. B. Psychologic analysis of stuttering. *Jour. Abnormal Psychology*, Vol. 10, No. 4, Oct.-Nov., 1915.
- 657 SWIFT, W. B. The voice in tables—voice sign. *Amer. Jour. Insanity*, Oct., 1915.
- 658 SWIFT, W. B. Studies in speech disorder. *Bost. Med. and Surg. Jour.*, Nov. 4-11, 1915.
- 659 TAYLOR, H. "Speech-elementary sounds." *Volta Rev.*, Vol., 17, No. 8, Aug., 1915.
- 660 VOORHEES, I. W. Some common voice troubles in singers. *Med. Rec.*, May 8, 1915.
- 661 WEINBERG, K. Stuttering among children in the public schools of Stockholm. *Skolhygien. undersokninger*, 2, 1915.
- 662 WORCESTER, A. E. Pronunciation at sight. How shall our children be taught to pronounce at sight the words of our written language? *Volta Rev.*, p. 85, March, 1915.
- 663 WRIGHT, A. R. Manual and oral combination. *Am. Ann. of Deaf*, p. 219, May, 1915.

Larynx.

- 664 ALERNSTEDT, W. Congenital stridor. *Nord. med. arch.*, Nos. 1-4, 1915.
- 665 ARNOLDI, PAUL. Aponia spastica (Beitrag zur Aponia spastica [Dysphonia]). *Quang. Diss.*, Wuerzburg, Aug., 1915.
- 666 ARMBRUSTER. The larynx of the new-born infant (Kehlkopf der Neugeborenen). *Der Kinderarzt*, p. 3, Jan., 1915.
- 667 ASCHMAN, G. A. Diagnosis and treatment of chronic laryngitis. *W. Va. Med. Jour.*, Oct., 1915.
- 668 BARTON, E. A. Condition of the larynx and trachea in the still-born infant and its bearing on artificial respiration. *Jour. Obst. and Gyn. of Brit. Emp.*, Vol. 27, Nos. 3-5, March-May, 1915.
- 669 BARWELL, H. Laryngeal tuberculosis. *Brit. Jour. f. Tub.*, Jan., 1915.
- 670 BARWELL, H. Two cases of cicatricial stenosis of larynx. *Lancet*, Jan. 1, 1915.
- §671 BETRAN Y CASTILLO, F. Partial laryngectomy. *Rev. de Med. y Cir. prac.*, Feb. 28, 1915. THE LARYNGOSCOPE, p. 656, Sept., 1915.
- 672 BILANCIONI, G. Case of foreign body in the air passages. *Arch. ital. di Laringol.*, p. 166, Vol. 35, No. 4, Oct., 1915.
- 673 BINNEY, E. H. Laryngeal obstructions (Surgical disorders in childhood). *Australia Med. Jour.*, July 15, 1915.
- 674 BLAESSIG, K. Vocal cord paralysis in war (Beitrag zur funktionellen Stimmbandlähmung im Felde). *Muench. med. Woch.*, p. 835, No. 24, June, 1915.
- 675 BLEYL, DR. Gunshot wounds of the larynx (Zur Kasuistik der Schussverletzungen des Kehlkopfs). *Ztschr. f. Ohrenheilk.*, Bd. 73, H. 1, p. 22, July, 1915.
- 676 BOEHLER, L. Gunshot wounds of the larynx. Report from the battlefield in Russian Poland. *Surg., Gyn. and Abst.*, p. 275, Vol. 21, No. 3, Sept., 1915.

- 677 BOEHLEH, DR. Bullet wounds of the larynx (Kehlkopfschüsse). *Munch. med. Woch.*, No. 24, p. 828, June, 1915.
- 678 BOTEY, R. Modern surgery of laryngeal cancer (La cirugía moderna del cancer laríngeo). *Arch. de rinol., laringol. y otolog.*, Vol. 26, No. 190, p. 5, Jan., 1915.
- 679 BRUZZONE, C. Histology, pathology and therapy of laryngeal papillomata. *Arch. ital. di Laringol.*, p. 13, Jan. 5, 1915.
- 680 CARPENTER, E. W. Report on forty-six cases of intubation. *Pediatrics*, p. 324, July, 1915.
- §681 CASSELBERRY, W. E. Infective lymphoid growths of the laryngopharynx, secondary to sinus suppuration. *Jour. A. M. A.*, p. 576, Feb. 13, 1915. THE LARYNGOSCOPE, p. 866, Dec., 1915.
- 682 CASTELLANI, L. Foreign bodies in the air passages (Contributo allo studio dei corpi stranieri delle vie aeree). *La Pratica Otorino-laringoiatrica*, No. 4, June, 1915.
- 683 COLLET, M. A new type of associated laryngeal hemiplegia (Glossolaryngoscapulo-pharyngo hemiplegia). *Ann. des Mal. de l'oreille*, p. 740, Vol. 40, No. 8, 1914-1915, and *Lyon Med.*, No. 4, April, 1915.
- *684 COLLET, M. Technique of laryngeal heliotherapy (Technique de l'héliothérapie laryngée). *Rev. de Laryngol., d'Otol. et de Rhinol.*, No. 38-39, Sept. 30, 1915.
- 685 CORDOBA, J. Accident during laryngeal operation (Herida de laringe en un acto operatorio). *Arch. de rinol., laringolog. y otolog.*, Vol. 26, No. 190, p. 40, Jan., 1915.
- 686 COX, HERMANN. Variations of the normal laryngeal picture (Varianten des normalen Kehlkopfbildes). *Inaug. Diss., Rostock*, Sept., 1915.
- 687 DAVIES, E. D. Laryngeal stenosis following bayonet wound treated by intubation. *Proc. Roy. Soc. Med., Laryngol. Sect.*, p. 120, Vol. 8, No. 8, June, 1915.
- 688 DUPUY, H. Cancer of the larynx. *N. O. Med. and Surg. Jour.*, July, 1915.
- 689 FRIEDBERG, S. O. Foreign bodies in the respiratory tract. *Ill. Med. Jour.*, Sept., 1915.
- 690 GERMONIG, E. Tuberculous laryngitis in a two-year-old child. *Wien. Klin. Woch.*, V. 28, No. 24, June, 17, 1915.
- 691 GLAS, E. Laryngeal cases from field hospital. *Monat. f. Ohrenheilk.*, Vol. 99, 1915.
- 692 GOLDBACH, L. J. Squamous-cell carcinoma of the larynx. *N. Y. Med. Jour.*, p. 513, Sept. 4, 1915.
- 693 GOLDSMITH, P. Traumatic paralysis of the left vocal cord. *Proc. Royal Soc. Med., Laryngol. Section*, Vol. 9, No. 2, Dec., 1915.
- 694 GRAEF, C. Two cases of laryngeal obstruction and one other. *Med. Record*, p. 604, April 10, 1915.
- 695 GREENE, J. B. Laryngeal tuberculosis. *South. Med. Jour.*, Nov., 1915.
- 696 GRUENWALD, L. Genesis and treatment of non-chondritic laryngeal fistulae. *Ztschr. f. Orhenh.*, Bd. 72, Heft 2, p. 105, 1915.

- 697 GUISEZ. Wounds of the larynx and trachea in war. *Paris Medical*, Nos. 20-21, Sept. 25, 1915.
- *698 HANSEN, T. BERGTRUP. Treatment of laryngeal tuberculosis in sanatoria for pulmonary tuberculosis. *Ugesk. f. Læger*, Sept. 23, 1915.
- 699 DE HASS, T. W. Laryngismus stridulus. *Indianap. Med. Jour.*, p. 286, July, 1915.
- 700 HARMS, H. Case of aspiration of a grain of rye with spontaneous cure in an infant of 8 months. *Ztschr. f. Ohrenh.*, Bd. 72, Heft 3, p. 160, 1915.
- 701 HASTINGS, H. Tuberculosis of the larynx with special reference to tuberculin. *Trans. Amer. Laryngol. Ass'n*, p. 353, 1915.
- 702 HOFFMANN, R. A phenomenon occasionally observed in recurrent paralysis (Über ein bei Rekurrensparalyse zuweilen zu beobachtendes Phänomen). *Ztschr. f. Laryngol. u. Rhinol.*, Bd. 7, H. 5, April, 1915.
- 703 HOLINGER, J. Tuberculosis of the larynx. *Ill. Med. Jour.*, June, 1915, and *Ann. Otol., Rhinol. and Laryngol.*, p. 327, Vol. 24, No. 2, June, 1915.
- 704 HORNE, W. J. Pachydermia laryngis. *Proc. Roy. Soc. Med., Laryngological Section*, Vol. 8, No. 5, March, 1915.
- †705 HUBBARD, THOMAS. Papilloma of the larynx. *THE LARYNGOSCOPE*, p. 652, Sept., 1915.
- †706 INGALLS, E. F. Symptoms and diagnosis of laryngeal tuberculosis. *THE LARYNGOSCOPE*, p. 13, Jan., 1915.
- 707 JONES, R. M. Papilloma of the larynx. *Jour. Ophthal., Otol. and Laryngol.*, June, 1915.
- 708 KAUFMAN, A. S. Laryngeal tuberculosis. *N. Y. Med. Jour.*, p. 460, Aug. 28, 1915.
- 709 KELSON, W. H. Case of laryngeal obstruction following specific infection in a young man. *Proc. Roy. Soc. Med., Laryngological Section*, Vol. 8, No. 5, March, 1915.
- 710 KOFLER, K. Twenty-five cases of foreign bodies in the larynx, trachea or bronchi. *Wiener Klin. Woch.*, Nos. 40-41, Oct., 1915.
- 711 KOFLER, K. and FRUEHWALD, V. Gunshot and shell wounds of the larynx and trachea; seventeen cases. *Wiener Klin. Woch.*, No. 49, Dec. 9, 1915.
- 712 KOERNER, O. Injuries of the larynx and vagus nerve sustained in the war. *Zeitschr. f. Ohrenheilk.*, Bd. 72, H. 2, p. 65, 1915.
- 713 KOERNER, O. Further experiments with military wounds of the larynx and nervus vagus. *Ztschr. f. Ohrenh.*, Bd. 72, Heft. 3, p. 125, 1915.
- 714 KOERNER, O. Gunshot wounds of the larynx (Beobachtungen über Schussverletzungen des Kehlkopfs). *Arch. f. Ohrenheilk.*, Bd. 73, H. 1, p. 27, July, 1915.
- 715 LASAGNA, F. A rare form of laryngeal stenosis (Di una rara forma laringea stenosa). *Arch. ital. di. otol., rinol. e laryngol.*, p. 323, Vol. 26, No. 4, Aug., 1915.
- 716 LEVESQUE, DR. Sub-glottic abscess (Abcès sous-glottique sténosant). *Rev. de laryngol., d'otol. et de rhinol.*, p. 179, No. 34-35, Aug., 1915.

- 717 LEVINSON, A. Differential diagnosis of membranous croup by auscultation of the larynx. *Muench. med. Wchnschr.*, Feb. 2, 1915.
- 718 LEWIS, DEAN. Laryngeal diverticula. *Ann. of Surg.*, Jan., 1915.
- 719 LIEBAULT, G. Abnormal projection of the cartilages of Santorini (Saillie anormale des cartilages de Santorini et corne laryngée). *Rev. de laryngol., d'otol. et de rhinol.*, p. 181., Nos. 34-35, Aug. 1915.
- 720 LOEWY, I. D. Tuberculosis of the larynx. *New Mexico Med. Jour.*, Dec., 1915.
- 721 LOEFLE, W. Alcoholic polyneuritis with paralysis of left side of diaphragm and left vocal cord. *Deut. Med. Woch.*, Oct. 28, 1915.
- 722 LOW, W. STUART. Dry cupping in laryngeal affections. *Lancet*, Jan. 23, 1915.
- 723 MACNAB, J. C. G. Congenital (?) closure of the laryngeal pharynx. *Jour. Laryngol., Rhinol. and Otol.*, p. 337, Vol. 30, No. 9, Sept., 1915.
- 724 MACNAB, J. C. G. False ankylosis of right crico-arytenoid articulation. *Jour. Laryngol., Rhinol. and Otol.*, p. 337, Vol. 30, No. 9, Sept., 1915.
- 725 MARTINEZ, E. Laryngeal papilloma cured by radium (Dos casos de papilomas de la glotis curado por el radium). *Soc. de Estudios Clin. de la Habana*, April, 1915.
- 726 MCCOY, J. Case of multiple laryngeal papillomata in a child. Treatment by direct laryngoplasty and fulguration. *Internat. Jour. Surg.*, April, 1915.
- 727 MERELLI, GINO. Investigations on motor endings in the laryngeal muscles (Ricerche sulle terminazioni nervosi motrici dei muscoli laringei). *Arch. ital. di otol., rinol., e laryngol.*, p. 326, Vol. 26, No. 4, Aug., 1915.
- 728 MEYER, E. L. Papilloma of the larynx. Demonstration of specimen and patient. *Bull. of St. Louis Med. Soc.*, p. 109, March 4, 1915.
- 729 MICHAEL, KARL. Phlegmonous inflammation of the pharynx and larynx (Phlegmonöse Entzündungen des Rachenringes und des Kehlkopfes). *Inaug. Diss., Wuerzburg*, Aug., 1915.
- 730 MILLER, EDWARD. Bilateral peripheral paralysis of the posterior crico-arytenoid muscles. *Zeitschr. f. Laryngol.*, p. 685, Vol. 7, H. 6, Sept., 1915.
- 731 MILLIGAN, W. Laryngeal papillomata in children. *Med. Chronicle*, p. 273, Feb., 1915.
- 732 MILLIGAN, W. Subglottic growth; removal; recovery. *Proc. Laryngol. Section, Royal Soc. Med.*, Vol. 8, No. 6, April, 1915.
- 733 MOLLISON, W. M. Case of fixation of right cricoarytenoid joint; traumatic (?). *Proc. Roy. Soc. Med. Laryngological Section*, Vol. 8, No. 5, March, 1915.
- 734 MOLLISON, W. M. Epithelioma of larynx; removal. *Proc. Laryngol. Section, Royal Soc. Med.*, Vol. 8, No. 6, April, 1915.
- 735 NADOLCZNY, DR. Bullet wounds of the larynx (Schussverletzungen des Kehlkopfes). *Muench. med. Woch.*, No. 24, p. 826, June, 1915.

- 736 O'MALLEY, J. F. Traumatic fixation of both vocal cords. *Proc. Roy. Soc. Med., Laryngol. Sect.*, p. 122, Vol. 8, No. 8, June, 1915.
- 737 O'MALLEY, J. F. Two cases of paralysis of the left vocal cord. *Proc. Roy. Soc. Med., Laryngol. Sect.*, p. 121, Vol. 8, No. 8, June, 1915.
- 738 PAPE, H. Paralysis of the vocal cords in the field (Funktionelle Stimmbandlähmung im Felde). *Muench. med. Woch.*, No. 11, March, 16, 1915.
- 739 PAYR. Plastic operation on thyroid cartilage in paralysis of vocal cord (Plastik am Schildknorpel zur Behebung der Folgen einseitiger Stimmbandlähmung). *Deut. med. Woch.*, Oct. 21, 1915.
- 740 POLLATSCHEK, E. Six cases of gunshot wounds of the larynx. *Orvosi hetilap.*, No. 3, 1915.
- 741 QUIST-HANSEN, C. Laryngeal tuberculosis and its treatment. *Medicinsk revue*, No. 3, 1915.
- 742 RETHI, A. The technic of sagittal roentgenization of the larynx (Zur Technik der sagittalen Kehlkopf-rontenaufnahmen). *Ztschr. f. Laryngol. u. Rhinol.*, Bd. 7, H. 5, April, 1915.
- 743 RETHI, A. Treatment of dysphagia in laryngeal tuberculosis (Die Therapie der Dysphagie bei der Kehlkopf-tuberkulose). *Arch. f. Ohrenheilk.*, Vol. 98, Nos. 2-3, p. 125, Aug., 1915.
- 744 RETHI, A. An indirect method of examination of the larynx (Eine indirekte Untersuchungsmethode des Kehlkopfes [Distractio Laryngis]). *Arch. f. Ohrenheilk.*, Vol. 98, Nos. 2-3, p. 131, Aug., 1915.
- 745 ROHN, A. Post-diphtheritic stenosis of the larynx. *Jahrbuch f. Kinderheilk.*, Vol. 81, p. 357, 1915.
- 746 RUEDA, F. Experiences with local anesthesia in laryngeal operations (Mi experiencia actual de la gran operatoria laringea con anestesia local). *Arch. de rinolog., laringol. y otolog.*, Vol. 26, 190, p. 22, Jan., 1915.
- *747 RUEDI, TH. Operative treatment of laryngeal tuberculosis (Beobachtungen aus Davos über operative Behandlung der Kehlkopf-tuberkulose). *Zeitschr. f. Ohrenheilk.*, p. 174, Bd. 73, H. 3, Nov., 1915.
- 748 SALKELD, C. Case of laryngeal sarcoma. *Brit. Med. Jour.*, p. 157, Jan. 23, 1915.
- 749 SAMENGO, L. Laryngeal mycosis. *Anales de la Sociedad Argent. de Otorino-laringologia*, 1, H. 2, 1915.
- 750 SARGNON, DR. Section of the larynx at the thyroid cartilage (Section du larynx au niveau du cartilage thyroïde). *Lyon med.*, No. 4, April, 1915.
- 751 SCHEIER, M. Gunshot wounds of the larynx (Schuss-verletzungen des Kehlkopfes). *Berl. Klin. Woch.*, May, 31, 1915.
- 752 SEIFFERT, O. Large piece of bone in the larynx (Grosses Knochenstück im Kehlkopf). *Arch. f. Laryngol. u. Rhinol.*, p. 321, Vol. 29, No. 3, 1915.
- *753 SHALLCROSS, I. G. and BAYLEY, W. D. A case of laryngeal papilloma treated and cured by Roentgen rays. *Jour. Ophthal., Otol. and Laryngol.*, p. 873, Oct., 1915.

- *754 SHAMBAUGH, G. E. and LEWIS, D. Laryngeal diverticula; report of cases. *Ann. of Surg.*, Jan., 1915.
- †755 SMITH, HARMON. Cancer of the larynx complicated with laryngocele. *THE LARYNGOSCOPE*, p. 560, Aug., 1915.
- 756 SOBOTKY, J. Alcoholic injection of the internal branch of the superior laryngeal nerve. *Bost. Med. and Surg. Jour.*, Jan. 21, 1915.
- 757 STEIN, O. J. The differential diagnosis of laryngeal motor paralysis. *Ann. Otol., Rhinol. and Laryngol.*, p. 669, Sept., 1915.
- *758 STEWART, G. D. Conservative treatment of intrinsic cancer of the larynx by thyrocriotomy or thyrotomy; report of three cases. *Ann. Surgery*, Dec., 1915.
- 759 SWETNAM, C. R. K. Tuberculous laryngitis. *Ariz. Med. Jour.*, p. 7, Feb., 1915.
- 760 TAPIA, DR. Laryngectomy by Gluck's method (Un caso de laringuectomia por el metodo de Gluck). *Los Prog. de la Clin.*, p. 105, Vol. 5, 1915.
- *761 THOMSON, ST. CLAIR. Granuloma of the vocal cord. *Jour. Laryngol., Rhinol. and Otol.*, p. 425, Vol. 30, No. 11, Nov., 1915.
- 762 THOMSON, ST. C. Intrinsic cancer of the larynx, one year after operation by laryngo-fissure. *Proc. Royal Soc. Med., Laryngol. Section*, Vol. 9, No. 2, Dec., 1915.
- 763 THORNVAL, A. Foreign bodies in the bronchi; sixteen cases. *Ugeskrift for Laeger*, Dec. 23, 1915.
- 764 TILLEY, H. Two cases of functional aphonia (one including functional deafness) following the bursting of a shell in close proximity to the patient. *Proc. Roy. Soc. Med., Laryngol. Sec.*, p. 115, Vol. 8, No. 8, June, 1915.
- 765 TURNER, A. LOGAN. Foreign bodies in the larynx. *Edinburgh Med. Jour.*, p. 261, Oct., 1915.
- 766 VAGNI, D. A. Electro cauterization for laryngeal tuberculosis (La galvano-cauterizacion en la tuberculosis laríngea). *Sem. Med.*, Vol. 22, No. 24, June 17, 1915.
- 767 WEINBERG, K. Studies on the vocal cords of school children. *Hygiea*, 1915.
- 768 WELLS, W. A. Vocal strain from the standpoint of the laryngologist; its causes and prevention. *South. Med. Jour.*, Dec., 1915.
- 769 WHALE, H. L. Functional aphonia. *Proc. Roy. Soc. Med., Laryngol. Sec.*, p. 117, Vol. 8, No. 8, June, 1915.
- 770 WILSON, D. S. Tuberculosis with laryngo-pharyngeal stenosis. Report of a case. *Pediatrics*, p. 320, July, 1915.

Trachea and Bronchi.

- 771 ADAMS, J. Case of tracheal tumor. *Jour. of Laryngol.*, p. 64, Feb., 1915.
- *772 BLECHER, DR. Lung gangrene from primary stone of bronchus (Lungengangrän bei Bronchialsteinen). *Mitteilungen a. d. Grenzgeb. d. Med., u. Chir.*, Bd. 28, H. 4, 1915.
- 773 BROSE, L. D. Tracheotomy. *Lan.-Clin.*, June 5, 1915.

- *774 CAILLE, A. Foreign body pneumonias. *Arch. Pediatrics*, Dec., 1915.
- 775 CHIARI, O. Circular resection and suture of the trachea and plastic reconstruction of large defects in the trachea. *Monatschr. f. Ohrenheilk.*, p. 337, Vol. 46, 1915.
- 776 CHIARI, O. Extraction of a ball from the left main bronchus by bronchoscopy (Extraktion eines Rundkugel aus dem linken Hauptbronchus mittelst der oberen Bronchoskopie ein Jahr nach ihrem Eindringen von der rechten Schulter aus). *Monatschr. f. Ohrenheilk. u. Laryngo-Rhinol.*, p. 577, Bd. 49, H. 9 and 10, Sept.-Oct., 1915.
- 777 FINKLESTONE, B. B. and ELLIS, T. L. Report of a case of esophago-tracheal fistula. *Jour. A. M. A.*, p. 2155, Dec. 18, 1915.
- 778 FINZI, G. Foreign bodies in the bronchi (Contributo alla casistica dei corpi estranei nei bronchi). *Arch. ital. di otol., rinol. e laringol.*, Vol. 26, No. 3, p. 204, June, 1915.
- 779 FRASER, J. S. Foreign bodies removed from the bronchus and esophagus. *Edinburgh Med., Jour.*, p. 417, Dec., 1915.
- 780 FREUDENTHAL, W. On the direct galvanization and faradization of the bronchi and their tropical medicinal treatment in bronchial asthma. *Am. Medicine*, p. 176, March, 1915.
- 781 FREUND, A. Osteoplastic tracheopathia. *Beitr. z. Anat., Physiol., Pathol., und Therap.*, Bd. 8, H. 11, 1915.
- 782 FRIEDRICH, P. L. Slitting the sternum to relieve pressure on the lower air passages. *Beitr. z. klin. Chir.*, Vol. 93, No. 2, 1915.
- 783 GOEPPERT, F. Inhalation of eucalyptus oil in treatment of excessive bronchial secretion in children. *Therap. Monatshefte*, Sept., 1915.
- 784 GRANT, J. D. Rabbit's vertebra in the right lower bronchus removed by superior bronchoscopy. *Proc. Laryngol. Section, Royal Soc. Med.*, Vol. 3, No. 6, April, 1915.
- 785 HART, C. Idiopathic necrotic tracheo-bronchitis. *Berl. klin. Woch.*, April 19, 1915.
- 786 HART, C. Perforation of aortic aneurysm into the trachea. *Berl. klin. Woch.*, Sept. 13, 1915.
- 787 HICKEY, P. M. Scarf pin in bronchus. *Det. Med. Jour.*, p. 781, May, 1915.
- 788 HILL, W. Tracheal obstruction due to long latent cervical abscess following wound by shrapnel fragment. *Proc. Roy. Soc. Med., Laryngol. Sec.*, p. 124, Vol. 8, No. 8, June, 1915.
- 789 HOERHAMMER, C. Rupture of trachea (Über isolierte subkutane Trachealrupturen). *Muench. med. Woch.*, No. 27, p. 911, July 6, 1915.
- 790 HOWELL, W. W. Studies in bronchial glands. *Amer. Jour. Dis. Child.*, Vol. 10, No. 2, Aug., 1915.
- 791 IGLAUER, S. Accidental pneumothorax during tracheotomy, with report of a case. *Ann. Otol., Rhinol. and Laryngol.*, p. 303, Vol. 24, No. 2, June, 1915.
- 792 JONES, SEYMOUR. Foreign body impacted at the bifurcation of the trachea extracted by aid of the bronchoscope. *Brit. Med. Jour.*, May 8, 1915.

- 793 KELLING, G. Suppurating bronchial glands perforating into esophagus. *Arch. f. Verdauungs-Kr.*, Feb., 1915.
- 794 KOERNER, O. Traumatic hematoma of the mediastinum with displacement of the trachea and esophagus but without paralysis of the recurrent laryngeal nerve (Ein traumatisches Hämatom im Mediastinum mit starker Verdrängung der Speise und der Luftröhre, aber ohne Rekurrenslähmung). *Ztschr. f. Ohrenheilk.*, Bd. 73, H. 1, p. 33, July, 1915.
- 795 LARGE, S. H. Removal of open safety pin, point up, from the left bronchus of a seven-year-old child. *Cleveland Med. Jour.*, July, 1915.
- 796 LIEBAULT, G. Piece of bone in the trachea (Morceau d'os enclavé dans la trachée depuis deux mois extirpation par voie externe). *Rev. de Laryngol., d'Otol. et de Rhinol.*, p. 345, Nos. 50-51, Dec., 1915.
- 797 LINKER, F. A case of broncho-lithiasis (Ein Fall von Broncho-lithiasis). *Wiener med. Woch.*, No. 23, 1915.
- †798 LYNNAH, HENRY L. Foreign body in left bronchus. *THE LARYNGOSCOPE*, p. 578, Aug., 1915.
- †799 LYNNAH, HENRY L. Two cases of foreign bodies in the bronchi complicated by general emphysema. *THE LARYNGOSCOPE*, p. 574, Aug., 1915.
- 800 MAACFIE, J. W. S. Bronchial spirochetosis. *Jour. Trop. Med. and Hyg.*, March 15, 1915.
- 801 MACNAB, J. C. G. Acute streptococcal infection of the trachea in an infant aged fifteen months. *Jour. Laryngol., Rhinol. and Otol.*, p. 336, Vol. 30, No. 9, Sept., 1915.
- 802 MALAN, ARNALDO. Transplantation of fascia lata in repair of trachea (Trapianti liberi di fascia lata per riparazione di perdita di sostanza della trachea). *Boll. delle Malatti dell'Orecchio*, p. 177, Vol. 33, No. 8, Aug., 1915.
- 803 MARINELLI, F. Tracheotomy in croup. *Gazz. d. osped. e d. clin.*, p. 483, Vol. 36, 1915.
- 804 MILLIGAN, W. Bone impacted in left bronchus for six months; removal; recovery. *Proc. Laryngol. Section, Royal Soc. Med.*, Vol. 8, No. 6, April, 1915.
- 805 MOORHEAD, R. L. Diagnosis and removal of foreign bodies in the trachea, bronchi and esophagus. *Long Island Med. Jour.*, Oct., 1915.
- 806 PANDALAI, G. Foreign body in the trachea. *Indian Med. Gaz.*, Vol. 50, No. 7, July 15, 1915.
- 807 PRICE, H. T. Clinical findings in foreign bodies in the air passages in children. *Penn. Med. Jour.*, Dec., 1915.
- 808 RHEA, L. J. Comparative pathology of tracheal and bronchial lesions produced in man by *B. pertussis* and those produced in dogs by *B. bronchisepticus*. *Jour. Med. Research*, Vol. 32, No. 3, July, 1915.
- 809 RICHARDS, G. L. Foreign body in the lung. *Jour. A. M. A.*, July 17, 1915.

- 810 RICKETTS, B. M. Intratracheal insufflation. *Amer. Jour. Surg., Anesthesia Supp.*, p. 147, Oct., 1915.
- 811 SARTORY, A. and LASSEUR, P. Streptothricosis involving the bronchi. Recovery. *Bull. de l'Acad. de Med.*, Feb. 2, 1915.
- 812 SOERENSEN, J. Two cases of total removal of the trachea because of carcinoma. *Arch. f. Laryngol.*, Bd. 29, Heft 2, p. 188, 1915.
- 813 STEINER, R. Foreign bodies in the lower air passages and their removal (Über Fremdkörper in den unteren Luftwegen und ihre Entfernung). *Prag. med. Woch.*, No. 23, 1915.
- 814 SYMES, W. L. Note on the treatment of the symptoms arising from inhalation of irritant gases and vapours. *Brit. Med. Jour.*, July 3, 1915.
- 815 TELEGA, V. I. Foreign body several years in bronchus. *Russky Frach.*, Vol. 14, No. 28, 1915.
- 816 THORN, V. Emergency tracheotomies. *Berl. klin. Wchnschr.*, Feb. 22, 1915.
- 817 TOFT, E. A case of foreign body in the lung. *Ugeskrift f. Laeger*, No. 17, 1915.
- 818 UYEDA, I. and TAKAKI, Y. Two cases of tubercular bronchial lymph glands breaking into the trachea. *Sei-I-Kucai Med. Jour.*, No. 5, May, 1915.
- 819 VOORHEES, I. W. The importance of early tracheotomy. *Internat. Jour. Surg.*, Vol. 28, p. 110, April, 1915.
- 820 WATT, C. H. Intratracheal ether anesthesia in the surgery of the brain and spinal cord. *Jour. A. M. A.*, p. 869, Sept. 4, 1915.
- 821 WEBB, G. B., FORSTER, A. M., and GILBERT, G. B. Trachea position. *Jour. A. M. A.*, p. 1017, Sept. 18, 1915.

Esophagus.

- †822 ARROWSMITH, H. Angioneurotic edema of the esophagus. A clinical history. *THE LARYNGOSCOPE*, p. 156, March, 1915.
- 823 BARCLAY, A. E. Peptic ulcer of the esophagus. *Proc. Royal Soc. Med., Electro-therap. Section*, p. 96, May, 1915.
- *824 BEELER, R. C. Fistula of the esophagus and bronchus. Report of a case, with Roentgenologic findings. *Jour. A. M. A.*, p. 1178, Oct. 2, 1915.
- 825 BERGER, W. Case of gunshot wound of the esophagus. *Muench. med. Woch.*, p. 1557, No. 45, Nov. 9, 1915.
- §826 BOOT, G. W. Report of four cases of foreign body in the air passages or esophagus. *Jour. A. M. A.*, p. 817, March 6, 1915. *THE LARYNGOSCOPE*, p. 799, Nov., 1915.
- †827 CHAMBERLIN, W. B. Removal of an open safety-pin from the esophagus under suspension. *THE LARYNGOSCOPE*, p. 18, Jan., 1915.
- 828 CHRISTIE, G. W. Unusual case of perforated ulcer of the esophagus. *Lancet*, July 3, 1915.
- 829 COATES, G. M. and GOEPP, R. M. Case of perforation of the esophagus followed by septic infection and ending in recovery. *Penn. Med. Jour.*, May, 1915.

- 830 DAVIS, B. F. Mechanism of production of hemorrhagic erosions of esophagus and of some associated lesions in fleus. *Ann. of Surg.*, March, 1915.
- 831 DECHERD, H. B. and JONES, W. D. Removal of half-dollar and quarter-dollar from esophagus of child. *Jour. A. M. A.*, p. 2006, Dec. 4, 1915.
- 832 ELSNER, and URY, H. Roentgen differentiation of disease of esophagus and stomach. *Arch. f. Verdauungskrank.*, Vol. 21, No. 4, Aug., 1915.
- §833 FERREYRA, E. Simple method for extracting foreign bodies in the esophagus. *Sem. Med.*, Vol. 22, No. 23, June 10, 1915. *THE LARYNGOSCOPE*, p. 123, Feb., 1916.
- 834 GAUB, O. C. and JACKSON, C. Pulsion diverticulum of the esophagus. *Surg., Gyn. and Obst.*, p. 52, July, 1915.
- 835 GOFFE, E. G. L. Perforation of arch of aorta by safety pin impacted in esophagus. *Brit. Jour., Children's Dis.*, Feb., 1915.
- 836 GOYANES, DR. Removal of foreign body from esophagus (Extraccion de un cuerpo extrano del esofago). *Los Prog. de la Clin.*, p. 105, Vol. 5, 1915.
- 837 GRANT, H. H. Esophageal stenosis due to typhoid fever. *Pediatrics*, p. 258, June, 1915.
- 838 HALPENNY, J. Esophageal obstruction due to enlarged thyroid. *Brit. Jour. Surg.*, April 11, 1915.
- *839 HOPE, C. W. M. Two cases of esophageal stricture in children. *Proc. Laryngol. Section, Royal Soc. Med.*, Vol. 8, No. 6, April, 1915.
- 840 JABLONS, A. and BANOWITCH, M. M. Foreign body in esophagus. *N. Y. Med. Jour.*, June 26, 1915.
- 841 JACKSON, J. B. Case of congenital atresia of esophagus. *Mich. State Med. Soc. Jour.*, March, 1915.
- 842 JEWELL, W. H. Half-penny imbedded in the esophagus. *Proc. Roy. Soc. Med., Laryngol. Sec.*, p. 108, Vol. 8, June, 1915.
- 843 LERCHE, WM. Diseases of the esophagus. *N. Y. Med. Jour.*, June, 1915.
- 844 LINDVALL, H. Diffuse dilatation of the esophagus combined with asthma. *Hygiea*, Vol. 77, No. 15, 1915.
- 845 MADELUNG, O. Wounds of the esophagus. *Deut. med. Wchnschr.*, Jan. 28, 1915.
- †846 MCKINNEY, R. Simple inflammatory stenosis of the esophagus. *THE LARYNGOSCOPE*, p. 354, June, 1915.
- 847 McNALLY, H. H. Foreign body in the esophagus. *Canad. Med. Ass'n, Jour.*, Sept., 1915.
- 848 MERRITT, E. A. Report of case with an open safety-pin in the esophagus. *Interstate Med. Jour.*, p. 160, Feb., 1915.
- §849 MEYER, W. The early diagnosis of cancer of the esophagus. *Amer. Jour. Surg.*, July, 1915. *THE LARYNGOSCOPE*, p. 866, Dec., 1915.
- 850 MEYER, W. Further experience with resection of the esophagus for carcinoma. *Surg. Gynecol. and Obstetr.*, p. 162, Feb., 1915.
- 851 MONNIER. A case of congenital closure of the esophagus with tracheo-esophageal fistula. *Corresp. Blatt. f. Schweiz Aerzte*, No. 19, 1915.

- 852 MOORE, I. Further notes on a case of malignant stricture of the esophagus. *Proc. Roy. Soc. Med., Laryngol. Sec.*, p. 105, Vol. 8, June, 1915.
- 853 MOORE, J. L. I. Malignant stricture of the esophagus. *Proc. Royal Soc. Med., Laryngol. Section*, Vol. 9, No. 2, Dec., 1915.
- †854 MOORHEAD, R. L. Post-typhoid ulceration and stricture of the esophagus. *THE LARYNGOSCOPE*, p. 848, Dec., 1915.
- 855 NEVEU. Esophageal polyps (Deux volumineux polypes de l'oesophage). *Rev. de Laryngol., d'Otol. et de Rhinol.*, p. 306, Nos. 46-47, Nov. 30, 1915.
- 856 NUTHALL, A. W. Bullet wound of the esophagus. *Lancet*, Dec. 18, 1915.
- 857 OTRICH, G. S. Safety-pin in esophagus of young child. *J. A. M. A.*, p. 419, July 31, 1915.
- 858 RASHB, I. E. Cases of communication between esophagus and air passages. *Russky, Vrach*, Vol. 13, No. 29, 1914.
- 859 ROBERTS, W. O. Esophageal stenosis. *Amer. Jour. Surg.*, Vol. 29, May, 1915.
- 860 ROTHENBERG, F. Malformations of the esophagus (Ein casuistischer Beitrag zu den Oesophagusmissbildungen). *Inaug. Dissert., Berlin*, 1915.
- 861 SANDBERG, O. Diffuse dilatation of the esophagus. *Svenska Laeker, foer Handlingar*, Jan. 12, 1915.
- 862 SCHILLING. Gunshot wound of the esophagus. *Munch. med. Woch.*, No. 32, p. 1000, Aug. 10, 1915.
- 863 SCHMIDGALL, G. Congenital atresia of the esophagus with esophago-tracheal fistula. *Arch. f. Kinderheilk.*, Bd. 64, H. 1-2, 1915.
- 864 SCHNABEL, A. G. Stricture of esophagus. *Arizona Med., Jour.*, p. 7, Jan., 1915.
- 865 SCHREIBER, J. Sphincter mechanism in upper esophagus (Die Verschlussvorrichtung am Beginn der Speiseröhre). *Arch. f. Verdauungskrank.*, Vol. 21, No. 3, June, 1915.
- 866 SCHUETZE, J. X-ray demonstration of normal and pathological esophagus. *Berl. klin. Woch.*, June 7, 1915.
- 867 SCHUELE, A. Diffuse idiopathic dilatation of the esophagus in a man sixty-five years old. *Arch. f. Verdauungskrank.*, Vol. 21, No. 4, Aug., 1915.
- 868 SECOR, WM. L. Esophageal diverticulum; operation followed by cardiospasm. *Med. Rev. Reviews*, p. 722, Dec., 1915.
- 869 SIMPSON, W. L. Cases of removal of foreign bodies from lower respiratory tract and esophagus. *Tenn. State Med. Ass'n Jour.*, Sept., 1915.
- 870 STEPHENSEN, F. B. Extraction of coin from child's esophagus. *Colorado Med.*, May, 1915.
- 871 TOREK, F. Operative treatment of carcinoma of esophagus. *Ann. of Surg.*, p. 385, April, 1915.
- 872 UNGER, DR. Surgery of carcinoma of esophagus (Zur Chirurgie des intrathorakalen Oesophaguskarzinoms). *Arch. f. klin. Chir.* Bd. 106, H. 1, 1915.

Asthma.

- §873 BABCOCK, R. H. Nature and treatment of bronchial asthma. *Jour. A. M. A.*, June 26, 1915. *THE LARYNGOSCOPE*, p. 108, Feb., 1916.
- *874 BERTUCCI, E. A. Asthmatic attacks and inhalation of flour dust. *N. O. Med. and Surg. Jour.*, Sept., 1915.
- 875 BROWN, O. H. Some observations on bronchial asthma. *Jour. Mo. State Med. Ass'n*, p. 19, Jan., 1915.
- 876 DAVIES, B. C. An etiological study of asthma as a basis of treatment. *Interstate Med. Jour.*, p. 111, Nov., 1915.
- 877 DUDLEY, W. H. Consideration of nasal conditions causing asthma. *Jour. Ophth. and Oto-Laryngol.*, p. 14, Jan., 1915.
- 878 FREUDENTHAL, W. Direct application of electricity as well as medicaments in bronchial-asthma (Über die direkte Application des elektrischen Stromes sowie medikamentöser Mittel bei der Behandlung des Bronchial asthmas). *Arch. f. Laryngol. u. Rhinol.*, p. 443, Vol. 29, No. 3, 1915.
- 879 GORDON, A. H. Asthma. *N. E. Med. Gaz.*, p. 124, March, 1915.
- 880 HOOVER, C. F. and TAYLOR, L. Ventilatory function of lung in emphysema and asthma. *Arch. of Int. Med.*, p. 1, Jan., 1915.
- 881 ISKOWITZ, H. Immunization with autogenous proteins in bronchial asthma. *N. Y. Med. Jour.*, p. 950, Nov. 6, 1915.
- 882 JACKSON, D. E. The pharmacology of bronchial asthma. *Jour. Lab. and Clin. Med.*, p. 126, Vol. 1, No. 2, Nov., 1915.
- 883 KLENK, C. L. The treatment of ozena and asthma with autogenous vaccines. *Interstate Med. Jour.*, p. 1209, Dec., 1915.
- 884 LEVISOHN, L. A. The treatment of horse asthma by repeated injections of normal horse serum. *N. Y. Med. Jour.*, p. 901, Oct. 30, 1915.
- 885 MORRISH, R. S. Treatment of asthma and hay fever by active immunization. *Mich. State Med. Soc. Jour.*, Aug., 1915.
- 886 SAENGER, M. Asthma and fitness for military service. *Muench. med. Woch.*, No. 51, p. 1752, Dec. 21, 1915.
- 887 SICARD, M. H. Recent conceptions of bronchial asthma. *U. S. Nav. Med. Bull.*, Vol. 9, No. 3, July, 1915.

Endoscopy.

- 888 BRODER, C. B. Direct laryngobronchoscopy and esophagoscopy. *N. Y. Med. Jour.*, p. 569, Sept. 11, 1915.
- 889 CHAMBERLIN, W. B. Suspension laryngoscopy. Review of recent literature. *Interstate Med. Jour.*, p. 180, Feb., 1915.
- 890 COLLIER, C. C. Peroral endoscopy. *Jour. Ophthal. Otol. and Laryngol.*, p. 838, Oct., 1915.
- 891 CUNNINGHAM, H. M. Bronchoscopy and esophagoscopy. *Mich. State Med. Soc. Jour.*, Sept., 1915.
- 892 EZELL, H. Practical consideration of tracheo-bronchoscopy. *Tenn. State Med. Ass'n*, Jan., 1915.
- 893 FRIEDBERG, S. A. Upper bronchoscopy for a foreign body in an infant three months old. *Jour. A. M. A.*, May 15, 1915.

- 894 INGALS, E. FLETCHER. Fluoroscopic bronchoscopy with case reports and one by Dr. Stanton A. Friedberg. *Jour. A. M. A.*, p. 528, Aug. 7, 1915.
- 895 INGALS, E. F. Fluorescopic bronchoscopy (Supplemental report). *Med. Rec.*, July 10, 1915.
- 896 ISRAEL, S. Killian's suspension laryngoscopy and its application in diagnosis and surgery of the larynx. *Southwest. Jour.*, p. 136, April, 1915.
- 897 JACQUES and DURAND. Laryngoscopy (Modification à l'instrumentation laryngologique). *Rev. de Laryngol., d'Otol. et de Rhinol.*, p. 337, Nos. 48-49, Dec., 1915.
- 898 JACKSON, CHEVALIER. A fence staple in the lung; a new method of bronchoscopic removal. *Jour. A. M. A.*, June 5, 1915.
- 899 JOHNSTON, R. H. Straight, direct laryngoscopy and esophagoscopy. Report of cases. *South. Med. Jour.*, Oct., 1915.
- 900 KILLIAN, G. History of endoscopy to Bozzini's time (Zur Geschichte der Endoscopie von den ältesten Zeiten bis Bozzini). *Arch. f. Laryngol. u. Rhinol.*, p. 347, Vol. 29, No. 3, 1915.
- 901 KNEEDLER, G. C. Direct laryngoscopy. *Pittsburg Med. Jour.*, April, 1915.
- 902 LERCHE, W. Esophagoscope in diagnosis of the gullet. *Jour.-Lancet*, Oct. 1, 1915.
- 903 LYNCH, R. C. Technic of suspension laryngoscopy. *THE LARYNGOSCOPE*, p. 840, Dec., 1915.
- 904 LYNCH, R. C. Suspension laryngoscopy and its accomplishments. *Trans. Amer. Laryngol. Ass'n*, 1915, and *Ann. of Otol., Rhinol. and Laryngol.*, p. 429, Sept., 1915.
- 905 MARION, G. Extraction of projectiles in the lungs. *Presse Med.*, Sept. 16, 1915.
- 906 ORLANDINI, A. Esophagoscopy (Contributo all' esofagoscopia; Disco di metallo con punte all periferia nell' esofago). *Boll. delle Malattie dell'Orecchio, della Gola e del Naso*, May, 1915.
- 907 PHOCAS, DR. Extraction of foreign body from the lung with Bergonie's apparatus. *Bull. et mem. Soc. de Chir. de Paris*, p. 1387, Vol. 56, 1915.
- 908 REITHI, A. Hypopharyngoscopy. (Über Hypopharyngoskopie.) *Arch. f. Laryngol. u. Rhinol.*, p. 437, Vol. 29, No. 3, 1915.
- 909 SCHOONMAKER, P. Use of bronchoscope in direct examination of larynx, trachea, bronchi and esophagus. *N. Y. State Jour. of Med.*, March, 1915.
- 910 STERNBERG, W. Certain physiologic and psychologic factors to be observed in esophagoscopy and study of the stomach. *Arch. f. Verdauungskrank.*, Vol. 21, No. 3, June, 1915.
- 911 STERNBERG, W. New views in modern esophagoscope technic (Neuere Gesichtspunkte für die moderne Oesophagoskopische Technic). *Arch. f. Verdauungskrank.*, Bd. 20. H. 5-6, 1915.
- 912 STERNBERG, W. Removal of secretion by esophagoscopy (Die Sekretentfernung bei der Oesophagoskopie). *Muench. med. Woch.*, No. 14, April, 1915.

- 913 STERNBERG, W. Knee-elbow position for esophagoscopy. *Muench. Med. Wchnschr.*, p. 289, March 2, 1915.
- 914 STERNBERG, W. Simplified and improved technic for esophagoscopy. *Muench. med. Woch.*, No. 14, April 6, 1915.
- 915 VAUGHAN, H. Laryngoscopy, bronchoscopy and esophagoscopy. *N. J. Med. Soc. Jour.*, Oct., 1915.

V. DIPHThERIA AND THYROID GLAND.

Diphtheria.

- 916 AUER, E. M. Cerebral hemiplegia a sequela of diphtheria. *N. Y. Med. Jour.*, May-8, 1915.
- 917 BELL, A. J. Modern dose of diphtheria antitoxin; report of a case of laryngeal diphtheria. *Lan-Clin.*, Vol. 114, No. 5, July 31, 1915.
- 918 BERGH, E. Primary nasal diphtheria. *Monatschr. f. Ohrenheilk. u. Laryngo-Rhinol.*, p. 584, Bd. 49, H. 9 and 10, Sept.-Oct. 1915.
- 919 BERLIN, H. Bacteriologic diagnosis of diphtheria. *Deut. med. Woch.*, Vol. 41, No. 29, July 15, 1915.
- 920 BESSAU, G. and SCHWENKE, J. Local diphtheria bacillus-broth-culture reaction in man. *Monatschr. f. Kinderheilk.*, Vol. 13, No. 9, 1915.
- 921 BRUNK, O. C. Diphtheria diagnosed therapeutically. *Old Dom. Jour. Med. and Surg.*, Nov., 1915.
- 922 BUNDESEN, H. N. Schick reaction with report of eight hundred tests. *Jour. A. M. A.*, p. 1203, April 10, 1915.
- 923 CABRERA, R. Treatment of diphtheria. *Semana Medica*, No. 52, Dec. 30, 1915.
- 924 CLAYTON, T. M. Petroleum in laryngeal diphtheria. *Brit. Med. Jour.*, p. 208, Jan. 30, 1915.
- 925 CUSHING, G. M. Laryngeal diphtheria. *Jour. Ophth., Otol. and Laryngol.*, p. 65, Jan., 1915.
- *926 DUPAQUIER, E. M. Intravenous injection of diphtheria antitoxin. *N. O. Med. and Surg. Jour.*, Sept., 1915.
- 927 ENGELMANN. The question of diphtheria carriers (Zur Frage der sogenannten Diphtheria-bazillenträger). *Muench. med. Woch.*, No. 12, March 23, 1915.
- 928 EWART, W. The control of "diphtheria-bacillus carrying;" and the Jasmin oil method. *Brit. Med. Jour.*, p. 854, Dec. 11, 1915.
- 929 FERGUSON, R. R. Diphtheria treatment in general practice. *Ill. Med. Jour.*, Dec., 1915.
- 930 FOUST, E. H. How epidemic of diphtheria was checked by thorough isolation of carriers. *Mich. State Med. Soc. Jour.*, Feb., 1915.
- 931 FOX, H. Studies in diphtheroids. (1) Nature and clinical importance of pseudo-diphtheria bacilli. *Jour. Med. Research*, Vol. 32, No. 3, July, 1915.
- 932 FOX, H. Studies in diphtheroids. (2) Source of the pseudo-diphtheria forms in the body. Experiments on mutation. *Jour. Med. Research*, Vol. 32, No. 3, July, 1915.

- 933 FOX, H. Studies in diphtheroids. *Arch. of Int. Med.*, Sept., 1915.
- 934 FUCHS, DR. Diphtheritic paralysis. *Wiener klin. Woch.*, No. 21, 1915.
- 935 GEHRMANN, A. Incubation time in culture diagnosis of diphtheria. *Amer. Jour. Pub. Health*, Oct., 1915.
- 936 GRAEF, C. and GINSBERG, G. Some observations of the Schick test. *Jour. A. M. A.*, p. 1205, April 10, 1915.
- 937 GULART and FORTINEAU. An epidemic of diphtheria. *Presse med.*, May 27, 1915.
- 938 HANLY, J. F. Diphtheria: Its dissemination, prevention and cure. *Can. Lan.*, Vol. 48, No. 12, p. 645, Aug., 1915.
- 939 HARRIS, W. H. and WADE, H. W. Widespread distribution of diphtheroids and their occurrence in various lesions of human tissues. *Jour. Exp. Med.*, May, 1915.
- 940 HEWLETT, R. T. Treatment of persistent diphtheria infection by means of diphtheria endotoxin. *Lancet*, Feb. 6, 1915.
- 941 JOHNSTONE, F. M. Primary diphtheria of the trachea. *Med. Jour. Australia*, May 22, 1915.
- 942 KEILTY, R. A. A study of the occurrence of diphtheria bacilli in normal throats. *Med. Rec.*, p. 311, Vol. 88, No. 8, Aug. 21, 1915.
- 943 KELLOGG, W. H. Carrier problem in diphtheria. *Cal. State Jour. of Med.*, p. 150, April, 1915.
- 944 KINYOUN, J. J. Modification of Ponder's stain for diphtheria. *Am. Jour. of Pub. Health*, March, 1915.
- 945 KLINGER, R. and SCHOCH, E. Investigations on diphtheria. (Weitere epidemiologische Untersuchungen über Diphtherie.) *Ztschr. f. Hyg. u. Infektionskrank.*, Bd. 80, H. 1, 1915.
- 946 KNÖSEFEL, L. Serum treatment of diphtheria. *Jahrbuch f. Kinderheilk.*, No. 3, March, 1915.
- 947 KOLMER, J. A., and MOSHAGE, E. L. The Shick test. *Jour. A. M. A.*, July 10, 1915.
- *948 KOLMER, J. A., and MOSHAGE, E. L. Shick toxin reaction for immunity in diphtheria. *Am. Jour. Dis. of Children*, p. 189, March, 1915.
- 949 LESCHKE, E. Local treatment of diphtheria with tribrom-beta naphthol (Providoform). *Munch med. Woch.*, p. 1383, No. 41, Oct. 12, 1915.
- 950 LEVINSON, A. A case of diphtheritic trachea-bronchial casts in a woman 50 years of age. *Medical Record*, May 1, 1915.
- 951 LEVINSON, A. Value and limitations of the Shick diphtheria reaction in general practice. *Ill. Med. Jour.*, Dec., 1915.
- 952 LEVINSON, A., and BLATT, M. L. Studies in Schick diphtheria reactions. *Arch. of Diag.*, Vol. 8, No. 3, July, 1915.
- 953 MOFFETT, R. D. and CONRAD, A. C. Observations on the intracutaneous reaction of Schick in 455 infants and children. *Jour. A. M. A.*, p. 1010, Sept. 18, 1915.
- 954 MOOD, G. MCF. The contact in diphtheria. *Jour. So. Carolina Med. Ass'n*, Aug., 1915.
- 955 MOODY, E. E. Intradermic diphtheria toxin test. *Jour. A. M. A.*, p. 1207, April 10, 1915.

- 956 MOODY, E. E. and WOODRUFF, M. C. Schick reaction in the handling of diphtheria epidemics. *Jour. Mo. State Med. Ass'n*, p. 148, April, 1915.
- 957 NEFF, F. C. Recent methods in treating diphtheria. *Jour. A. M. A.*, p. 585, Aug. 14, 1915.
- 958 NIELSEN, R. Staining diphtheria bacilli for microscopic examination. *Hospitalstien*, p. 225, March 10, 1915.
- 959 PARK, W. H. and ZINGHER, A. Active immunization with diphtheria toxin-antitoxin, and with toxin-antitoxin combined with diphtheria bacilli. Second paper; late results. *Jour. A. M. A.*, p. 2216, Dec. 25, 1915.
- 960 PLANGE, SMITZ and RALL. Distribution of diphtheria bacilli in the body. *Muench. med. Woch.*, March 23, 1915.
- 961 RANDEGGER, G. F. Nasal diphtheria and nervous complications of diphtheria (Contributo alla clinica della difterite nasale e sulle complicanze nervose della difterite). *Rivista di Clinica Pediatrica*, July, 1915.
- 962 RIEMSDYK, M. VAN. Diphtheria and pseudodiphtheria bacilli. *Nederl. Tydschr. v. Geneesk.*, Aug. 28, 1915.
- 963 ROLLESTON, J. D. and GUNSON, E. B. Extensive occlusion of the cerebral arteries in diphtheria. *Rev. Neurol. and Psychiat.*, Aug., 1915.
- 964 SANFORD, B. J. Diphtheria, a consideration of its diagnosis and rational therapy. *Jour. of Ophth., Otol. and Laryngol.*, p. 221, March, 1915.
- 965 SAWYER, W. A. Reduction of diphtheria morbidity. *Cal. State Jour. of Med.*, p. 134, April, 1915.
- 966 SCHATZ, H. A. Primary nasal diphtheria. *N. Y. Med. Jour.*, p. 23, Jan. 2, 1915.
- 967 SCHMIDT, H. Primary nasal diphtheria in children. *Beit. zur Anat., Physiol. u. Pathol.*, Vol. 8, 1915.
- 968 SCHUMACHER, J. Detoxication of diphtheria and tetanus toxins. *Deut. med. Wchnschr.*, March 11, 1915.
- 969 SCHUERMANN, W. and PRINGSHEIM, E. G. Gins' modification of Neisser's stain for diphtheria bacilli (Zum Nachweis von Diphtherie bacillen in Original (tupferausstrich). *Med. Klinik.*, Oct. 17, 1915.
- 970 SEIDEL, O. Treatment of diphtheria (Zur Behandlung der Diphtherie). *Muench. mediz. Woch.*, p. 1209, No. 36, Sept. 7, 1915.
- 971 SMITH, W. M. Case of primary bronchial diphtheria. *Brit. Jour. Child. Dis.*, Nov., 1915.
- 972 STAMMEL, C. A. Newer antitoxin treatment of diphtheria. *Lancet*, June 12, 1915.
- 973 VEEDER, B. S. Intravenous and intramuscular administration of diphtheria antitoxin. *Jour. Mo. State Med. Ass'n*, p. 145, April, 1915.
- 974 VESSELOFF, V. S. Alkali and acid production of diphtheria bacilli. *Russky Vrach*, Vol. 14, No. 9, Feb. 28, 1915.
- 975 WADE, H. W. Bacteremia due to bacillus diphtheriae. Report of case. *Jour. of Infect. Dis.*, March, 1915.

- 976 WEAVER, G. H. and MAHER, L. K. Diagnostic value of intracutaneous injection of diphtheria toxin (Schick reaction). *Jour. of Infect. Dis.*, March, 1915.
- 977 ZINGHER, A. The Schick test. *Jour. A. M. A.*, July 24, 1915.
- 978 ZINGHER, A. and SOLETSKY, D. An economical intracutaneous method of testing the virulence of diphtheria bacilli. *Jour. Inf. Dis.*, Nov., 1915.
- 979 ZUCKERMAN, JEROME. The Schick test. One year's institutional experience with its relation to passive and active immunization in diphtheria. *N. Y. Med. Jour.*, p. 808, Oct. 16, 1915.

Thyroid Gland.

- 980 ABRAHAM, R. Medical management of exophthalmic goiter. *Med. Mag.*, p. 70, Feb., 1915.
- 981 BAINBRIDGE, W. S. Question of anesthesia in goiter operations. *Med. Press. and Circ.*, p. 265, March 17, 1915.
- 982 BAIRD, R. W. Medical side of exophthalmic goiter. *Tex. State Jour. of Med.*, Feb., 1915.
- 983 BARTON, W. M. Partial myxedema, with report of case. *Jour. A. M. A.*, p. 986, March, 20, 1915.
- 984 BEEBE, S. P. Serum treatment of hyperthyroidism. *Jour. A. M. A.*, p. 413, Jan. 30, 1915.
- 985 BENJAMIN, A. E. Goiter operations with simplified technique. *Surg., Gyn., and Obst.*, p. 111, July, 1915.
- 986 BEVERIDGE, J. W. Thyroid disease. *N. Y. Med. Jour.*, p. 1275, Dec. 25, 1915.
- 987 BIALOKUR, F. P. Exophthalmic goiter and tuberculosis. *Russky Vrach*, Nos. 26-27, 1915.
- 988 BLACKWOOD, A. L. Clinical observation of hypothyroidism. *N. E. Med. Gaz.*, May, 1915.
- 989 BLAIR, V. P. Indications for operative interference in goiter. *Jour. A. M. A.*, June 5, 1915.
- 990 BLAIR, V. P. Factors of safety in goiter operations. *Jour. Mo. State Med. Ass'n*, p. 446, Oct., 1915.
- 991 BLAU, A. A few unusual foreign bodies in the esophagus. *Ztschr. f. Laryngol.*, Bd. 7, Heft 4, p. 467, 1915.
- 992 BOYD, F. D. Treatment of exophthalmic goiter, especially in relation to control of protein metabolism. *Edin. Med. Jour.*, Vol. 15, No. 2, Aug., 1915.
- 993 BRAM, I. The non-surgical treatment of exophthalmic goiter. *N. Y. Med. Jour.*, p. 1095, Nov. 27, 1915.
- 994 BREAKSTONE, B. H. Thyroidectomy under local anesthesia. *Clin. Med.*, p. 523, June, 1915.
- 995 BROWN, A. G. Heart in diseases of the thyroid and other glands of internal secretion. *Va. Med. Semi-Monthly*, p. 469, Jan. 8, 1915.
- 996 BUTCHEL, F. C. Toxic goiter. *Denver Med. Times*, March, 1915.
- 997 BUFORD, C. G. Goiter in children. *Surg. Gynecol. and Obstetr.*, p. 35, Jan., 1915.

- 998 BULL, P. and HARBITZ, F. A case of osteomalacia associated with tumor of the parathyroid glands. *Norsk Mag. for Laegevid.*, No. 4, 1915.
- 999 BUNSTED, C. V. R. Some clinical features of exophthalmic goiter. *Med. Rec.*, March 20, 1915.
- 1000 BUNSTED, C. V. R. Some clinical features of exophthalmic goiter. *Med. Rec.*, p. 467, March 20, 1915.
- 1001 CERVO, D. Finer structure of the thyroid cell. *Policlin.*, Jan. Med. Sec., 1915.
- 1002 CLOW, F. E. Thyroid insufficiency. *Amer. Med.*, p. 887, Dec., 1915.
- 1003 COOPERNAIL, G. P. What patients with toxic goiter should do and what they should not do. *Med. Rec.*, Oct. 2, 1915.
- 1004 CORNING, E. Medical therapeutics of exophthalmic goiter. *Albany Med. Ann.*, p. 114, March, 1915.
- 1005 COTTIS, G. W. Present status of the goiter problem. *Buffalo Med. Jour.*, March, 1915.
- 1006 CRAIN, F. M. Exophthalmic goiter; report of a case. *Jour. Lancet*, Oct. 15, 1915.
- 1007 CRANK, A. C. The surgical treatment of goiter. *Jour. Mo. State Med. Ass'n.*, p. 420, Vol. 12, No. 9, Sept., 1915.
- 1008 DALE, H. H. Physiology of thyroid. *Practitioner*, Jan., 1915.
- 1009 DAUNEHL, DR. Military duty and thyrotoxia. *Deutsch. Milit. Zeit.*, Bd. 44, p. 41, 1915.
- 1010 DOUGLAS, M. Histology of thyroid in animals fed on various diet. *Jour. of Pathol. and Bacteriol.*, Jan., 1915.
- *1011 DRAPER, E. L. A case of aberrant thyroid. *Albany Med. Ann.*, p. 576, Dec., 1915.
- 1012 EASON, JOHN. A case of myxedema with tetany and suprarenal insufficiency. *Edinburgh Med. Jour.*, p. 273, Oct., 1915.
- 1013 FAWCETT, G. G., ROGERS, J., RAHE, J. M. and BEEBE, S. P. Effect of thyroid extracts on blood pressure. *Am. Jour. of Physiol.*, Jan., 1915.
- 1014 FISHER, M. R. X-ray treatment of exophthalmic goiter. *N. Y. Med. Jour.*, p. 455, March 6, 1915.
- 1015 FRIEDMAN, G. A. The influence of removal of the adrenal and one-sided thyroidectomy upon the gastric and duodenal mucosa. The experimental production of lesions, erosion and acute ulcers. *Jour. Med. Research*, May, 1915.
- 1016 FULCI, F. and GRANNUZZI, A. Regenerative power of the parathyroid. *Centralbl. f. allgem. Pathol.*, Bd. 26, 1915.
- 1017 GEYELIN, H. R. Carbohydrate metabolism in hyperthyroidism as determined by examination of blood and urine. *Arch. Internal Medicine*, Dec., 1915.
- 1018 GINSBURG, N. Surgical anatomy of upper and lower poles of thyroid with reference to thyroidectomy. *Ann. of Surg.*, March, 1915.
- 1019 GRANGER, J. P. Thyroid insufficiency in general practice. *Practitioner*, Feb., 1915.
- 1020 GULEKEE, N. Thyroid transplantation in the bone marrow. *Arch. f. klin. Chir.*, Vol. 106, No. 2, 1915.

- 1021 GURNEY, HELEN, M. Exophthalmic goiter. An abstract of the results obtained from the investigation of 93 cases. *Brit. Med. Jour.*, May 29, 1915.
- 1022 HABERER, H. VON. Failures in the treatment of Basedow's disease (Kasuistisches zur Frage therapeutischer Misserfolge bei Morbus Basedowii). *Wien. klin. Woch.*, p. 1, Bd. 28, 1918.
- 1023 HAEBERLIN, J. B. Surgical treatment of hyperthyroidism. *New York Med. Jour.*, May 29, 1915.
- 1024 HAEBERLIN, J. B. Surgical considerations of thyroid. *Ill. Med. Jour.*, Jan., 1915.
- 1025 HAGGARD, W. D. Exophthalmic goiter. *South. Pract.*, Feb., 1915.
- 1026 HANES, E. L. The parathyroids and tetany. *N. Y. Med. Jour.*, June 26, 1915.
- 1027 HARBITZ, F. On tumors of the parathyroid glands. *Jour. Med. Research*, Vol. 32, p. 361, July, 1915.
- 1028 HARDT, H. G. Treatment of hypothyroidism. *Ill. Med. Jour.*, March, 1915.
- 1029 HART, C. Exophthalmic goiter. *Med. Klin.*, p. 388, Bd. 11, 1915.
- 1030 HASKINS, J. B. Hyperthyroidism—its etiology, symptomatology and treatment. *Tenn. State Med. Ass'n*, Jan., 1915.
- 1031 HAWN, C. B. Exophthalmic goiter. *Albany Med. Ann.*, p. 111, March, 1915.
- 1032 HELLER, E. Ossification in connection with goiter. *Beitr. z. klin. Chir.*, Vol. 94, No. 2, 1915.
- 1033 HERTOEGHE, E. Thyroid insufficiency. *Practitioner*, Jan., 1915.
- 1034 HERTZ, A. F. Parathyroid glands. *Practitioner*, Jan., 1915.
- 1035 HERTZLER, A. E. Thyroidectomy under local anesthesia. *Jour. Mo. State Med. Ass'n*, p. 45, Feb., 1915.
- *1036 HESSELBERG, C. Comparison of autoplasmic and homeoplasmic transplantation of thyroid tissue in guinea-pigs. *Jour. Exper. Med.*, Feb., 1915.
- 1037 HILL, ROLAND. Surgery of goiter. *Jour. Mo. State Med. Ass'n*, p. 449, Oct., 1915.
- 1038 HIXSON, J. S. Hypo- and hyperthyroidism. *Texas Med. Jour.*, p. 223, Dec., 1915.
- 1039 HOLMGREN, I. Slight influence from thyroid preparations on growth of puppies and guinea-pigs. *Nordisch. med. Arch.*, Vol. 48, Nos. 1-2, 1915.
- 1040 HOWARD, R. M. Goiter. *Okla. State Med. Ass'n Jour.*, Feb., 1915.
- 1041 HUBERER, V. VON. Bad therapeutic results in exophthalmic goiter. *Wien. klin. Woch.*, Bd. 28, H. 1, 1915.
- 1042 HUNTER, A. and SIMPSON, S. Influence of diet of marine algae on iodine content of sheep's thyroid. *Jour. of Biol. Chem.*, Feb., 1915.
- 1043 IFLAND, H. Amyloid substance in adenoma of the thyroid (Über Amyloid in Adenomen der Schilddrüse). *Frankfurter Ztschr. f. Pathol.*, Vol. 16, No. 3, 1915.
- 1044 JAMESON, THOMAS. The indications for surgical treatment of hyperthyroidism. *N. Y. Med. Jour.*, p. 616, Sept. 18, 1915.

- 1045 JEHN, W. Removal of large intrathoracic goiters. *Deut. Zeitschr. f. Chir.*, No. 1, March, 1915.
- 1046 JOHNSTON, C. B. and BUDD, S. W. Tetany following para-thyroidectomy; report of case following partial thyroidectomy. *S. Med. Jour.*, Feb., 1915.
- 1047 KAHANE, M. Hyperthyroidism from the standpoint of military medicine. *Wt. klin. Wchnschr.*, Feb. 11, 1915.
- 1048 KASTNER, A. L. Juvenile hyperthyroidism. *Wisc. Med. Jour.*, Dec., 1915.
- 1049 KEELERT, E. Pathology of exophthalmic goiter. *Albany Med. Ann.*, p. 101, March, 1915.
- *1050 KENDALL, E. C. The isolation of the compound containing iodine which occurs in the thyroid. Its chemical nature and physiologic activity. *Jour. A. M. A.*, June 19, 1915.
- 1051 KENNEDY, D. Surgical treatment of goiter. *Med. Press and Circ.*, p. 344, April 7, 1915.
- 1052 KEY, E. Successful operation for large intra-thoracic goiter. *Nordisches med. Arch.*, Vol. 47, Surg., No. 1, 1915.
- 1053 KLINGER, R. and MONTIGEL, T. Endemic goiter. *Correspondenz-Blatt. f. Schweizer Aerzte*, April 24, 1915.
- 1054 KNUDSON, A. Physiology and pathological chemistry of exophthalmic goiter. *Albany Med. Ann.*, p. 105, March, 1915.
- 1055 KRAUS, R. Possible connection between goiter, cretinism and Chagas' disease. *Wiener klin. Woch.*, Bd. 28, No. 33, 1915.
- 1056 BOXWELL, W. Carcinoma of the thyroid. *Med. Press and Circular*, Dec. 29, 1915.
- 1057 LATHROP, W. Rectal anesthesia in thyroidectomy. *N. Y. Med. Jour.*, p. 996, Nov. 13, 1915, and *Penn. Med. Jour.*, p. 78, Nov., 1915.
- 1058 LEVI, L. Thyroid insufficiency. *Practitioner*, Feb., 1915.
- 1059 LILIENTHAL, H. Case of mediastinal thyroid removed by trans-sternal mediastinotomy. *Surg. Gynecol. and Obstetr.*, p. 589, May, 1915.
- 1060 MACALISTER, C. J. The influence of thyroid gland substance in some cases of rheumatoid disease. *Amer. Med.*, Vol. 10, p. 490, June, 1915.
- 1061 MACKENTY, J. E. Thyroid disease in its relation to laryngology. *Trans. Amer. Laryngol. Ass'n*, p. 292, 1915.
- 1062 MACY, MARY S. The overworked thyroid. *Med. Rev. of Rev.*, May, 1915.
- 1063 MACY, M. S. Some observations on clinical treatments for systemic goiter (Grave's disease). *Med. Record*, p. 431, March 13, 1915.
- 1064 MARIMON Y CARBONELL, D. J. Studies on the thyroid and para-thyroids (Estudia quirurgico del bocio y glandulas tiroideas y para-tiroideas). *Los Prog. de la Clin.*, pp. 129 and 218, May, 1915.
- 1065 MARINE, D. Etiology of goiter in brook trout. Effect of feeding with fresh and stale liver. *Jour. Exp. Medicine*, May, 1915.
- 1066 MARINE, D. Quantitative studies on *in vivo* absorption of iodine by dog's thyroid glands. *Jour. Biolog. Chem.*, Oct., 1915.

- 1067 MASSERLI, F. Treatment of goiter by continuous intestinal disinfection. *Rev. Med. de la Suisse Rom.*, Vol. 35, p. 148, 1915.
- 1068 MASSERLI, F. Etiology of endemic goiter. *Centf. f. Bakteriöl. u. Parasitol.*, Bd. 75, p. 211, 1915. *Schweiz. Rundschau f. Med.*, Vol. 15, p. 139, 1915.
- 1069 MAYER, C. P. Present status of exophthalmic goiter. *Semana Medica, Buenos Aires*, No. 11, 1915.
- 1070 MAYO, C. H. and PLUMMER, H. S. Goiter and life expectancy. *Lan.-Clin.*, June 12, 1915.
- 1071 McCARRISON, R. Distribution of goiter in India. *Indian Jour. of Med. Research*, Jan., 1915.
- 1072 McCARRISON, R. Endemic goiter. *Practitioner*, Jan., 1915.
- 1073 McCARRISON, R. Endemic goiter. *Brit. Med. Jour.*, p. 70, Jan., 1915.
- 1074 McGLANNAN, A. Some interesting experiences in thyroid surgery. *Sur., Gyn. and Obst.*, p. 696, Dec., 1915.
- 1075 MARINE, D., and FEISS, H. O. Absorption of potassium iodid by perfused thyroids and factors modifying it. *Jour. Phar. and Exp. Therap.*, Dec., 1915.
- 1076 MEYER, M. Relations between ovaries and parathyroids. *Beitr. z. klin. Chir.*, Vol. 94, No. 2, 1915.
- 1077 MILLER, S. R. and FAIRBANK, R. E. Complement fixation in thyroid diseases. *Bull. Johns Hop. Hosp.*, Vol. 26, p. 245, July, 1915.
- 1078 MILLER, JAMES. Case of myxedema. Studies from the Patholog. Dept., Univ. of Edinburgh. *Edinburgh Med. Jour.*, p. 253, Oct., 1915.
- 1079 MORI, F. Experimental production of thyroid toxic symptoms. *Frank. Zeit. f. Pathol.*, Vol. 16, No. 123, 1915.
- 1080 NOERREGAARD, K. Thyroiditis. *Ugesk. f. Leger*, p. 51, Jan. 14, 1915.
- 1081 O'DAY, J. C. Boiling water in hyperthyroidism. *N. Y. Med. Jour.*, p. 681, April 3, 1915.
- 1082 ODIN, M. Cerebellar symptoms in myxedema. *Hygiea*, Vol. 77, No. 6, 1915.
- 1083 OLSEN, E. T. Goiter, its prevalence in Chicago. *Ill. Med. Jour.*, Jan., 1915.
- 1084 O'NEILL, A. A. Hyperthyroidosis. *Ill. Med. Jour.*, Jan., 1915.
- 1805 O'NEILL, A. A. Goiter and indications for its treatment. *Ill. Med. Jour.*, Dec., 1915.
- 1086 OSOKIN, N. E. The thyroid secretion in normal and pathologic conditions. *Russky Vrach*, Vol. 14, No. 13, March 28, 1915.
- 1087 OSWALD, ADOLF. Basedow's disease (Zur Theorie des Basedow). *Muench. med. Woch.*, No. 27, p. 907, July 6, 1915.
- 1088 OSWALD, A. Constitutional iodism and its relation to exophthalmic goiter. *Rev. Med. de la Suisse Romande*, Sept., 1915.
- 1089 OUTLAND, J. H. and CLENDENING, L. The relation between the clinical symptoms and the pathologic histology of exophthalmic goiter and a plea for the more radical removal of goiter. *Med. Rev. Rev.*, p. 460, Aug., 1915.
- 1090 POPE, C. The non-surgical treatment of hyperthyroidism. *Advanced Therap.*, p. 102, March, 1915.

- 1091 POPOFF, N. M. Early signs of exophthalmic goiter. *Russky Vrach*, No. 32, 1915.
- 1092 PORTER, M. F. Boiling water injections into the thyroid gland for hyperthyroidism. *Surg., Gynecol. and Obstetr.*, p. 1, Jan., 1915.
- 1093 PORTER, M. F. Diseases of the thyroid with special reference to treatment. *Internat. Jour. Surg.*, p. 82, March, 1915, and *Detroit Med. Jour.*, p. 83, March, 1915.
- §1094 RAUTMANN, H. Pathologic anatomy of exophthalmic goiter. (Pathologisch-anatomische Untersuchungen über die Basedowsche Krankheit.) *Mitteilung. a. d. Grenzgeb. der Med. u. Chir.*, Vol. 28, No. 3, 1915. *THE LARYNGOSCOPE*, p. 577, Aug., 1915.
- 1095 RAUTMANN, H. Pathologic-anatomic studies in Basedow's disease. (Pathologisch-anatomische Untersuchungen über die Basedowsche Krankheit.) *Ztschr. f. physikal. u. diätet. Therap.*, H. 2, 1915.
- 1096 RAY, R. H. Some impressions of the goiter question. *Jour. Lancet*, Sept. 15, 1915.
- 1097 REEDER, C. L. Abnormal conditions of the thyroid gland and its relationship to neurasthenia and mental defects. *Ill. Med. Jour.*, Oct., 1915.
- 1098 RIBBERT, H. Basedow's goiter. *Virchow's Arch.*, Vol. 219, p. 246, 1915.
- 1099 ROGERS, J. Exophthalmic goiter and its treatment. *N. Y. State Jour. of Med.*, Jan., 1915.
- 1100 ROTHMANN, M. Familial occurrence of Friedrich's ataxia myxedema and dwarf growth. *Berl. klin. Wchnschr.*, Jan. 4, 1915.
- 1101 SALZMAN, S. The treatment of goiter of puberty. *Med. Rev. of Rev.*, p. 658, No., 1915.
- 1102 SANDELIN, E. Tetany after thyroid operations. *Finska Laekaresaell. Handlingar*, Vol. 57, No. 1, 1915.
- 1103 SATTERTHWAITE, T. E. Some of the modern problems in Grave's disease. *N. O. Med. and Surg. Jour.*, p. 823, April, 1915.
- 1104 SCOTT, A. C. Indications and contra-indications for surgical treatment of goiter. *Tex. State Jour. of Med.*, Feb., 1915.
- 1105 SEXTON, L. Enlarged thyroid; report of successful case of thyroidectomy. *N. O. Med. and Surg. Jour.*, No. 11, May, 1915.
- 1106 SHERRILL, J. G. Thyroid gland with special reference to goiter. *L'ville Monthly Jour. of Med. and Surg.*, p. 321, April, 1915.
- §1107 SMITH, D. Observations from twenty-four cases of exophthalmic goiter. A new symptom. *Canada Lancet*, No. 10, June, 1915. *THE LARYNGOSCOPE*, p. 176, March, 1916.
- 1108 SPITZIG, B. L. Goiter as it was known 100 years ago. *Cleveland Med. Jour.*, Jan., 1915.
- 1109 STAEMMLER, M. Post-mortem findings on the thyroid in the Charity Hospital of Berlin. *Virchow's Arch.*, Vol. 207, p. 174, 1915.
- 1110 STAEMMLER, M. Congenital goiter and its relation to disturbances of internal secretion. *Virchow's Arch.*, Vol. 219, p. 226, 1915.
- 1111 STARK, H. Indications for and results of operative treatment of exophthalmic goiter (Morbus Basedowii und Operationserfolge). *Deut. med. Woch.*, No. 28, July 8, 1915.

- 1112 STEINEGGER, A. Experiences in 400 operations on the thyroid. *Beitr. z. klin. Chir.*, Vol. 94, No. 2, 1915.
- 1113 SUMMERS, J. E. Incidence of heredity as a factor in goiter. *Ann. Surg.* Aug., 1915.
- 1114 SWAN, J. M. Cases of dysthyroidism. *N. Y. State Jour. Med.*, Vol. 15, No. 8, Sept., 1915.
- 1115 TANSINI, I. Local injections of iodine in treatment of goiter. *Gaz. degli Osped. e dell. Clin.*, Jan. 21, 1915.
- 1116 TRAVER, A. H. Surgical treatment of exophthalmic goiter. *Albany Med. Ann.*, p. 122, March, 1915, and *L'ville Monthly Jour. of Med. and Surg.*, p. 336, April, 1915.
- 1117 UNWIN, H. R. The thyroid. *Practitioner*, Nov., 1915.
- 1118 VOGEL, H. Pathological histology of the thyroid. *Virchow's Arch.*, Vol. 207, p. 204, 1915.
- 1119 WATHEN, J. R. Surgery of the thyroid gland, experimental and applied. *Lan.-Clin.*, June 5, 1915.
- 1120 WATKINS, R. Clinical reports of serum treatment for physiologically defective thyroids. *Ill. Med. Jour.*, Dec., 1915.
- 1121 WATTS, C. F. Changes in iodine content of the thyroid following changes in the blood flow through the gland. *Amer. Jour. Physiol.*, Vol. 38, No. 3, Sept., 1915.
- 1122 WEILAND, W. Treatment of exophthalmic goiter. *Therap. d. Gegenw.* No. 5, May, 1915.
- 1123 WHITMAN, L. B. Anatomy and embryology of the thyroid gland. *Chironian*, p. 321, Vol. 31, 1915.
- 1124 WILLARD, L. M. Two cases of exophthalmic goiter with ocular muscle complications. *Ophthal.*, Jan., 1915.
- 1125 WILLIAMS, L. Exophthalmic goiter. *Practitioner*, Jan., 1915.
- 1126 WILSON, D. W., STEARNS, T. and JANNEY, J. H. Excretion of acids and ammonia after parathyroidectomy. *Jour. Biol. Chem.*, Nov., 1915.
- 1127 WILSON, D. W., STEARNS, T. and THURLOW, M. D. Acid-base equilibria in blood after parathyroidectomy. *Jour. Biol. Chem.*, Nov., 1915.
- 1128 WRIGHT, H. W. Functional defects of the thyroid in relation to neurotic conditions. *Calif. State Jour. Med.*, May, 1915.
- 1129 WYETH, JOHN A. The cure of goiter by the injection of boiling water into the enlarged thyroid. *Med. Rec.*, May 29, 1915.

VI. EAR.

External Ear.

- 1130 ANZIGER, F. P. A suggestion for removing foreign bodies from the external auditory canal. *J. A. M. A.*, p. 1908, June 5, 1915.
- 1131 BANE, W. C. Eczema of the auricle due to excessive carbohydrates in diet. *Colo. Med.*, No. 6, June, 1915.
- 1132 BASILE, G. Auricular dermoid cysts (Cisti dermoidi del padiglione dell'orecchio). *Arch. ital. di Otol., Rinol. e Laringol.*, p. 478, Vol. 26, Fasc. 6, Dec., 1915.
- 1133 BEAN, R. B. Some characteristics of the external ear of American whites, American Indians, American negroes, Alaskan Eskimos and Filipinos. *Amer. Jour. Anatomy*, Vol. 18, No. 2, Sept., 1915.

- 1134 BROWN, S. H. Note concerning an artificial ear. *Ann. Otol., Rhinol. and Laryngol.*, March, 1915.
- 1135 CLAY, JOSEPH, V. F. Foreign bodies in the external auditory canals. *Jour. Ophthalm., Otol. and Laryngol.*, May, 1915.
- 1136 GUCCIARDELLO, S. Restoration of the external ear (Plastica del padiglione dell' orecchio). *Policlinico*, Vol. 22, No. 29, July 18, 1915.
- 1137 KOERNER, O. Facial paralysis following diseases of the auricle. (Über Fazialislähmung infolge von Erkrankungen der Ohrmuschel Herpes, perichondritis und Othämatom). *Arch. f. Ohrenheilk.*, Bd. 72, H. 4, May, 1915.
- 1138 MALAN, A. Tuberculoma of lobe of ear. *Boll. delle mal. dell' Orecchio*, p. 1, Jan., 1915.
- 1139 MCKENZIE, DAN. Herpes zoster oticus combined with recurrent laryngeal paralysis. *Jour. Laryngol., Rhinol. and Otol.*, p. 339, Vol. 30, No. 9, Sept., 1915.
- 1140 SARENSON, T. Periauricular abscess from scarlet fever. *Ugeskrift for Læger*, Bd. 77, p. 175, 1915.
- 1141 SCHUMACHER, H. Loose exostosis in bony auditory canal with its cutaneous pedicle. *Monatschr. f. Ohrenh.*, Heft 1, p. 14, 1915.
- 1142 SOERENSEN, S. T. Phlegmons around the ear in scarlet fever. *Ugesk. f. Læger*, Feb. 4, 1915.
- 1143 TAKANARITA, W. Post-mortem findings in a case of congenital unilateral atresia of the auditory canal. *Arch. f. Ohrenh.*, Bd. 96, Heft 3-4, p. 186, 1915.
- 1144 TWEEDIE, A. R. Herpes zoster oticus. *Jour. Laryngol., Rhinol. and Otol.*, July, 1915.

Tympanic Cavity.

- §1145 ADAM, J. Infection of middle ear with Vincent's organisms. *Brit. Jour. Children's Dis.*, p. 40, Feb., 1915. THE LARYNGOSCOPE, p. 52, Jan., 1916.
- 1146 ALEXANDER, G. J. Nasal obstruction as the cause of diseases of the middle ear. *Jour. Ophthalm., Otol. and Laryngol.*, p. 36, Jan., 1915.
- 1147 AGAZZI, B. Atrophy of tensor tympani muscle; remarks on its physiology (Un caso di grave atrofia del tensor tympani con alcune considerazioni sulla fisiologia di questo muscolo). *Arch. ital. di Otol., Rinol. e Laryngol.*, p. 457, Vol. 26, Fasc. 6, Dec., 1915.
- 1148 AYNESWORTH, H. T. Chronic purulent otitis media and complications; report of cases. *Texas State Jour. of Med.*, Aug., 1915.
- 1149 BACON, GORHAM. The difficulties of closing a perforation of the drum membrane in certain cases of acute suppurative otitis media. *Ann. Otol., Rhinol. and Laryngol.*, March, 1915.
- 1150 BECK, O. Acute otitic dyscrasia. *Ztschr. f. Ohrenh.*, Bd. 72, Heft 2, p. 97, 1915.
- 1151 BELLOW, H. P. Internal remedies in acute inflammatory conditions of the middle ear. *Jour. Ophthalmol., Otol. and Laryngol.*, p. 996, Dec., 1915.

- 1152 BLEDSOE, R. W. Treatment of middle ear through Eustachian tube. *Ky. Med. Jour.*, Jan. 1, 1915.
- 1153 BLEDSOE, R. W. Acute purulent otitis media complicated by mastoiditis and temporo-sphenoidal abscess. *South. Med. Jour.*, Vol. 8, No. 9, Sept., 1915.
- 1154 BORDEN, C. R. C. Systemic infection of middle ear origin in the exanthemata. *Ann. Otol., Rhinol. and Laryngol.*, March, 1915.
- 1155 BOETTGER, K. Technic of Politzeration (Zur Technik des Politzer-verfahrens). *Muench. mediz. Woch.*, p. 1215, No. 36, Sept. 7, 1915.
- 1156 CHARLTON, C. C. The treatment of acute suppurative otitis media in infants and children. *N. J. Med. Jour.*, Vol. 12, p. 221, May, 1915.
- 1157 CLERC, L. Observations on the middle ear glands of some mammifers (Osservazioni su le ghiandole dell'orecchio medio di alcuni mammiferi). *Arch. ital. di otol., rinol. e laringol.*, Vol. 26, No. 3, p. 177, June, 1915.
- 1158 CLERC, L. Anatomy of the fetal middle ear (Contributo all'anatomia dell'orecchio medio fetale). *Arch. ital. di otol., rinol. e laringol.*, p. 269, Vol. 26, No. 4, Aug., 1915.
- †1159 COCKS, G. H. and DWYER, J. G. The isolation and cultivation of the tubercle bacillus from the discharging ear in cases of chronic purulent otitis media (Preliminary report). *THE LARYNGOSCOPE*, March, 1915.
- *1160 EMERSON, W. R. P. Acute otitis media in infancy and early childhood; avoidable mistakes in diagnosis, prevention and treatment. *Bost. Med. and Surg. Jour.*, Oct. 21, 1915.
- *1161 FRASER, J. S. and OWEN, J. L. A case of sub-acute purulent otitis media labyrinthitis and purulent leptomeningitis due to a capsulated streptococcus: spontaneous recovery. *Edinburgh Med. Jour.*, p. 269, Oct., 1915.
- 1162 FRASER, J. S. and DICKIE, J. K. M. A reconstruction model of the right middle and inner ear. *Jour. Laryngol., Rhinol. and Otol.*, p. 457, Dec., 1915.
- 1163 GIVENS, L. S. Complications of middle ear suppuration. *Kentucky Med. Jour.*, Dec., 1915.
- *1164 KOBRAK, F. Prognosis and treatment of severe acute middle ear disease. *Therap. der Gegenw.* Bd. 55, H. 6, June, 1915.
- 1165 KOERNER, O. Facial paralysis following operations on the middle ear and temporal bone (Über Fazialislähmung infolge von Operationen im Mittelohr und am Schläfenbein). *Arch. f. Ohrenheilk.*, Bd. 72, H. 4, May, 1915.
- 1166 LANG, J. Acute middle ear suppuration after removal of nasopharyngeal tumor with the galvanocautery (Über akute eitrige Mittelohrentzündung nach der Entfernung der Nasenrachentumoren mittelst der galvanokaustischen Schlinge). *Monatschr. f. Ohrenheilk.*, p. 289, Vol. 46, 1915.
- †1167 LARGE, S. H. Gold-platinum inserted in middle ear for adhesive processes in the middle ear. *THE LARYNGOSCOPE*, p. 370, June, 1915.

- 1168 LEWIS, E. R. Acute suppurative otitis media with unusual complications. *Trans. Amer. Otol. Soc.*, p. 561, Vol. 8, part 3, 1915.
- 1169 LEWY, A. The treatment of acute otitis media by the general practitioner. *Clinique*, May, 1915.
- 1170 LILLIE, H. J. Fulminating otitis media, mastoiditis, extensive sigmoid sinus thrombosis, ligation of internal jugular vein, recovery. *Mich. State Med. Soc. Jour.*, March, 1915.
- 1171 LINCK, A. Headache from unsuspected middle ear disease. *Therap. d. Gegenwart*, March, 1915.
- 1172 LOMBARD, BLOCH, A. and MOULONGUET, A. Abscess of frontal lobe on opposite side to chronic, suppurative otitis (Un cas d'abcès du lobe frontal du côté opposé à une otite, suppurée chronique). *Ann. des mal. de l'oreille*, p. 749, Vol. 40, No. 8, 1914-1915.
- 1173 STUART-LOW, W. Non-suppurative forms of middle-ear deafness. *Pract.*, p. 503, April, 1915.
- 1174 LUNGHINI, O. Comparative anatomy of the middle-ear. (Contributo all'anatomia comparata dell'orecchio medio.) *Arch. ital. di Otol., Rinol. e Laringol.*, p. 537, V. 26, Fasc. 6, Dec., 1915.
- 1175 MACFARLAN, D. So-called stock vaccines and their value in suppurative middle-ear disease. *Jour. Ophthal., Otol. and Laryngol.*, p. 691, Aug., 1915.
- 1176 MACKENZIE, G. W. Acute secretory catarrh of the middle-ear. *The Hahnemannian Monthly*, Oct., 1915.
- 1177 MACNAB, J. C. G. Two cases of neuritis of the lesser occipital nerve following acute otitis media suppurativa. *Jour. Laryngol., Rhinol. and Otol.*, p. 335, V. 30, No. 9, Sept., 1915.
- 1178 MAYER, OTTO. Transplantation of skin flaps in complete opening of the middle-ear cavity. (Eine Methode der Transplantation von Hautlappen bei der Totalaufmeisselung der Mittelohrräume.) *Arch. f. Ohrenheilk.*, V. 98, No. 2-3, p. 146, Aug. 1915.
- 1179 MCKENZIE, D. Otogenic pharyngeal abscess. *Jour. of Laryngol.*, p. 12, Jan., 1915.
- 1180 MCKINNEY, R. Epidemic otitis media; study of its manifestations. *Tenn. State Med. Ass'n Jour.*, Aug., 1915.
- 1181 MURPHY, J. W. Acute middle-ear inflammations. *Jour.-Lancet*, Nov. 6, 1915.
- 1182 OPPENHEIMER, S. Metastatic complications of suppurative otitis media with especial reference to secondary inflammatory processes in the joints. *N. Y. Med. Jour.*, p. 49, Jan. 9, 1915.
- 1183 SAWREY, E. R. Notes on the causation and diagnosis of suppurative otitis. *Med. Jour. Austral.*, V. 1, p. 286, 1915.
- 1184 SCHENCK, H. G. Pneumo-massage in diseases of the middle-ear. *Jour. Ophth., Otol. and Laryngol.*, p. 342, April, 1915.
- *1185 SCHERER, F. and KUTVIRT, O. Otitis media in relation to infant pathology. (Beziehung der Mittelohrentzündung zu den Krankheiten des Säuglingsalters.) *Jahrbuch f. Kinderheilk.*, Bd. 82, H. 2, Aug., 1915.
- 1186 SHAMBAUGH, G. E. Clinical problems in connection with some of the complications of acute otitis media. *Ill. Med. Jour.*, Sept., 1915.

- *1187 SMITH, C. H. Acute otitis media. *N. Y. Med. Jour.*, p. 1951, Nov. 20, 1915.
- *1188 SPINETTO, D. J. Radium treatment in diseases of the middle and internal ear. (Radium terapia en algunas afecciones del oído.) *Semana Medica*, Aug. 5, 1915.
- 1189 STICKNEY, O. D. Complications of suppurative inflammation of the middle-ear and mastoid. *Hahnemanian Monthly*, p. 199, March, 1915.
- 1190 TENNER, A. S. Otitic pneumococcic bacteremia. *Jour. A. M. A.*, July 17, 1915.
- 1191 TOMSON, J. and FRASER, J. S. A case of broncho-pneumonia, purulent otitis media and interna, and leptomeningitis. Post-mortem: microscopic examination of the ear. *Edinburgh Med. Jour.*, p. 265, Oct., 1915.
- 1192 TORRINI, UMBERTO L. Epitympanic infection; paralysis of external oculomotor. (Infezioni delle cellule squamose epitimpaniche. Paralisi dell'oculomotore esterno.) *Boll. delle Mal. dell'Orecchio*, May, 1915.
- 1193 TRIBLE, G. B. Suppurative otitis media and its complications. *Milit. Surg.*, June, 1915.
- *1194 TURNER, A. LOGAN, and FRASER, J. S. Tuberculosis of the middle-ear cleft in children: a clinical and pathological study. *Jour. Laryngol., Rhinol. and Otol.*, June, 1915.
- 1195 WALES, E. DE WOLFE. Treatment of chronic suppurative otitis media, with report of cases. *Trans. Amer. Otol. Soc.*, p. 534, V. 8, Part 3, 1915.
- 1196 WEST, C. E. Tuberculosis of the auditory apparatus. *Brit. Jour. Child. Dis.*, June, 1915.
- 1197 WHALE, H. LAWSON. Septicemia of otitic origin. *British Med. Jour.*, pp. 723-24, April 24, 1915.
- 1198 WILKINSON, O. The treatment of acute otitis media. *Virginia Med. Semi-Monthly*, p. 292, Sept. 24, 1915.
- *1199 WILLIAMS, C. E. A new treatment for middle-ear disease. *Jour. Ophthalm., Otol. and Laryngol.*, June, 1915.
- 1200 WINCKLER, E. Infection with streptococcus mucosus. *Arch. f. Ohrenh.*, Bd. 96, Heft 3-4, p. 193, 1915.
- 1201 WOODSIDE, E. E. Acute infantile otitis. *Ill. Med. Jour.*, Feb., 1915.

Ossicles.

- 1202 AGAZZI, B. Function of the tensor tympani muscle. (La funzione del "Musculus tensor tympani.") *Arch. ital. di otol., rinol. e laringol.*, V. 26, No. 3, p. 224, June, 1915.
- †1203 BECK, J. C. The Roentgenographic diagnosis in otosclerosis. Preliminary report. *THE LARYNGOSCOPE*, p. 154, March, 1915.
- 1204 BOOR, G. W. The diagnosis of oto-sclerosis. *Ill. Med. Jour.*, Sept., 1915.
- 1205 BRYANT, W. S. The etiology of otosclerosis. *Jour. Laryngol., Rhinol. and Otol.*, p. 389, V. 30, No. 10, Oct., 1915.
- 1206 HARRIS, T. J. A brief consideration of certain recent views regarding otosclerosis. *New Albany Med. Herald*, Aug., 1915.

Eustachian Tube.

- 1207 COULTER, C. F. and PIERCE, C. H. The bacteriology of the Eustachian tube. *Jour.-Lan.*, V. 35, p. 177, 1915.
- 1208 DENMAN, I. O. Pharyngo-Eustachian infections. *Jour. of Ophth., Otol. and Laryngol.*, p. 198, March, 1915.
- 1209 HAYS, H. Eustachian salpingitis. *Med. Times*, V. 43, p. 320, 1915.
- *1210 JACOD, DR. Sarcoma of the Eustachian tube. (Sur les sarcomes de la trompe d'eustache cartilagineuse.) *Rev. de laryngol., d'otol. et de rhinol.*, p. 169, No. 34-35, Aug., 1915.
- 1211 MINK, P. J. Massage of the Eustachian tube. (Die von mir geübten Striche bei der Massage der Ohrtrumpete.) *Zeitschr. f. Ohrenheilk.*, Bd. 72, H. 3, p. 177 and Bd. 73, H. 1, p. 53, July, 1915.
- 1212 SITES, J. M. Acute tubo-tympanic catarrh. *W. Va. Med. Jour.*, Jan., 1915.
- 1213 SOWERS, A. Pneumomassage of the Eustachian tube and tympanum. *Jour. Ophthal., Otol. and Laryngol.*, p. 856, Oct., 1915.
- 1214 STIMSON, G. W. Breakage and removal of Eustachian applicator. *Jour. A. M. A.*, p. 430, Jan. 30, 1915.
- †1215 YANKAUEB, SIDNEY. Report of a collective investigation on the curettement of the Eustachian tube in chronic aural suppuration. *THE LARYNGOSCOPE*, p. 675, Oct., 1915.

Labyrinth.

- 1216 ALBRECHT, W. Vestibular irritability in deaf-blind individuals. *Beitr. zur Anat., Physiol. Pathol. u. Therap.*, Bd. 8, p. 192, 1915.
- 1217 ALT, F. Neuritis of auditory nerves following carbon monoxide gas intoxication. *Arch. f. Ohrenh.*, Bd. 96, Heft 3-4, p. 183, 1915.
- 1218 ANDRADE, C. DE. Keratosis of ear-shell. *Brazil Medico*, No. 18, 1915.
- 1219 BACHRACH. Intensity of sound perception at various times in the day. *Zeitschr. f. Sinnesphysiol.*, V. 49, No. 2, 1915.
- 1220 BALENGER, W. L. Labyrinthitis with illustrative cases. *New Albany Med. Herald*, July, 1915.
- 1221 BARRIE, T. S. A case of rotatory nystagmus with recovery under optical treatment. *Brit. Med. Jour.*, p. 469, Sept. 25, 1915.
- 1222 BECK, K. Fixation of labyrinth during life for histological examination. *Ztschr. f. Ohrenheilk.*, Bd. 72, H. 2, p. 113, 1915.
- 1223 BERITOFF, J. S. On the reciprocal innervation in tonic reflexes from the labyrinths and the neck. *Jour. Physiology*, V. 49, No. 4, May 12, 1915.
- 1224 BERRY, G. Labyrinthitis following operation for atresia. *Boston Med. and Surg. Jour.*, May 13, 1915.
- 1225 BING, A. Tuning-fork tests and their diagnostic value. *Arch. f. Ohrenh.*, Bd. 96, Heft 3-4, 1915.
- 1226 BUYS and HENNEBERT. Galvanic destabilization. (De la destabilisation galvanique.) *Rev. de Laryngol., d'Otol. et de Rhinol.*, p. 245, No. 40-41, Oct. 15, 1915.
- 1227 CLAY, JOSEPH V. F. A case of bilateral auditory neuritis due to poisoning by illuminating gas. *Jour. Ophth., Otol. and Laryngol.*, May, 1915.

- 1228 COLE, W. F. Some investigations concerning mysterious form and function of internal and middle-ear. *Texas State Jour. Med.*, Oct., 1915.
- 1229 DABNEY, V. Bell's palsey and neuritis of the eighth and twelfth nerves following salvarsan injection. *Interstate Med. Jour.*, p. 478, May, 1915.
- †1230 DENCH, EDWARD B. The labyrinth operation; the frequency with which it is demanded as determined by hospital and private statistics, extending over a period of ten years. *THE LARYNGOSCOPE*, p. 556, Aug., 1915.
- 1231 EDGAR. Clinical investigations on diseases of the ear, especially of the internal ear, in diabetes. (Klinische Untersuchungen über die Erkrankungen des Gehörorgans bei Diabetes Mellitus, mit besonderer Berücksichtigung der Erkrankungen des inneren Ohres.) *Monatschr. f. Ohrenheilk.*, Bd. 49, H. 4-5, 1915.
- 1232 ELLIS, A. W. M., and SWIFT, H. Involvement of the eighth nerve in syphilis of the central nervous system. *Jour. A. M. A.*, p. 1471, May 1, 1915.
- 1233 EPPLEN, F. Cerebellar localization with reference to nystagmus and vertigo as localizing signs. *Northwest Med.*, Nov., 1915.
- 1234 FOSTER, J. H. Two cases of labyrinth disease with operations. *Texas State Jour. Med.*, Nov., 1915.
- 1235 FOWLER, E. P. The origin of labyrinthine rest-tone. *Jour. A. M. A.*, Jan. 9, 1915.
- *1236 FOWLER, E. P. Origin of labyrinthine rest-tone. *Jour. A. M. A.*, p. 118, Jan. 9, 1915.
- 1237 GEZES, R. Rinné's test. (La pratique due Rinné Avec quels diapasons faut-il pratiquer le Rinne?) *Rev. de Laryngol., d'Otol. et de Rhinol.*, p. 297, Nos. 46-47, Nov. 30, 1915.
- 1238 GIRARD. Circumscribed labyrinthitis. (Un cas de labyrinthite circonscrite au limaçon avec paralysie faciale; évidemment pétromastoidien; guérison.) *Rev. de Laryngol., d'Otol. et de Rhinol.*, p. 304, Nos. 46-47, Nov. 30, 1915.
- *1239 GOT, A. War deafness from lesions of the internal ear. (Sourdité de guerre des lésions d l'oreille interne.) *Gaz. hebdomadaire des sciences med.* July, 1915.
- 1240 GRADENIGO, G. Anomalies of sound perception. (Sulla anomalia di percezione dei suoni.) *Arch. ital. di otol., rinol. e laringol.*, V. 26, No. 1, p. 43, Jan., 1915.
- †1241 GUGGENHEIM, LOUIS K. Bárány's cerebellar localization. *THE LARYNGOSCOPE*, p. 568, Aug., 1915.
- 1242 HARDESTY, I. Model to illustrate probable action of the tectorial membrane. *Amer. Jour. Anatomy*, Nov., 1915.
- 1243 HARDESTY, I. On the proportions, development and attachment of the tectorial membrane. *Amer. Jour. Anatomy*, p. 1, V. 18, No. 1, July 15, 1915.
- *1244 HASTINGS, H. Syphilis of the internal ear (hereditary). *Jour. A. M. A.*, p. 607, Aug. 14, 1915.
- 1245 HEAD, J. W. Important symptoms which require labyrinthine operation; report of cases. *Texas State Jour. Med.*, Nov., 1915.

- *1246 HIRSCH, C. Isolated vestibular neuritis following typhoid inoculation. *Deutsch med. Woch.*, Aug. 19, 1915.
- 1247 HOFFMAN, R. Detonation injuries of the internal ear. (Detonations labyrinthosen.) *Munch. med. Woch.*, p. 1269, No. 34, Sept. 14, 1915.
- 1248 HOLMES, E. M. Space sense and the labyrinth. *Maine Med. Ass'n Jour.*, Dec., 1915.
- †1249 IDE, C. EDWARD. A case of suppurative labyrinthitis, meningismus, irritation of the Gasserian ganglion. *THE LARYNGOSCOPE*, p. 129, March, 1915.
- 1250 KELLEY, I. D. JR. Nystagmus. *Interstate Med. Jour.*, p. 891, V. 22, No. 9, Sept., 1915.
- 1251 LANG, JAX. Can we determine by differences in the irritability of both vestibular labyrinths which side is involved in non-suppurative labyrinthine disease? (Kann man in den Fällen von nichteitrigen Labyrinthkrankungen bei konstatierter Differenz der Erregbarkeit beider Vestibularlabyrinthe eventuell auch beider Vestibularnerven feststellen, welche Seite betroffen ist und ob es sich dann um eine Erhöhung oder Abnahme der Erregbarkeit handelt? *Passow's Beiträge*, p. 193, V. 8, No. 4, Aug. 12, 1915.
- 1252 LORENZEN, P. Functional disturbances in gait in the young. (Infantile funktionelle Gangforstyrrelser.) *Hospitalstidende*, Nov. 3, 1915.
- 1253 MACKENZIE, G. W. The practical value of the labyrinth tests. *Jour. Ophthal., Otol. and Laryngol.*, p. 787, Sept., 1915.
- 1254 MACKENZIE, G. W. The galvanic test as an aid in the differential diagnosis of inner-ear from eighth nerve affections. *Jour. Ophthal., Otol. and Laryngol.*, p. 858, Oct., 1915.
- 1255 MAXEY, E. E. Case of syphilis of internal and middle-ear. *N. W. Med.*, Jan., 1915.
- †1256 MAYBAUM, J. L. Labyrinthine fistula; operation; recovery. *THE LARYNGOSCOPE*, p. 774, Nov., 1915.
- 1257 MAYER, O. Hearing capacity with suppuration of the labyrinth. *Monatschr. f. Ohrenh.*, Heft 1, p. 1, 1915.
- 1258 MEYER, DR. Acoustic trauma. (Das akustische Trauma.) *Arch. f. Ohrenheilk.*, V. 98, No. 2-3, p. 152, Aug., 1915.
- *1259 MEYERS, I. LEON. Galvanometric studies of the cerebellar function. *Jour. A. M. A.*, p. 1348, Oct. 16, 1915.
- 1260 MUNDT, G. H. Syphilis of the internal ear. *Ill. Med. Jour.*, Sept., 1915.
- 1261 NAKAMURA, Y. Experimental study of the effects of ethyl and methyl alcohol upon the acusticus. *Beitr. z. Anat., Physiol., Pathol. u. Therap.*, p. 41, 1915.
- 1262 NICCOL, W. A pedigree of hereditary nystagmus. *Ophthalmoscope*, May, 1915.
- 1263 NOEHTE. Nystagmus after injury of foot of second frontal convolution. Report of nine cases. *Deut. mediz. Woch.*, Oct. 7, 1915.
- 1264 PANSE, R. Remarks on testing the labyrinth (Einige Bemerkungen zur Prüfung des Labyrinthes). *Zeitsch. f. Ohrenheilk.*, p. 161, Vol. 73, No. 12, 1915.

- 1265 PROVERA, GIOVANNI. Considerations on caloric nystagmus and Brünings's oto-calorimetry (Considerazioni sul nistagmo calorico e sull'otocalorimetria alla Brünings's). *Boll. delle malat. dell'orecchio*, Vol. 33, No. 2, p. 30, February, 1915, and March, 1915.
- 1266 PUSATERI, S. Inequality of the pupils in suppurative labyrinthitis (L'anisocoria nelle piolabirintiti). *Arch. ital. di Otol., Rinol. e Laringol.*, p. 467, Vol. 26, fasc. 6, Dec., 1915.
- 1267 RANDALL, B. ALEX. The ear-tests of Barany in locating cerebellar and other encephalic lesions. *Trans. Amer. Otol. Soc.*, p. 511, Vol. 8, part 3, 1915.
- 1268 RHESE, DR. Origin and clinical significance of vestibular unbalance (Die Entstehung und klinische Bedeutung der vestibulären Fallbewegungen). *Zeitsch. f. Ohrenheilk.*, p. 94, Vol. 73, No. 12, 1915.
- *1269 SCHMIEGELOW, E. Translabyrinthine removal of acoustic tumors (Beitrag zur translabyrinthären Entfernung der Akustikustumoren). *Ztschr. f. Ohrenheilk.*, Bd. 73, H. 1, p. 1, July, 1915.
- 1270 SCHMIEGELOW, E. Removal through petrous bone of tumor involving the auditory nerve. *Hospitalstidende*, p. 201, March 3, 1915.
- 1271 SCHROEDER, P., and HINSBERG, V. The specific action of quinin and salicylic acid on the spiral ganglion. (Zur Frage der spezifischen Wirkung von Chinin und Salizylsäure auf das Ganglion spirale). *Zeitsch. f. Ohrenheilk.*, p. 65, Vol. 73, No. 12, 1915.
- 1272 SHAMBAUGH, G. E. Focal infection in the etiology of labyrinth disease. — *Trans. Amer. Otol. Soc.*, 1915, *Ann. of Otol., Rhinol. and Laryngol.*, p. 479, Sept., 1915.
- 1273 SHINKLE, C. E. The utility of the Barany functional tests. *Lancet-Clin.*, Sept. 18, 1915.
- 1274 SIMON, R. M. Case of Meniere's disease. *Brit. Med. Jour.*, p. 282, Feb. 13, 1915.
- 1275 STEFANINI, A. Experimental confirmation of the Cotugno-Helmholtz theory on the perception of sound (Conferma sperimentale della teoria di Cotugno-Helmholtz sulla percezione dei suoni). *Arch. ital. di otol., rinol. e laringol.*, Vol. 26, No. 1, p. 34, January, 1915.
- 1276 STEFANINI, A. Sound perception (La teoria della risonanza per la percezione dei suoni). *Arch. ital. di otolog.*, p. 364, Vol. 26, No. 5, Oct., 1915.
- 1277 WEINBERG, MAX. Blood examinations in otitis media, especially in the first stage (Blutuntersuchungen bei Otitis media, besonders in den ersten Stadien). *Inaug. Dissert.*, Heidelberg, Feb., 1915.
- 1278 WILLCUTT, G. H. Disturbances of the acoustic nerve in the early stages of syphilis. *Jour. A. M. A.*, No. 602, Aug. 14, 1915.
- 1279 WILSON, J. G., and PIKE, F. H. Mechanism of labyrinthine nystagmus and its modifications by lesions in cerebellum and cerebrum. *Arch. of Int. Med.*, Jan., 1915.
- *1280 WILSON, J. G., and PIKE, F. H. The differential diagnosis of lesions of the labyrinth and of the cerebellum. *Jour. A. M. A.*, p. 2156, Dec. 18, 1915.

- 1281 WILSON, J. A. Nystagmus and allied conditions. *Lancet*, Oct. 23, 1915.
- 1282 WILSON, J. GORDON. The differential diagnosis of lesions of the labyrinth and cerebellum. *Trans. Amer. Otol. Soc.*, p. 470, Vol. 8, Part 3, 1915.
- 1283 WILSON, J. G., and PIKE, F. H. Vertigo. *Jour. A. M. A.*, p. 561, Feb. 13, 1915.
- 1284 WINTERMUTE, G. P. Auditory neuritis. *Jour. A. M. A.*, p. 603, Aug. 14, 1915.
- 1285 ZAUGE, J. Psychic origin of functional nervous disturbance in the ear. (Hysterische Funktionsstörungen des nervösen Ohrapparates im Kriege). *Munch. med. Woch.*, No. 28, July 13, 1915.
- 1286 ZANGE, J. Injuries to the nervous apparatus of the ear in war (Organische Schädigungen des nervösen Ohrapparates im Kriege). *Munch. med. Woch.*, p. 1091, No. 32, Aug. 10, 1915.

Defects of Hearing.

- 1287 ADAMS, MARY B. Visits to the homes of pupils in the Horace Mann school in Boston. *Volta Review*, July, 1915.
- †1288 BALDWIN, R. N. Responsibilities of the community in the education of the deaf. *THE LARYNGOSCOPE*, April, 1915.
- 1289 BARNES, F. G. Seaside school and camp for the deaf. *Volta Rev.*, June, 1915.
- 1290 BARNES, F. G. Teacher question. *Teacher of Deaf*, p. 22, Feb., 1915.
- 1291 BEMIS, L. M. Yours for the deaf in China. *Volta Rev.*, p. 18, Jan., 1915.
- 1292 BOLTON, T. L. Psychology of education of deaf-blind. *Am. Ann. of Deaf*, p. 222, May, 1915.
- 1293 BRUHN, MARTHA E. The hard-of-hearing adult. *Volta Rev.*, Vol. 17, No. 10, Oct., 1915.
- 1294 BRUHN, MARTHA E. The Müller-Walle method of lip-reading. *Volta Rev.*, Vol. 17, No. 8, August, 1915.
- 1295 BUELL, E. M. Reading for the deaf. *Am. Ann. of Deaf*, p. 1, Jan., 1915.
- 1296 CARTER, MISS A. E. Teaching speech to the deaf. *Volta Rev.*, Vol. 17, No. 8, August, 1915.
- 1297 CASE, L. E. Conversation-class work in lip-reading. *Volta Rev.*, p. 2, Jan., 1915.
- 1298 CASTELLANI, L. Auditory re-education in progressive deafness (Note critiche sulla rieducazione uditiva nei casi di sordità progressiva). *Arch. ital. di Otol., Rinol. e Laringol.*, Vol. 26, April, 1915.
- 1299 CASTEX, A. Deafness from shell explosion. *Bull. de l'Academie de Med. de Paris*, Nov. 16, 1915.
- 1300 CLARK, LLOYD H. The prevention of deafness. *Jour. Ophth., Otol. and Laryngol.*, May, 1915.
- 1301 DROUOT, E. Great European war and schools for the deaf in Germany. *Am. Ann. of Deaf*, Jan. and May, 1915.
- 1302 DUFF, J. Advantages of an oral day school for the deaf. *Volta Rev.*, p. 138, April, 1915.

- 1303 DUFOUR, C. R. A plea for co-operation of physicians to prevent deafness. *Va. Med. Semi-Monthly*, p. 34, April 23, 1915, and *Volta Rev.*, July, 1915.
- 1304 DYER, H. L. Vocational suggestions for the adult deaf. *Volta Rev.*, p. 9, Jan., 1915.
- †1305 GOLDSTEIN, M. A. Medical co-operation in the education of the deaf. *THE LARYNGOSCOPE*, April, 1915.
- 1305 GRUVER, E. A. Correlation of liberal and vocational education for the deaf. *Am. Ann. of Deaf*, p. 165, March, 1915.
- 1306 GRUVER, E. A. Correlation of liberal and vocational education for the deaf. *Am. Ann. of Deaf*, p. 165, March, 1915.
- 1307 HAENLEIN. Lip-reading for deaf soldiers. (Fürsorge für ertaubte und Schwerhörige Krueger.) *Med. Klin.*, May 30, 1915.
- 1308 HARRIS, T. J. The re-education of the deaf ear. *Volta Rev.*, July, 1915.
- 1309 HAYS, H. Can deafness be prevented? *Amer. Med.*, p. 122, Feb., 1915, and *Volta Rev.*, Vol. 17, No. 10, Oct., 1915.
- 1310 HAYS, H. Deafness versus stupidity. *Med. Times*, Mar., 1915.
- †1311 HILLIARD, ETHEL M. Practical demonstration with pupils of the Central Institute for the Deaf. *THE LARYNGOSCOPE*, April, 1915, and *Volta Rev.*, Sept., 1915.
- 1312 HIRSH, A. B. Chronic infective deafness relieved by the static brush discharge (Yates method). *Adv. Therap.*, p. 300, June, 1915.
- 1313 HUBBARD, THOMAS. Nerve deafness and mumps. *Trans. Amer. Otol. Soc.*, p. 451, Vol. 8, Part 3, 1915.
- 1314 HUBBY, L. M. Severely impaired audition in childhood. *Med. Record*, p. 1089, Dec. 25, 1915.
- 1315 HURD, ANNA M. The Montessori method applied to deaf children. *Volta Rev.*, June, 1915.
- †1316 KERRISON, P. D. The psychology of deafness; with suggestions as to a more comprehensive plan of treatment. *THE LARYNGOSCOPE*, p. 257, May, 1915.
- 1317 KERRISON, P. D. Treatment of advanced tympanic deafness. Clinical observations bearing on the problems involved. *Jour. A. M. A.*, p. 109, Jan. 9, 1915.
- 1318 KITSON, H. D. Psychological tests for lip-reading ability. *Volta Review*, p. 471, Dec., 1915.
- 1319 LANNOIS and CHAVANNE. Total deafness from explosion of shell. *Bull. de l'Acad. de Med.*, Jan. 14, 1915.
- *1320 LANNOIS and CHAVANNE. Prognosis of deafness from war injuries. (Le pronostic des surdités de guerre-d'après 1,000 cas.) *Bull. de l'Acad. de Med., Paris*, Dec. 21, 1915.
- 1321 LEDERMAN, I. A. Social and economic aspect of deafness. *Ky. Med. Jour.*, Jan. 1, 1915.
- 1322 LERMOYEZ, M. War deafness. *Presse Med.*, Feb. 25, 1915, and *Progres Medicale*, Dec., 1915.
- 1323 MAURICE, A. Treatment of deafness by re-education of the powers of hearing. *Monopaph*, Paris, 1915.
- *1324 MOLL, A. M. A case of hysterical deafness. *Medical Jour. of South Africa*, p. 3, Aug., 1915.

- 1325 MONRO, S. J. Resume of the rhythmic work in the Horace Mann school, Boston. *Volta Rev.*, p. 133, April, 1915.
- 1326 MORGENSTERN, LOUISE I. Teaching lip-reading to the adult hard of hearing in public school evening classes. *Volta Review*, July, 1915, and Aug., 1915.
- 1327 NAGER, F. R. Modern methods of diagnosis and treatment of deafness. *Corresp. Bl. f. Schweizer. Aerzte*, March 6, 1915.
- 1328 NITCHIE, EDWARD B. What a deaf adult should do to acquire the art of lip-reading. *Volta Review*, July, 1915.
- 1329 NITCHIE, E. B. Detective possibilities of lip-reading. *Volta Rev.*, p. 81, March, 1915.
- 1330 PETERS, E. V. Where do deaf get their "mixed language?" *Am. Ann. of Deaf*, p. 228, May, 1915.
- 1331 PETERSON, P. N. Influence of manual training upon the mental development of the deaf. *Am. Ann. of Deaf*, p. 119, March, 1915.
- *1332 RHESE. Typhoid deafness. (Die Typhus-Schwerhörigkeit.) *Med. Klinik*, Nov. 7, 1915.
- 1333 ROBERTS, EMMA. Training the deaf child. *Volta Rev.*, Vol. 17, No. 8, August, 1915.
- 1334 RODGER, T. R. Noise-deafness, a review of recent experimental work and a clinical investigation into the effect of loud noise upon the labyrinth in boiler-makers. *Jour. of Laryngol.*, p. 91, March, 1915.
- 1335 SEARCH, M. Public schools best for deaf. *Volta Rev.*, p. 112, March, 1915.
- 1336 SMITH, J. C. Development of ability to work independently and of the power of application in young deaf children. *Volta Rev.*, p. 49, Feb., 1915.
- 1337 STEIN, C. and POLLACK, R. Influence of vaso-motor disturbances in children on hearing. *Arch. f. Ohrenh.*, Bd. 96, Heft 3-4, p. 216, 1916.
- 1338 STEVENS, J. E. Schools for the deaf in Europe. *The Volta Rev.*, June, 1915.
- 1339 STORY, A. J. Lip-reading. *Volta Rev.*, V. 17, No. 10, Oct, 1915.
- 1340 STRUCK, A. N. The deaf and the dictionary. *Am. Ann. of Deaf*, p. 233, May, 1915.
- 1341 SZASZ, T. Functional disturbances of hearing. (Über funktionelle Hörstörungen). *Wiener klin. Woch.*, No. 30, 1915.
- 1342 TAYLOR, H. Speech teaching in American schools for the deaf. *Volta Rev.*, p. 56, Feb. 1915.
- 1343 TORREY, G. Lip-reading for the slightly deaf. *Jour. Ophth., Otol. and Laryngol.*, p. 70, Jan., 1915, and *Volta Rev.*, p. 57, Feb., 1915.
- 1344 WALKER, S. T. Utilizing the alumni of our state schools for the deaf. *Am. Ann. of Deaf*, p. 199, May, 1915.
- 1345 WALKER, D. H. Chronic deafness; the present status of its treatment. *Jour. A. M. A.*, p. 663, Aug. 21, 1915.
- 1346 WEINSTEIN, J. Deafness following mumps. *Med. Rec.*, p. 353, Feb. 27, 1915.
- 1347 WINTERMUTE, G. P. Report of an unusual case of labyrinthine deafness. *Cal. State Jour. of Med.*, p. 120, March, 1915.

- †1348 WRIGHT, J. D. Hard of hearing classes in London. *THE LARYNGOSCOPE*, p. 714, Oct., 1915.
- †1349 WRIGHT, J. D. Prophylaxis of deafness. *THE LARYNGOSCOPE*, p. 113, Feb., 1915.
- 1350 WRIGHT, JOHN D. Better speech for the deaf; reform needed in their education. *Jour. Ophthal., Otol. and Laryngol.*, p. 744, Sept., 1915.
- 1351 YEARSLEY, M. The preventive treatment of catarrhal deafness. *Practitioner*, p. 804, June, 1915.
- 1352 YEARSLEY, M. What factor determines ear to be first attacked in chronic middle-ear catarrh. *Jour. Laryngol., Rhinol. and Otol.*, London, No. 5, May, 1915.
- 1353 YEARSLEY, M. Classification of the deaf child for educational purposes. *Teacher of Deaf*, p. 8, Feb., 1915.
- 1354 YEARSLEY, M. Hard of hearing classes in London. *Volta Rev.*, p. 41, Feb., 1915.

General.

- 1355 ALEXANDER, G. Auditory organs of cretins. (Das gehörorgan der kretinen). *Arch. f. Ohrenheilk.*, V. 98, No. 2-3, p. 122, Aug., 1915.
- 1356 BROECKAERT, JULES. Supra-auricular intervention in cases of otic paralysis of the external oculomotor. (Intervention par la voil supra-auriculaire en cas de paralysie otique du moteur oculaire externe associée a la neuralgie du trijumeau). *Rev. de Laryngol., d'Otol. et de Rhinol.*, No. 38-39, Sept. 30, 1915.
- 1357 BULSON, G. A. Ear complications in infections from zymotic fevers. *Mich. State Med. Soc. Jour.*, Jan., 1915.
- *1358 CLAY, J. V. F. Syphilitic lesions of the ear. *Jour. Ophth., Otol. and Laryngol.*, p. 99, Feb., 1915.
- 1359 COX, C. N. Some relations of the ear to general medicine. *L. I. Med. Jour.*, p. 117, Mar., 1915.
- 1360 CRANE, C. G. The conservation of the ears. *L. Island Med. Jour.*, p. 131, Mar., 1915.
- 1361 DAVIS, G. E. Rheumatism in relation to ear affections. *Amer. Med.*, p. 431, June, 1915.
- 1362 DAVIS, J. L. Progress in otology. *American Jour. Dis. Child.*, June, 1915.
- 1363 DENKER, A. Military injuries of ear, upper air tract and the border regions. *Arch. f. Ohrenh.*, Bd. 98, Heft 1, p. 1, 1915.
- 1364 DIGHTON, A. Aural rarities. *Liverpool Med. Chir. Jour.*, Oct., 1915.
- 1365 DOWNEY, J. W. Corrective otoplasty. *Ann. Surg.*, Oct., 1915.
- 1366 EDMONDSON, E. E. Conservation of hearing. *Ill. Med. Jour.*, Feb., 1915.
- 1367 FERRERI, G. Wounds of the ears in war. (Traumi di guerra dell' orecchio). *Polichinico*, No. 34, Aug. 22, 1915.
- 1368 FRASER, J. S. Tubercular disease of the ear. *Proc. Roy. Soc. Med.*, No. 8, Otol. Sect., No. 17, 1915.
- 1369 FRIEDLAENDER, WALTER. Gunshot wounds of the ears (Die Schädigungen des Gehörorgans durch Schusswirkung). *Arch. f. Ohrenheilk.*, V. 98, No. 2-3, p. 158, Aug., 1915.

- 1370 FROST, S. K. Examination of ears of institutional children. *Long Island Med. Jour.*, March, 1915.
- 1371 GRADENIGO, G. Use of the encephaloscope in post-operative treatment of ear cavities. (Sull' impiego dell' encefaloscopio nel trattamento postoperatorio degli ascessi otitici). *Arch. ital. di otol., rinol. e laringol.*, V. 26, No. 3, p. 189, June, 1915.
- 1372 GRADENIGO, G. Memory of sound. (Sulla memoria dei suoni). *Arch. ital. di otol., rinol., e laringol.*, V. 26, No. 4, p. 295, Aug., 1915.
- 1373 GRADENIGO, G. and NUZZI, G. Physiology of the ear (Sulla fisiologia dell' organo uditivo). *Arch. ital. di otolog.*, V. 26, No. 5, p. 403, Oct., 1915.
- 1374 GRIVOT, M. The ear and traumatism of war. *Paris Medical*, Nos. 20-21, Sept. 25, 1915.
- §1375 GUARNACCIA, E. and SBORNI, V. Auricular hematidrosis (Ematidrosi auricolare). *Arch. ital. di otol., rinol. e laringol.*, V. 26, No. 3, p. 193, June, 1915; *THE LARYNGOSCOPE*, p. 684, Oct., 1915.
- 1376 HAGEMANN J. A. Some ocular manifestations of aural disturbances and their interpretation. *Med. Rec.*, p. 100, Jan. 16, 1915.
- *1377 HAMM, O. Treatment of concussion of the ear. (Zur Behandlung der Oherschütterungen). *Munch. Med. Woch.*, No. 48, p. 1664, Nov. 30, 1915.
- 1378 HAUGHEY, W. Bacterial vaccine therapy in diseases of the ear. *Ann. Otol., Rhinol. and Laryngol.*, March, 1915.
- 1379 HAYMAN, L. Gunshot wounds of the ear. *Internat. Zentralbl. f. Ohrenheilk.*, Bd. 13, H. 8; Bd. 13, H. 9; Bd. 13, H. 10; Bd. 13, H. 11; p. 127, 1915.
- 1380 HAYS, H. Ear complications in influenza. *Arch. of Diag.*, April, 1915.
- 1381 HETRICK, L. E. Electrical methods in aural inspection. *Jour. Ophthol., Otol. and Laryngol.*, p. 860, Oct., 1915.
- 1382 HEUSER, C. Technic for diathermy for the ear. *Sem. Med.*, Jan. 21, 1915.
- 1383 HOFER, I. Injuries of the ear in war. (Kriegsverletzungen des Ohres). *Wiener klin. Woch.*, Nov. 11, 1915.
- 1384 KING, J. J. A review of the recent advances in otology. *Med. Times*, Jan., 1915.
- 1385 KREBS, G. Wounds of the ear. *Munch. Med. Wchnschr.*, March 9, 1915.
- 1386 LINNELL, E. H. Some ear cases. *Jour. Ophth., Otol. and Laryngol.*, p. 125, Feb, 1915.
- 1387 LOUGHRAN, R. L. The dangers of neglect in aural diseases. *Woman's Med. Jour.*, Aug., 1915.
- 1388 MATSUMOTO, K. Injection of salt solution in tinnitus. *Sei-I-Kwai Med. Jour.*, Sept., 1915.
- 1389 MCCARTY, V. W. Functional testing in ear disease. *Mo. State Med. Assn. Jour.*, No. 5, May, 1915
- 1390 MCCLELLAND, L. O. Effect of la grippe on the human ear. *L. I. Med. Jour.*, p. 263, June, 1915.
- 1391 NEWARK, W. E. Hemorrhage from ear. *Mich. State Med. Soc. Jour.*, Jan., 1915.

- †1392 PERKINS, C. E. The chorda tympani nerve in otology. *THE LARYNGOSCOPE*, p. 341, June, 1915.
- 1393 PUSATERI, S. So-called "papillomata" of the ear and of the upper respiratory tract. (I cosiddetti "papillomi" dell' orecchio e delle prime vie aeree). *Arch. ital. di otolog.*, Vol. 26, No. 5, p. 453, Oct., 1915.
- 1394 QUIX, F. H. Combined radio-therapy of ear disease. *Nederl. Tijdschr. v. Geneesk.*, Jan. 30, 1915.
- 1395 RAIMY, J. J. A few more of the more frequent ear conditions. *Albany Med. Ann.*, p. 56, Feb., 1915.
- 1396 ROBB, J. M. Aural conditions following contagious disease. *Mich. State Med. Soc. Jour.*, Jan., 1915.
- 1397 ROY, D. More rational methods of treating aural diseases. *Ga. Med. Assn. Jour.*, p. 257, Jan., 1915.
- 1398 SEWELL, D. L. The indications for operative interference in cases of aural discharge. *Medical Chronicle*, p. 49, Nov., 1915.
- 1399 KAHN, A. Myringotomy. *West. Med. Times*, V. XXV, p. 278, 1915.
- 1400 TRIBLE, G. B. Functional testing of the ears. *U. S. Nav. Bull.*, V. 9, No. 3, p. 400, 1915.
- 1401 WEIL, E. Aural injuries in war. (Über Kriegsschädigungen der Ohren). *Munch. Med. Woch.*, No. 48, p. 1661, Nov. 30, 1915.
- 1402 WELTY, C. F. Gun shot wound of the ear with report of two cases. *Cal. State Jour. of Med.*, p. 119, March, 1915.
- *1403 WINGRAVE, W. Lumbar puncture in aural and nasal cases; pathology of the fluid. *Jour. Laryngol., Rhinol. and Otol.*, V. 30, No. 7, July, 1915.

VII. MASTOID AND INTRA-CRANIAL COMPLICATIONS.

Mastoid.

- 1404 ALEXANDER, G. J. Indications for the acute mastoid operation. *Hahneman, Month.*, June, 1915.
- 1405 ALLPORT, F. and ROCHESTER, A. S. Hammond's mastoid skin grafting. *Ann. Otol., Rhinol. and Laryngol.*, V. 24, No. 2, p. 335, June, 1915.
- 1406 ALTER, F. W. Case of mastoiditis with complications and discussion of the value of blood cultures in aiding diagnosis. *Ohio State Med. Jour.*, Sept., 1915.
- 1407 AMBERG, EMIL. Roentgen ray assistance in diagnosing mastoiditis. *Detroit Med. Jour.*, Aug., 1915.
- 1408 AMBERG, E. Some remarks on the X-ray diagnosis of mastoiditis. *Trans. Amer. Otol. Soc.*, Vol. 8, Part 3, p. 598, 1915.
- †1409 BONNER, H. and DUTROW, H. V. Report of a case of primary mastoiditis. *THE LARYNGOSCOPE*, p. 244, April, 1915.
- 1410 BRODER, C. B. Primary mastoiditis: a report of two cases. *Med. Rec.*, p. 482, Sept. 18, 1915.
- 1411 BRYAN, J. H. Report of two cases of mastoid abscess, infective thrombosis of the lateral sinus and jugular vein; resection of the jugular; recovery. *Trans. Amer. Otol. Soc.*, Vol. 8, Part 3, p. 424, 1915, and *Ann. Otol., Rhinol. and Laryngol.*, p. 543, Sept., 1915.

- †1412 CARPENTER, E. R. Preliminary report on the use of plastic operations in large mastoid cavities. *THE LARYNGOSCOPE*, p. 1, Jan., 1915.
- 1413 CARPENTER, E. R. Use of fascia in plastic mastoid work. *Texas State Jour. Med.*, Nov., 1915.
- 1414 CHEATLE, A. Specimens of tuberculosis of the temporal bone. *Proc. Roy. Soc. Med., Otol. Section*, 30, Aug., 1915.
- 1415 CLAY, J. V. F. Four interesting atypical cases of mastoiditis; two with intra-cranial complications. *Jour. Ophthalm., Otol. and Laryngol.*, p. 700, Aug., 1915.
- 1416 COLBURN, F. W. Notes on the modern radical mastoid operation. *Jour. Ophthalmol., Otol. and Laryngol.*, p. 1022, Dec., 1915.
- §1417 DABNEY, V. Idiopathic mastoid abscess. *Jour. A. M. A.*, p. 501, Aug. 7, 1915; *THE LARYNGOSCOPE*, p. 191, Mar., 1916.
- †1418 DENCH, E. B. An improved technique in the application of the Thiersch graft in the radical operation for chronic middle-ear suppuration. *THE LARYNGOSCOPE*, p. 755, Nov., 1915.
- 1419 GOLDSMITH, P. Case of bullet entering below right mastoid and emerging at left orbit. *Proc. Royal Soc. Med., Laryngol. Section*, V. 9, No. 2, Dec., 1915.
- 1420 HARDWICKE, L. C. V. Interesting case of mastoiditis. *Practitioner*, p. 888, June, 1915.
- 1421 HAYS, H. Skiagraphs of the mastoid in certain atypical cases. *Ann. Otol., Rhinol. and Laryngol.*, p. 660, Sept., 1915.
- 1422 HETRICK, L. E. Atypical mastoiditis. *Jour., Ophthalm., Otol. and Laryngol.*, June, 1915.
- 1423 HOLMES, G. and SARGENT, P. Injuries of the superior longitudinal sinus. *Brit. Med. Jour.*, p. 493, Oct. 2, 1915.
- 1424 HUNTINGTON, W. H. Report of case of latent mastoiditis with sinus thrombosis. *Va. Med. Semi-Monthly*, p. 533, Feb. 12, 1915.
- 1425 IMPERATORI, C. J. Report of a case of traumatic mastoiditis. *Ann. Otol., Rhinol. and Laryngol.*, p. 660, Sept., 1915.
- 1426 JONES, W. D. Report of a case of double mastoiditis, with pure culture of streptococcus capsulatus mucosus from the mastoid cells of both ears. *Ann. Otol., Rhinol. and Laryngol.*, March, 1915.
- 1427 JONES, W. D. and DECHERD, H. B. Foreign body in mastoid cells; foreign body in left bronchus. *Texas State Jour. of Med.*, Aug., 1915.
- 1428 JONES, E. L. Relation of rhino-pharynx to the middle-ear and mastoid. *Jour. A. M. A.*, p. 115, Jan. 9, 1915.
- †1429 KIEFFER, G. F. Mastoiditis; a probable cause of acute nephritis. *THE LARYNGOSCOPE*, p. 287, May, 1915.
- 1430 KYLE, J. J. Some observations upon the modern mastoid operation. *Ann. Otol., Rhinol. and Laryngol.*, March, 1915.
- 1431 KYLE, J. J. The early diagnosis of mastoiditis. *Jour. A. M. A.*, p. 496, Aug. 7, 1915.
- 1432 LOEB, H. W. Acute hemorrhagic mastoiditis. *Trans. Amer. Otol. Soc.*, p. 576, Vol. 8, Part 3, 1915, and *Ann. Otol., Rhinol. and Laryngol.*, p. 636, Sept., 1915.

- 1433 MACKENZIE, G. W. Treatment of acute mastoid abscess from the etiologic standpoint. *Jour. Ophth. Otol. and Laryngol.*, p. 154, Feb., 1915.
- 1434 MACNAB, J. C. G. Two cases of the "Gradenigo" syndrome in children eight and ten years respectively. *Jour. Laryngol., Rhinol. and Otol.*, p. 335, Vol. 30, No. 9, Sept., 1915.
- 1435 MAHU, G. Radical cure of chronic mastoiditis. (Ce que devient les évidés.) *Ann. des mal. de l'oreille*, p. 754, Vol. 40, No. 8, 1914-1915.
- 1436 McLOONE, J. J., and NELSON, R. M. Mastoid operation without artery forceps or ligatures. *Ga. Med. Assn. Jour.*, Jan., 1915.
- 1437 MOLLISON, W. M. Acute mastoiditis. *Practitioner*, Oct., 1915.
- 1438 PALEN, G. J. An anatomical consideration of mastoiditis. *N. E. Med. Gaz.*, April, 1915, and *Jour. Ophthal. and Oto-Laryngol.*, May, 1915.
- 1439 PALEN, G. J. Bezold's mastoiditis, a third series of cases. *Jour. Ophth., Otol. and Laryngol.*, p. 35, Jan., 1915.
- 1440 PALEN, G. J. Recent series of mastoid cases. *Jour. Ophth., Otol. and Laryngol.*, p. 144, Feb., 1915.
- 1441 PERKINS, C. E. Facial contractions during the radical mastoid operation. *Ann. Otol., Rhinol. and Laryngol.*, p. 664, Sept., 1915.
- 1442 PIKE, J. B. Permeating mastoiditis. *Practitioner*, Oct., 1915.
- 1443 PLISSON, L. Primary mastoiditis. (Contribution à l'étude des mastoïdites primitives des ostéites primitives infectieuses localisées à la portion mastoïdienne du temporal et en particulier de la forme ostéo-myélique de cette lésion.) *Rev. de Laryngol., d'Otol., et de Rhinol.*, p. 321, Nos. 48-49, Dec., 1915.
- 1444 SECO, D. I. FERNANDEZ. Acute primary mastoiditis. *Rev. de Med. y. Cirur. practicas*, Oct. 14, 1915.
- 1445 SHAPLEIGH, J. B. Insidious mastoiditis. *Mo. State Med. Assn. Jour.*, No. 5, May, 1915.
- 1446 SHEPARD, G. A. Interesting case of mastoiditis. *Jour. Ophthal., Otol. and Laryngol.*, June, 1915.
- 1447 SHEPARD, G. A. An interesting case of mastoiditis. *Jour. Ophthal., Otol. and Laryngol.*, June, 1915.
- †1448 SMITH, S. M. The end-results of the radical mastoid operation. *THE LARYNGOSCOPE*, p. 332, June 1915.
- 1449 SMITH, C. H. Double mastoiditis with septic serous thrombosis. *N. Y. Med. Jour.*, Feb. 27, 1915.
- 1450 SMITH, C. H. Mastoiditis. *N. Y. Med. Jour.*, June, 19, 1915.
- 1451 SOURDILLE, M. Mastoid operation. (Trepanation mastoïdienne élargie et atticotomie bransmastoïdienne [évidemment partiel].) *Rev. de Laryngol., d'Otol. et de Rhinol.*, p. 448, Nos. 48-49, Dec., 1915.
- 1452 STUCKY, J. A. A modified radical mastoid operation. *Trans. Amer. Otol. Soc.*, p. 553, Vol. 8, Part 3, 1915.
- 1453 TODD, F. C. Mastoid abscesses; some indications for operation. *Journal-Lancet*, No. 9, May 1, 1915.
- 1454 WELTON, C. B. Mastoiditis and mastoid abscess without suppuration from the middle-ear and without any apparent middle-ear in 1915.

- 1455 WELTY, C. F. Indications for mastoid operations as shown in one hundred cases. *Jour. A. M. A.*, p. 504, Aug. 7, 1915.
- 1456 WOLFE, C. T. The after-treatment of the mastoid wound. *Louisville Med. Jour.*, June, 1915, and *Jour. Ophthal. and Oto-Laryngol.*, p. 283, Vol. 9, Sept., 1915.

Lateral Sinus.

- 1457 BECCO, R. Suppuration in lateral sinus cured by ligation and curretting. *Sem. Med.*, p. 35, Jan. 14, 1914.
- 1458 BOOT, G. W. Cases of lateral sinus thrombosis. *Ann. of Otol., Rhinol. and Laryngol.*, p. 554, Sept., 1915.
- 1459 CARY, E. H. Spontaneous rupture of lateral sinus with general septicemia in ulcerating sinusitis. *S. Med. Jour.*, Feb., 1915.
- 1460 DANZIGER, E. Thrombosis of the sigmoid sinus. *Internat. Jour. Surgery*, p. 87, March, 1915.
- †1461 DAY, E. W. Report of spontaneous cure of six cases of unrecognized sinus thrombosis accidentally discovered during operation. *THE LARYNGOSCOPE*, p. 757, Nov., 1915.
- 1462 DENCH, E. B. Sinus thrombosis complicating middle-ear suppuration with unusual symptoms. *New Albany Med. Herald*, Nov., 1915.
- 1463 FREUDENTHAL, H., and DOWNEY, J. W. Decompression for the relief of choked disc following thrombosis of the lateral sinus. *Trans. Amer. Otol. Soc.*, p. 503, Vol. 8, Part 3, 1915, and *Ann. Otol., Rhinol. and Laryngol.*, p. 613, Sept., 1915.
- 1464 GRAHAM, H. B. Difficulties in the diagnosis of sinus thrombosis. *Jour. A. M. A.*, p. 848, Sept. 4, 1915.
- 1465 HAAS, E. A syndrome of syphilitic origin simulating thrombophlebitis of the sinuses. *Am. Jour. Ophthal.*, p. 48, Feb., 1915.
- 1466 HALL, G. C. Septic Sinus Thrombosis. *Lancet-Clin.*, June 19, 1915.
- 1467 HEIMAN, H., and BALLIN, M. J. Unusual case of sinus thrombosis. *Jour. A. M. A.*, p. 903, March 20, 1915.
- 1468 JONES, H. E. Some considerations which determine the extent of an operation in septic invasion of the lateral sinus. *Jour. of Laryngol.*, p. 105, March, 1915.
- †1469 KOPETZKY, S. J. A brief consideration of some factors concerned in cases of atypical sinus thrombosis. Preliminary report. *THE LARYNGOSCOPE*, p. 165, March, 1915.
- *1470 LANNOIS and PATEL. Obliteration of lateral sinus to arrest venous static measure in wounds involving large vessels at the base of the brain. *Bull. de l'Acad. de Med. de Paris*, April 27, 1915.
- 1471 LANNOIS and PATEL. Obliteration of lateral sinus to arrest venous hemorrhage in wounds of the neck. *Lyon Chirurgial*, Oct., 1915.
- 1472 PRINGSTEN, C. F. Otitic sinus thrombosis. *Jour. Mo. State Med. Assn.*, June, 1915, and *Jour. Ophthal. and Oto-Laryngol.*, p. 213, Vol. 9, No. 7, July, 1915.
- 1473 SATYARANJAU, S. Hemorrhage from lateral sinus. *Indian Med. Gaz.*, Sept., 1915.

- 1474 SCRUTON, WM. A. Accidental injuries of the sigmoid sinus inflicted in simple mastoidectomy. *Ann. Otol., Rhinol. and Laryngol.*, p. 310, Vol. 24, No. 2, June, 1915.
- 1475 SEN, S. Treatment of hemorrhage from lateral sinus. *Indian Med. Gaz.*, Sept., 1915.
- 1476 SIMPSON, J. R. Thrombosis of lateral sinus. Report of four cases. *Pa. Med. Jour.*, March, 1915.
- 1477 SKROWACZEWSKI, P. The treatment of otitic sinus diseases. (Beitrag zur Behandlung der otitischen Sinuserkrankungen.) *Monatschr. f. Ohrenheilk. u. Laryngo-Rhinol.*, p. 514, Bd. 49, Heft 9 and 10, Sept.-Oct., 1915.
- 1478 STREISSLER. Surgery of the cavernous sinus. (Ein Beitrag zur Chirurgie des Sinus cavernosus.) *Arch. f. klin. Chir.*, Bd. 106, H. 1, 1915.
- 1479 ZAUN, G. F. Unusual case of lateral sinus thrombosis. *Wisc. Med. Jour.*, Dec., 1915.

Intra-cranial Complications and Brain Abscess.

- 1480 ALIN, E. Tumor of the brain simulating eclampsia. (Fall von Gehirntumor.) *Nord. Mediz. Arch.*, Stockholm, Vol. 47, No. 3, 1915.
- 1481 ANDREWS, A. H. Brain and sinus complications of otitis media, with special reference to symptomatology. *Jour. Ophthal. and Oto-Laryngol.*, p. 383, Dec., 1915, and *Texas State Jour. Med.*, Oct., 1915.
- 1482 ANDREWS, B. F. Some reflex manifestations of intranasal origin; suggested nerve paths through which they may travel and operations for their relief. *Jour. Ophthalmol. and Oto-Laryngol.*, p. 354, Nov., 1915.
- 1483 ARCHAMBAULT, L. Diagnosis of central tumor. *Albany Med. Ann.*, May, 1915.
- 1484 AUVRAY, DR. Cranial surgery in military hospitals. *Presse Med.*, Jan. 21, 1915.
- 1485 AYER, W. D. Pathology of brain tumors. *Albany Med. Ann.*, May, 1915.
- 1486 BAKER, D. Treatment of brain tumor. *Albany Med. Ann.*, May, 1915.
- 1487 BALL, C. R. Focal brain disease. *Jour. A. M. A.*, Aug. 14, 1915.
- 1488 BARANY, R. Open and closed treatment of gunshot wounds of the brain (Die offene und geschlossene Behandlung der Schussverletzungen des Gehirns). *Beitr. z. klin. Chir.*, Bd. 97, 1915.
- 1489 BARANY, R. Drainage of brain abscess with gutta percha with some observations on the operative treatment of brain and ear gunshot wounds. *Muench. med. Woch., feldärztliche Beilage* No. 4, 1915.
- 1490 BATTEN, F. B. and VON WYSS, W. H. The atonic form of cerebral diplegia. *Brit. Jour. Child. Dis.*, p. 65, March, 1915.
- 1491 BAUNES, DR. Brain abscess following tooth extraction. *Med. Klin.*, Bd. 11, 392, 1915.

- 1492 BECK, O. Meningitic suppurative extradural abscess of the posterior fossa of the skull after purulent tonsillitis. *Ann. Otol., Rhinol. and Laryngol.*, March, 1915.
- 1493 BECK, J. C. Diagnosis of intracranial complications in diseases of middle ear and accessory sinuses of nose. *Ill. Med. Jour.*, Jan., 1915; *Jour.-Lancet*, March 1, 1915.
- 1494 BIELLER, G. After-treatment of brain abscess with special reference to cerebellar abscess. *Arch. f. Ohrenheilk.*, Vol. 98, p. 58, 1915.
- 1495 BOERNER, DR. Prevention of prolapse of brain after gunshot wounds of the skull (Ein operatives Verfahren zur Verhütung des Hirnprolapses nach Schädelsschüssen). *Muench. med. Woch.*, p. 599, No. 17, April 27, 1915.
- 1496 BRAMWELL, BYROM. Bilateral lesion in the occipital lobes with histo-pathological report showing the position of the cortical area for macular as distinct from non-macular or panoramic vision. *Edinburgh Med. Jour.*, p. 4, July, 1915.
- 1497 BRAMWELL, BYROM. Lesions of the occipital lobe and affection of vision. *Edinburgh Med. Jour.*, p. 165, Sept., 1915.
- †1498 BROSE, L. D. Nose, throat, ear orbital and intra-cranial complications in accessory sinus disease. *THE LARYNGOSCOPE*, p. 35, Jan., 1915.
- 1499 BUELLER, G. After-treatment of brain abscess with especial reference to cerebellar abscess. *Arch. f. Ohrenh.*, Bd. 98, Heft 1, p. 58, 1915.
- 1500 BURKE, G. and NUZUM, J. Differential diagnosis between cerebral hemorrhage and softening due to thrombosis. *Med. Record*, p. 701, Oct. 23, 1915.
- 1501 CASTEX, M. R. and BOLO, P. O. Sarcoma of the left motor region. *Arch. f. Psychiat.*, Vol. 55, p. 479, 1915.
- 1502 COLE, H. P. Case of decompression under local anesthesia. *Ann. Surgery*, Dec., 1915.
- 1503 COUNCILMAN, W. T. The gliomatous tumors of the brain. *Amer. Med. Jour.*, Vol. 53, p. 71, March, 1915.
- 1504 COUNCILMAN, W. T. Anatomic consideration of tumors of brain with special reference to gliomata. *Colo. Med.*, Oct., 1915.
- 1505 DE MONTET and DE LA HARPE. Tumor of the bulbar olive. *Rev. med. de la Suisse Romande*, June 20, 1915.
- 1506 DUKEN, J. Two cases of intracranial pneumatocele (Über zwei Fälle von intrakranieller Pneumatocele nach Schussverletzung). *Muench. med. Woch.*, p. 598, No. 17, April 27, 1915.
- 1507 DUVAL, P. Three cases in which metal plates were used to repair skull defects. *Bull. et mem. Soc. de chir. de Paris*, p. 1228, Vol. 41, 1915.
- 1508 EVANS, C. A. Unlocalized intracranial injuries. *American Jour. Surg.*, p. 441, Vol. 29, No. 12, Dec., 1915.
- 1509 EWING, S. A. Difficulties in diagnosis of intracranial extension of suppurative otitis. *Med. Jour. Australia*, March 27, 1915.
- 1510 FACIO, L. J. and BORGHI, J. A. Abscess of the brain with meningitic syndrome. *Semana Medica*, Sept. 9, 1915.
- 1511 FEARNSIDES, E. G. Case of dyspraxia and paraphasia; necropsy; tumor in the supramarginal convolution. *Brain*, March, 1915.

- §1512 FRASER, J. S. Two cases of otitic extradural abscess. *Edin. Med. Jour.*, p. 38, Jan., 1915. *THE LARYNGOSCOPE*, p. 127, Feb., 1916.
- 1513 FRASER, J. S. and LOGAN, W. R. Two cases with multiple intracranial complication of chronic otitis media: recovery after operation. *Edinburgh Med. Jour.*, p. 350, Nov., 1915.
- 1514 FRIEDMAN, E. D. Idiopathic hypertrophy of the left ventricle; its relationship to status-thymo-lymphaticus. *Interstate Med. Jour.*, p. 1128, Nov., 1915.
- 1515 FUMAROLA, G. Clinical syndrome of tumors of the cerebello-pontine angle. *Arch. f. Psychiat.*, p. 781, Vol. 56, 1915.
- 1516 GIERLICH, DR. Papilloma in fourth ventricle with attempted operation and necropsy findings. *Deut. mediz. Woch.*, Vol. 41, No. 30, July 22, 1915.
- 1517 GLEASON, E. B. and PFAHLER, G. E. A case of fracture confined to the petrous position of the temporal bone. *Amer. Jour. Roentgen.*, p. 604, Jan., 1915.
- 1518 GOERDT, WILHELM. Abscess of temporal lobe with involvement of lateral ventricle (Ein Fall von Heilung eines Schläfenlappenabscesses mit Durchbruch in den Seitenventrikel). *Arch. f. Ohrenheilk.*, Vol. 98, No. 2-3, p. 101, Aug., 1915.
- 1519 GOETZ, J. Infections within the head. *St. Paul. Med Jour.*, p. 105, Vol. 17, 1915.
- 1520 GOLDSTEIN, K. Cerebellar symptom-complex in cranial injuries (Über den zerebellaren Symptomkomplex in seiner Bedeutung für die Beurteilung von Schädelverletzungen). *Muench. med. Woch.*, p. 1439, Oct. 19, 1915.
- *1521 GREY, E. G. Studies on the localization of cerebellar tumors. III. Posterior new growths without nystagmus. *Jour. A. M. A.*, p. 1341, Oct. 16, 1915.
- 1522 GREY, E. G. Studies on localization of cerebellar tumors. *Jour. Nervous and Mental Dis.*, Oct., 1915.
- 1523 GRUENEWALD, L. Gunshot wounds of the cranial cavity (Schussverletzungen der pneumatischen Schädelhöhlen). *Muench. med. Woch.*, No. 24, p. 823, June, 1915.
- *1524 GUTHRIE, THOMAS. Bilateral lesion of the auditory centre. *Jour. Laryngol., Rhinol. and Otol.*, Vol. 30, No. 5, May, 1915.
- *1525 GUTHRIE, L. Tuberculoma of pons varolii. *Brit. Jour. Child. Dis.*, Vol. 12, No. 140, Aug., 1915.
- †1526 GUTTMAN, J. Report of a case of multiple abscess of the brain; operation; recovery. *THE LARYNGOSCOPE*, p. 284, May, 1915.
- 1527 HARMS, H. Cured case of multiple brain abscess formation after acute middle ear suppuration. *Ztschr. f. Ohrenh.*, Bd. 72, Heft 2, p. 118, 1915.
- *1528 HASELTINE, B. A rare case of cerebellar abscess. *Jour. Ophthol., Otol. and Laryngol.*, p. 650, Aug., 1915.
- 1529 HAWTHORNE, C. O. Localizing brain symptoms (hemiplegia and hemispasm) as early events in tuberculous meningitis. *Brit. Jour. Child. Dis.*, Aug., 1915.
- 1530 HAYWARD, E. Gunshot wounds of the skull. *Berliner Klin. Woch.*, No. 45, No. 8, 1915.

- 1531 HOUSE, W. The relation of special sense function to intracranial disease. *Ophthalmology*, p. 279, Jan., 1915.
- 1532 HOWLAND, G. W. Diagnosis of subtentorial tumors. *Can. Pract. and Rev.*, p. 51, 1915.
- 1533 JEGER, E. Plastic closure of the dura with fascia in gunshot wounds of the brain (Über primäre Fascienplastik bei Schussverletzungen der Dura). *Beitr. z. klin. Chir.*, Bd. 97, 1915.
- †1534 JOHNSON, G. B. The ocular symptoms of brain abscess and sinus thrombosis of otitic origin; report of a case. *THE LARYNGOSCOPE*, p. 7, Jan., 1915.
- 1535 KENYON, J. H. Cerebral surgery. *Ann. Surg.*, Jan., 1915.
- 1536 KERN, L. C. Case of brain abscess. *Iowa State Med. Soc. Jour.*, Jan., 1915.
- 1537 KRAMER, S. P. Cerebellar surgery. *Ohio State Med. Jour.*, May, 1915.
- *1538 KROHN, G. H. M. Diagnosis of tumor in cerebello-pontine angle. *Norsk. Mag. for Lægevid.*, Vol. 76, No. 6, June, 1915.
- 1539 LASSEN, F. Brain abscess of otitic origin. *Colo. Med.*, Jan., 1915.
- †1540 LESHURE, J. A case of temporosphenoidal abscess with unusual complications. *THE LARYNGOSCOPE*, p. 281, May, 1915.
- 1541 LOEFBERG, O. Covering defects in the skull (Zur Deckung von Kranialdefekten). *Militärarzt*, Vol. 46, p. 274, 1915.
- 1542 LOUGHRAN, R. L. A case of acute exacerbation of a chronic mastoiditis with cholesteatoma, extensive epidural abscess, purulent pachymeningitis, with exploratory incision of the dura and lateral sinus; recovery. *Ann. Otol., Rhinol. and Laryngol.*, p. 676, Sept., 1915.
- 1543 MACRIS, E. Albuminuria not constant with meningeal hemorrhage. *Grece med.*, Athens, Vol. 17, Nos. 9-10, May 1-15, 1915.
- 1544 MANASSE, P. Therapy of brain abscess. *Muench. med. Woch.*, p. 1475, Oct. 26, 1915.
- 1545 MAYER, OTTO. Plastic covering of dural defects (Über die plastische Deckung von Duradefekten nach Abtragung von Hirnprolapsen in der Otochirurgie). *Ztschr. f. Ohrenheilk.*, Bd. 73, H. 1, p. 37, July, 1915.
- 1546 MILLIGAN, W. Some practical considerations in the diagnosis and treatment of abscess of the cerebellum with a record of cases subjected to operation. *Jour. of Laryngol.*, p. 4, Jan., 1915.
- *1547 MOORE, J. W. Fracture of base of skull with escape of cerebrospinal fluid from ear. Effect of atropin and epinephrin upon the secretion. *Am. Jour. Med. Sci.*, p. 580, April, 1915.
- †1548 MURPHY, J. W. Brain infection of otic origin with report of five cases. *THE LARYNGOSCOPE*, p. 701, Oct., 1915.
- 1549 O'SHEA, H. V. Traumatic subdural hemorrhage. *Practitioner*, Oct., 1915.
- *1550 PAGE, J. R. Case of temporosphenoidal abscess discovered by exploration through multiple small incisions in the dura; drained and cured. Remarks on technique of operation. *Surg., Gyn. and Obst.*, p. 718, June, 1915.

- 1551 PERTHES, G. Removal of bone splinters and foreign bodies in cranial injuries and brain abscess (Schonende Entfernung von Knochensplittern und Fremdkörpern bei Schädelchüssen und Heirnsabszessen). *Muench. med. Woch.*, p. 1706, No. 49, Dec. 7, 1915.
- 1552 PORTER, L. Apparently complete recovery after operation for cerebellar tumor. *Orch. Pediatrica*, Oct., 1915.
- 1553 REIK, H. O. Case of cerebral abscess secondary to chronic suppurative otitis media, cured by surgical intervention. *South. Med. Jour.*, Vol. 8, No. 8, Aug., 1915.
- 1554 RICHARDS, G. L. Report of a case of brain abscess with unusual findings. *Trans. Amer. Otol. Soc.*, 1915, *Ann. of Otol., Rhinol. and Laryngol.*, p. 561, Sept., 1915.
- *1555 SACHS, E. Intracranial telangiectasis: symptomatology and treatment with report of two cases. *Amer. Jour. Med. Sc.*, p. 565, Oct., 1915.
- 1556 SANZ, E. F. Injuries to the medulla in war (Los traumatismos de la medula segun la experiencia de la actual guerra europea). *Los Prog. de la Clin.*, p. 344, No. 30, June, 1915.
- 1557 SANZ, E. F. Diagnosis of cerebral tumors (Diagnostico de los tumores cerebrales). *Revista Ibero-Americana de Ciencias Medicas*, Aug., 1915, and *Los Progresos de la Clinica*, p. 108, No. 33, Sept., 1915.
- 1558 SATTLER, ROBERT. Double papillo-edema. Optic neuritis. The clinical constant of brain and cerebellar tumors. *Jour. of Ophthal. and Oto-Laryngol.*, p. 243, Vol. 9, No. 8, Aug., 1915.
- 1559 SAUER, W. E. Otitic brain abscess. *Mo. State Med. Ass'n Jour.*, No. 5, May, 1915.
- 1560 SCHMINCKE, A. Glioblastic sarcoma of the cerebellum with metastasis (Ein glioblastisches Sarkom des Kleinhirns mit Metastasenbildung in Hirn und Rückenmark). *Frankfurt. Zeitschr. f. Pathol.*, Bd. 16, H. 3, 1915.
- 1561 SHEMELY, WM. G. Successful removal of a bullet lodged in the region of the carotid artery in the skull. *Jour. Ophthalmol., Otol. and Laryngol.*, p. 1048, Dec., 1915.
- 1562 SHERMAN, THEODORE. Miliary aneurysms in relation to cerebral hemorrhage. *Edinburgh Med. Jour.*, p. 245, Oct., 1915.
- 1563 SHUTER, R. E. Notes on a case of tumor of the auditory nerve. *Med. Jour. Australia*, Feb. 27, 1915.
- 1564 SHUTER, R. E. Intracranial extension of middle ear disease. *Med. Jour. Australia*, March 27, 1915.
- 1565 SOUDER, C. R. Cerebral abscess with recovery. Case report. *Ind. Med. Jour.* p. 148, April, 1915.
- 1566 SPENCER, M. Osteoplastic resection of the skull in decompression operations on the brain. *Lancet*, Jan. 9, 1915.
- 1567 TETER, C. K. Anesthesia for brain surgery. *Amer. Jour. Surg., Anesthesia Supp.*, p. 144, Oct., 1915.
- 1568 THROCKMORTON, T. B. Cerebral abscess, probably primarily due to suppurative tonsillitis. *Iowa State Med. Soc. Jour.*, Oct., 1915.

- 1569 TODDE, C. Echinococcus cyst of the brain. *Lancet*, Sept. 25, 1915.
- 1570 TROTTER, W. The principles of the operation of traumatic cerebral lesions. *Brit. Jour. Surg.*, p. 520, April, 1915.
- *1571 WEISENBURG, T. H. and WOEK, P. Diagnosis of tumors in the posterior cranial fossa. *Jour. A. M. A.*, p. 1345, Oct. 16, 1915.
- 1572 WIESE, B. Methods of covering cranial defects (Methoden zur Deckung von knöchernen Schädeldefekten). *Inaug. Diss. Strassburg*, July-Oct., 1915.
- 1573 WOODBURY, M. S. Cerebellar symptomatology. *Med. Rec.*, p. 369, Sept. 4, 1915.
- 1574 ZDRODOVSKY, P. P. Plastic surgery of defects of the dura. *Russky Vrach*, Vol. 14, No. 6, Feb. 7, 1915.

Meningitis.

- §1575 ABECHUCO, J. L. Surgical treatment of acute serous meningitis. *Rev. de med. y cir. prac.*, June 14, 1915. *THE LARYNGOSCOPE*, p. 583, Aug., 1915.
- 1576 ASHBY, H. T. Eight cases of epidemic cerebro-spinal meningitis in infants. *Brit. Med. Jour.*, May, 1915.
- 1577 BATTEN, F. E. Epidemic cerebro-spinal meningitis. *Lancet*, May 8, 1915.
- 1578 BERENS, T. P. Ambulant otitic meningitis. *Am. Jour. of Surg.*, p. 147, April, 1915.
- 1579 BOLAND, C. V. Case of influenzal meningitis. *Lancet*, Sept. 25, 1915.
- 1580 BROWN, A. Influenzal meningitis; report of two cases. *Canad. Med. Ass'n Jour.*, Dec., 1915.
- 1581 BRYANT, W. S. Treatment of purulent streptococcic cerebro-spinal meningitis. *Surg., Gyn. and Obst.*, p. 240, Vol. 20, 1915.
- 1582 CALDERA, C. and FINZI, G. The treatment of otitic meningitis. *Arch. ital. di Otol., Rinol. e Laryngol.*, Vol. 26, April, 1915.
- 1583 CHIARI, H. Pathogenesis of meningitis after gunshot wounds of the brain (Zur Pathogenese der Meningitis bei Schussverletzungen des Gehirns). *Muench. med. Woch.*, p. 596, No. 17, April 27, 1915.
- 1584 CHIRAY. Pus in the ventricles of the brain in meningitis. *Presse medicale*, Dec. 2, 1915.
- 1585 ELY, F. A. Influenzal meningitis; report of a case. *Iowa State Med. Soc. Jour.*, Oct., 1915.
- 1586 FEARNSIDE, E. G. and GUTHRIE, L. Syphilis meningo-vascularis; congenital syphilis; choroiditis; optic atrophy; herpes zoster; multiple root lesions. *Brit. Jour. Child. Dis.*, p. 206, July, 1915.
- 1587 HOSKINS, W. D. Modern methods in diagnosis and treatment of cerebro-spinal meningitis. *Jour. Ind. State Med. Soc.*, March, 1915.
- 1588 JACQUES, P. Otitic meningitis (Contribution à l'étude de la meningite otogène). *Rev. de Laryngol., d'Otol. et de Rhinol.*, p. 233, Nos. 40-41, Oct. 15, 1915.

- 1589 JEMMA, R. Meningococcus meningitis in infants and serotherapy. *Pediatrics*, Nov., 1915.
- 1590 KOPETZKY, S. J. A case of otitis media and mastoiditis followed by meningitis—operation—recovery. After eight years, development of cerebral abscess—operation—death. *Ann. of Otol., Rhinol. and Laryngol.*, p. 566, Sept., 1915.
- 1591 KURAK, MARIE. Treatment of cerebrospinal meningitis with large doses of serum. *Med. Klinik*, Sept. 19, 1915.
- 1592 KURIBAYASHI, K. Present conditions of treatment of suppurative meningitis. *Sei-I-Kwai Med. Jour., Tokyo*, Oct., 1915.
- 1593 LACOSTE and BODIN. A typical acute meningitis. *Paris Med.*, Vol. 5, No. 9, 1915.
- 1594 LANNOIS and GIGNOUX. Fulminating otitic meningitis (La meningite otogène foudroyante.) *Ann. des mal de l'oreille*, Vol. 40, No. 7, 1914-1915.
- 1595 LOUGHRAN, R. L. A case of acute meningitis associated with acute suppurative otitis media, both produced by streptococcus mucosus capsulatus. *Ann. Otol., Rhinol. and Laryngol.*, p. 673, Sept., 1915.
- 1596 LUNDIE, A., THOMAS, D. J., FLEMING, S. Cerebro-spinal meningitis; diagnosis and prophylaxis. *British Med. Jour.*, March, 20, 1915.
- 1597 MYGIND, H. Otogenous meningitis. Statistics, prognosis, treatment. *Ztschr. f. Ohrenh.*, Bd. 72, Heft 2, p. 73, 1915.
- 1598 NEAL, JOSEPHINE, B. Summary of four years of clinical and bacteriologic experience with meningitis in New York City. *Amer. Jour. Dis. Child.*, Jan., 1915.
- §1599 PETERS, E. A. Note on the nasal conditions of cerebro-spinal meningitis. *Jour. Laryngol., Rhinol. and Otol.*, Vol. 30, No. 7, July, 1915. *THE LARYNGOSCOPE*, p. 187, March, 1916.
- 1600 ROSENBAUM, N. Fulminating meningitis with atypical symptoms. *Medizin. Klinik*, Dec. 26, 1915.
- 1601 RUTELLI, G. Streptothrix cerebro-spinal meningitis in a six-months infant. *Pediatrics*, Oct., 1915.
- 1602 SVETKA, V. Meningococcus sepsis. *Wiener klin. Woch.*, Dec. 2, 1915.
- 1603 WEGELIN, C. Actinomycotic purulent meningitis. *Correspond. Bl. für Schweizer Aerzte*, May 1, 1915.

VIII. NEW INSTRUMENTS.

Ear.

- 1604 FENTON, R. A. Simple adaptation of pocket flash-light for ear and eye examinations. *Jour. A. M. A.*, p. 240, Jan. 16, 1915.
- 1605 GATSCHER, S. Apparatus for demonstrating and proving the pointing test. *Monatschr. f. Ohrenh.*, Heft 8, p. 1112, 1915.
- 1606 KINDLMANN, J. An apparatus for distinguishing between true and simulated deafness. (Apparat um Taubheit-Simulierende zu entlarven.) *Wiener klin. Woch.*, Nov. 11, 1915.

- 1607 KOFLER, K. Auricle elevator and dissector. *Monatschr. f. Ohrenheilk. u. Laryngo. Rhinol.*, p. 602, Bd. 49, H. 9 and 10, Sept.-Oct., 1915.
- 1608 MALLOCK, A. Mallock-Armstrong ear defender. *Brit. Med. Jour.*, p. 25, Jan. 2, 1915.
- 1609 MEYERDING, HENRY, W. A simple operating mallet. *Jour. A. M. A.*, p. 1181, Oct. 2, 1915.
- 1610 SPENCER, F. R. A labyrinth probe. *Ann. Otol., Rhinol. and Laryngol.*, p. 680, Sept., 1915.
- 1611 STOCKER, A. An apparatus for testing the hearing (Neuer Gehörmesser). *Muench. mediz. Woch.*, Nov. 6, 1915.

Nose and Naso-pharynx.

- †1612 CARMODY, T. E. A new suction apparatus for nose, throat and ear operators. *THE LARYNGOSCOPE*, p. 777, Nov., 1915.
- †1613 COCKS, G. H. Improved Glatzell mirror. *THE LARYNGOSCOPE*, p. 135, March, 1915.
- 1614 CULLOM, M. MCT. New instrument for the removal of adenoids. *Jour. A. M. A.*, p. 1067, March 27, 1915.
- 1615 HAYS, H. Suction tubes for use in nose, throat and ears. *Jour. A. M. A.*, p. 421, July 31, 1915.
- 1616 HUBBY, L. M. A simple manometric apparatus for measuring Eustachian tubal patency and tympanic ossicular motility. *Ann. Otol., Rhinol. and Laryngol.*, p. 307, Vol. 24, No. 2, June, 1915.
- 1617 HUTCHINSON, A. J. A short note on a simple apparatus for applying ionic medication to the nasal cavities. *Jour. Laryngol., Rhinol. and Otol.*, p. 385, Vol. 30, No. 10, Oct., 1915.
- 1618 KOFLER, K. Radium carrier for the anterior part of nasal septum. *Monatschr. f. Ohrenheilk. u. Laryngo-Rhinol.*, p. 610, Bd. 49, H. 9 and 10, Sept.-Oct., 1915.
- 1619 KOFLER, K. A curette for Rosenmüller's groove. *Monatschr. f. Ohrenheilk. u. Laryngo-Rhinol.*, p. 602, Bd. 49, H. 9 and 10, Sept.-Oct., 1915.
- 1620 KOFLER, K. A nasal speculum for operations on the hindermost parts of the nose. *Monatschr. f. Ohrenheilk. u. Laryngo-Rhinol.*, p. 600, Bd. 49, H. 9 and 10, Sept.-Oct., 1915.
- 1621 KOFLER, K. A vomer forceps. *Monatschr. f. Ohrenheilk. u. Laryngo-Rhinol.*, p. 601, Bd. 49, H. 9 and 10, Sept.-Oct., 1915.
- 1622 KOPFLER, K. A sharp septum forceps. *Monatschr. f. Ohrenheilk. u. Laryngo-Rhinol.*, p. 600, Bd. 49, H. 9 and 10, Sept.-Oct., 1915.
- 1623 LUBMAN, MAX. Protective shield for the Holmes naso-pharyngoscope. *N. Y. Med. Jour.*, March, 1915.
- 1624 MCKENZIE, DAN. Nasal forceps. *Jour. Laryngol., Rhinol. and Otol.*, Vol. 30, No. 11, p. 456, Nov., 1915.
- 1625 ORENDORFF, O. Nasal dressing forceps. *Jour. A. M. A.*, p. 1316, April 17, 1915.
- 1626 PRINCE, A. E. New nasal punch. *Ophthalmic Record*, Dec., 1915.
- 1627 SHEEHAN, JOSEPH E. A self-retaining retractor for the submucous resection of the nasal septum. *Jour. A. M. A.*, p. 1183, Oct. 2, 1915.

- 1628 SHUMWAY, E. A. A self-illuminated Berger loupe. *Jour., A. M. A.*, p. 1363, Oct. 16, 1915.
- 1629 STOCKARD, C. Convenient face shield for nose and throat work. *Ga. Med. Ass'n Jour.*, Jan., 1915.

Mouth and Pharynx.

- 1630 ALTER, F. W. Some new instruments to simplify the modern tonsillectomy operation. (Mouth gag with lip retractor; combined tonsil-seizing forceps and tongue depressor; suction apparatus). *Jour. A. M. A.*, June 5, 1915.
- 1631 BORGMAYER, J. G. L. and LARKEY, C. J. An appliance for producing simultaneous suction and pressure in tonsil and adenoid operations. *Medical Record*, May 1, 1915.
- †1632 BYRNE, WM. D. Tonsil enucleator (and tonsil volsellum). *THE LARYNGOSCOPE*, p. 778, Nov., 1915.
- *1633 GLIDDEN, S. C. A new tonsil-holding forceps. *Surg., Gyn. and Obst.*, p. 662, Nov., 1915.
- †1634 HAGEMANN, J. A. An improved tonsillar hemostat. *THE LARYNGOSCOPE*, p. 588, Aug., 1915.
- *1635 HEACOCK, J. H. A new tonsil depressor. *N. Y. Med. Jour.*, p. 1234, 1234, Dec. 18, 1915.
- 1636 KAHN, A. An electrical mouth light. *Med. Rec.*, p. 609, April 10, 1915.
- 1637 KOFLER, K. Radium carrier for the tonsillar region. *Monatschr. f. Ohrenheilk. u. Laryngo-Rhinol.*, p. 610, Bd. 49, H. 9 and 10, Sept.-Oct., 1915.
- 1638 KOFLER, K. Radium carrier for the naso-pharynx. *Monatschr. f. Ohrenheilk. u. Laryngo-Rhinol.*, p. 609, Bd. 49, H. 9 and 10, Sept.-Oct., 1915.
- 1639 KOFER, K. Sharp naso-pharyngeal forceps. *Monatschr. f. Ohrenheilk. u. Laryngo-Rhinol.*, p. 602, Bd. 49, H. 9 and 10, Sept.-Oct., 1915.
- 1640 LARKEY, C. J. and BORGMAYER, J. G. L. An appliance for producing simultaneous suction and pressure in tonsil and adenoid operations. *Medical Record*, May 1, 1915.
- 1641 LUMBEARD, J. E. A controller of the tongue and palate during general anesthesia. *Jour. A. M. A.*, May 22, 1915.
- 1642 MEGRAIL, W. P. Tonsil forceps. *Jour. A. M. A.*, p. 1365, Oct. 16, 1915.
- 1643 MURPHY, F. G. New instrument and technic for circumcision of the tonsil. *Jour. A. M. A.*, p. 1551, Oct. 30, 1915.
- *1644 PINNEO, F. W. A pharyngeal tube combining rebreathing and insufflation. *Jour. A. M. A.*, p. 1279, Oct. 9, 1915.
- 1645 ROCKEY, A. E. A pharyngeal inhaler for anesthesia in operations of the jaws, head, neck and cervical spine. *Amer. Jour. Surg.*, p. 165, Quart. Supp., Oct., 1915.
- 1646 SANDERSON, W. New guillotine. *Jour. of Laryngol.*, p. 88, Feb., 1915.
- 1647 TITUS, N. E. A suction tongue depressor. *Med. Rec.*, p. 922, Nov. 27, 1915.

Accessory Sinuses.

- *1648 LAW, F. M. Accessory sinus angle finder. *Amer. Jour. Roentgenol.*, Nov., 1915.
- †1649 NEW, G. B. Antrum trocar and irrigating canula. *THE LARYNGOSCOPE*, p. 864, Dec., 1915.
- 1650 PRENN, J. Suction and irrigation apparatus for diagnosis and treatment of antrum of Highmore. *Bost. Med. and Surg. Jour.*, Sept. 16, 1915.
- 1651 SCHUMACHER, H. New instrument for maxillary sinus through inferior nasal meatus. *Monatschr. . Ohrenh.*, Heft 1, p. 47, 1915.
- 1652 VAIL, D. T. New retractor for extirpation of tear sac operation, exterior frontal sinus operation and infant mastoid operation. *Ophthalmic Rec.*, Vol. 24, No. 7, July, 1915.

Larynx. Trachea. Bronchi. Esophagus.

- 1653 DRESSLER, C. E. and KAHN, A. An ideal naso-laryngeal inhaler. *Med. Record*, p. 997, Dec. 11, 1915.
- †1654 GLOGAU, OTTO. A direct laryngoscopic spatulum. *THE LARYNGOSCOPE*, p. 860, Dec., 1915.
- †1655 IGLAUER, S. A new support for the suspension laryngoscope. *THE LARYNGOSCOPE*, p. 672, Sept., 1915.
- 1656 INGEBRIGSTEN, R. Overpressure apparatus for general anesthesia by intratracheal insufflation. *Norsk. Mag. for Laegevidenskaben*, Dec., 1915.
- †1657 JACKSON, CHEVALIER. An anterior commissure laryngoscope. *THE LARYNGOSCOPE*, p. 589, Aug., 1915.
- 1658 KOFLER, K. Radium carrier for the esophagus. *Monatschr. f. Ohrenheilk. u. Laryngo-Rhinol.*, p. 609, Bd. 49, H. 9 and 10, Sept.-Oct., 1915.
- 1659 KOFLER, K. Laryngeal dilators. *Monatschr. f. Ohrenheilk. u. Laryngo-Rhinol.*, p. 604, Bd. 49, H. 9 and 10, Sept.-Oct., 1915.
- 1660 KOFLER, K. An elliptical tracheobronchoscope for children. *Monatschr. f. Ohrenheilk. u. Laryngo-Rhinol.*, p. 605, Bd. 49, H. 9 and 10, Sept.-Oct., 1915.
- 1661 KOFLER, K. A cutting epiglottis forceps. *Monatschr. f. Ohrenheilk. u. Laryngo-Rhinol.*, p. 603, Bd. 49, H. 9 and 10, Sept.-Oct., 1915.
- 1662 KOFLER, K. A cutting laryngeal forceps. *Monatschr. f. Ohrenheilk. u. Laryngo-Rhinol.*, p. 603, Bd. 49, H. 9 and 10, Sept.-Oct., 1915.
- 1663 LYNNAH, H. L. Intubation apparatus and tubes for treatment of chronic stenosis of the larynx. *Jour. A. M. A.*, p. 1066, March 27, 1915.
- †1664 LYNNAH, HENRY L. A new bivalve laryngeal speculum. *THE LARYNGOSCOPE*, p. 673, Sept., 1915.
- 1665 PESCHER, J. A bottle spiroscope for exercising the respiratory apparatus. *Paris Medical*, Dec. 4, 1915.
- 1666 POPE, S. T. Apparatus for pharyngeal and intratracheal anesthesia. *Amer. Jour. Surg.*, p. 164, Quart. Supp., Oct., 1915.
- 1667 ROBINSON, S. Intratracheal ether anesthesia; a new apparatus and intracheal tube. *Surg., Gyn. and Obst.*, p. 774, Dec., 1915.

- 1668 ROEHR, H. Universal apparatus for compression therapy in laryngeal affections. *Arch. f. Laryngeol.*, Bd. 29, Heft 2, p. 179, 1915.
- 1669 ULLMANN, H. J. Apparatus for utilizing city water pressure to furnish compressed air for atomizers. *Jour. A. M. A.*, July 17, 1915.
- 1670 VOORHEES, I. W. A new atomizer bracket. *Jour. A. M. A.*, March 6, 1915.

General.

- 1671 ALBERT, H. A new museum jar for the preservation of eye specimens. *Jour. A. M. A.*, p. 1278, Oct. 9, 1915.
- 1672 BERNHEIM, B. M. A simple instrument for the indirect transfusion of blood. *Jour. A. M. A.*, p. 1278, Oct. 9, 1915.
- 1673 BETTMAN, A. G. A new neck bandage. *Jour. A. M. A.*, p. 1363, Oct. 16, 1915.
- 1674 BOWERS, L. G. and ROSS, L. F. A simple and satisfactory closed ether apparatus. *Surg., Gyn. and Obst.*, p. 521, Vol. 21, No. 4, Oct., 1915.
- 1675 BROWN, O. H. A practical sterilizing pocket thermometer case. *Jour. Mo. State Med. Ass'n*, p. 498, Nov., 1915.
- 1676 BROWN, H. C. Simple and convenient water bath for use in hemolytic experiments. *Indian Jour. Med. Research, Calcutta*, July, 1915.
- 1677 COBURN, R. C. An improved inflatable face cushion. *Jour. A. M. A.*, July 17, 1915.
- 1678 DEBERER, C. A new artery forceps. *Jour. A. M. A.*, p. 1364, Oct. 16, 1915.
- 1679 GIFFORD, H. Improved form of brain explorer. *Jour. A. M. A.*, June 26, 1915.
- 1680 GWATHMEY, J. T. An apparatus for vapor anesthesia and resuscitation. *Amer. Jour. Surg., Anesthesia Supp.*, p. 161, Oct., 1915.
- 1681 HORN BROOK, R. W. Simple and safe somnoform or ethyl chlorid inhaler for minor operations. *Jour. Roy. Nav. Med. Service*, No. 3, July 1, 1915.
- 1682 HULSE, J. A. VAN DER. Irrigating tip. *Jour. A. M. A.*, p. 1551, Oct. 30, 1915.
- 1683 ILIN, I. P. New sterilizing apparatus for dressings and instruments. *Russky Vrach*, Vol. 14, No. 7, Feb. 14, 1915.
- 1684 JACOBSON, J. H. An all-metal syringe for use in local anesthesia. *Amer. Jour. Surg.*, p. 349, Sept., 1915.
- 1685 KOFLER, K. Paraffin syringe. *Monatschr. f. Ohrenheilk. u. Laryngo-Rhinol.*, p. 608, Bd. 49, H. 9 and 10, Sept.-Oct., 1915.
- 1686 KOFLER, K. T-cannulae with removable upper parts. *Monatschr. f. Ohrenheilk. u. Laryngo-Rhinol.*, p. 607, Bd. 49, H. 9 and 10, Sept.-Oct., 1915.
- 1687 KOFLER, K. An electric head mirror. *Monatschr. f. Ohrenheilk. u. Laryngo-Rhinol.*, p. 607, Bd. 49, H. 9 and 10, Sept.-Oct., 1915.
- 1688 KOFLER, K. A canula attachment. *Monatschr. f. Ohrenheilk. u. Laryngo-Rhinol.*, p. 604, Bd. 49, H. 9 and 10, Sept.-Oct., 1915.

- 1689 LAYTON, O. M. Simple apparatus for the administration of chloroform and ether by the vapor method. *Jour. A. M. A.*, p. 824, March 6, 1915.
- 1690 LUKE, H. C. A new gas-oxygen-ether apparatus. *Medical Record*, June 12, 1915.
- 1691 LYON, JR., W. W. An easily filled capsule for collecting blood. *Jour. A. M. A.*, p. 1813, Nov. 20, 1915.
- 1692 MONTGOMERY, F. The etherometer a means of mechanical anesthesia. *Amer. Jour. Obstet.*, p. 133, Vol. 72, 1915.
- 1693 MUENNICH. A simple apparatus for infusion of sterile salt solution. *Muench. med. Woch.*, p. 1552, No. 45, Nov. 9, 1915.
- 1694 POOL, E. H. A device for breaking glass tubes containing suture material. *Med. Rec.*, p. 320, Aug. 21, 1915.
- 1695 SAXON, G. J. and DRUMMOND, L. E. A combination holder and aspirator for use with the blood pipet. *Jour. A. M. A.*, p. 1182, Oct. 2, 1915.
- 1696 SCHMIDT, C. L. A. Electric heater for staining the tubercle bacillus. *Jour. A. M. A.*, p. 823, March 6, 1915.
- 1697 SHATTOCK, S. G. An adaptable eye shade for microscopic work. *Brit. Med. Jour.*, p. 504, Oct. 2, 1915.
- 1698 SHERMAN, H. M. A paper-bag holder for soiled surgical dressings. *Calif. State Jour. Med.*, Nov., 1915.
- 1699 STEPHENS, R. A practical operating mask. *Jour. A. M. A.*, p. 1813, Nov. 20, 1915.
- 1700 SUFFA, G. A. A new ophthalmotrope. *Jour. Ophthal., Otol. and Laryngol.*, March, 1915.
- 1701 THIEMANN, H. Gunshot wounds of the skull (Schädelschüsse). *Muench. med. Woch.*, p. 593, No. 17, April 27, 1915.
- 1702 WEGEFORTH, H. M. and P. Universal surgical motor. *Jour. A. M. A.*, p. 420, July 31, 1915.
- 1703 WEISS, RICHARD. A simple portable apparatus for the Wassermann reaction. *Muench. med. Woch.*, p. 1355, No. 40, Oct. 5, 1915.
- 1704 WHITEFORD, C. H. Apparatus for rectal administration of saline solution. *Brit. Med. Jour.*, p. 538, Oct. 9, 1915.
- 1705 WILLIAMS, A. W. Portable positive-pressure apparatus for administration of ether by intra-tracheal insufflation. *Jour. A. M. A.*, p. 138, Jan. 9, 1915.
- 1706 WOLFSON, W. L. Improved ether mask. *Jour. A. M. A.*, p. 823, March 6, 1915.

IX. MISCELLANEOUS.

- 1707 AGNEW, F. F. Diseases of the eye and ear as influenced by general conditions or diseases of other organs. *Iowa State Med. Soc. Jour.*, Dec., 1915.
- 1708 ALBRECHT, W. Gunshot wounds of the neck. Über Schussverletzungen des Halses). *Arch. f. Ohrenheilk.*, Vol. 98, Nos. 2-3, p. 138, Aug., 1915.
- 1709 ALEXANDER, G. J. Influenzal infections of the nose, throat and ear. *Jour. of Ophth., Otol. and Laryngol.*, p. 231, Mar 'h, 1915.

- 1710 ALLPORT, F. State legislation concerning the examination of school children's eyes, ears, nose and throats. *Ophthalmology*, July, 1915.
- 1711 ANDERSON, J. Case of polyglandular syndrome with adrenal hypernephroma and adenoma of the pituitary gland, both of small size. *Glasgow Med. Jour.*, p. 178, March, 1915.
- 1712 ARCE, JULIAN. Leishmaniasis of Peru (with report of cases showing lesions of face and mouth and destruction of the nose). *Revista Espanola de Urolog. y Dermatolog.*, p. 503, Sept., 1915.
- 1713 AUBERTIN, C. and CHABANIER, H. Differential leucocyte count in mumps. *Arch. des Mal. du Coeur*, p. 1, Jan.-Feb., 1915.
- 1714 BABCOCK, W. W. Osteoplastic surgery of the face. *Jour. A. M. A.*, Vol. 64, 1915.
- 1715 BADERSCHER, J. A. Development of the thymus in the pig: (1) Morphogenesis. *Amer. Jour. Anat.*, March, 1915.
- 1716 BADERSCHER, J. A. Development of the thymus in the pig: (2) Histogenesis. *Amer. Jour. Anat.*, May, 1915.
- 1717 BARNDOLLAR, W. P. Surgical diseases of the superior maxilla. *Ann. of Otol., Rhinol. and Laryngol.*, p. 516, Sept., 1915.
- 1718 BASILE, G. Experimental sporotrichosis with special reference to oto-rhino-laryngology (Sporotricosi sperimentale con particolare riguardo alla oto-rino-laringologia). *Ann di Clinica medica*, Anno 6, fasc. 2, 1915.
- 1719 BECK, J. C. Borderland topics in ophthalmology and oto-laryngology. *Pa. Med. Jour.*, p. 251, Jan., 1915, and *Ind. State Med. Ass'n Jour.*, No. 5, May, 1915.
- *1720 BECK, J. C. and HASSIN, G. B. A case of combined extracranial paralysis of cerebral nerves. *Med. Rec.*, p. 308, Aug. 21, 1915.
- 1721 BECKLEY, C. C. Border-line cases of mental defect with special reference to hysterical symptoms. *Bost. Med. and Surg. Jour.*, No. 14, April 8, 1915.
- 1722 BELBEZE, R. The use of colloidal gold in surgery. *Bull. de l'Acad. de Med.*, Jan. 19, 1915.
- 1723 BENJAMINS, C. E. and ROCHAT, G. F. Physiology of the lacrymal apparatus. *Nederl. Tijdschr. v. Geneesk.*, No. 18, May 1, 1915.
- 1724 BILANCIONI, G. Unedited letters of Leopoldo Marcantonio Cالداني. *Arch. ital. di otol., rinol. e laringol.*, Vol. 26, No. 3, p. 234, June, 1915.
- 1725 BILANCIONI, G. Unedited letters of Leopoldo Marcantonio Cالداني. *Arch. ital. di otol., rinol. e laringol.*, Vol. 26, No. 3, p. 234, June, 1915.
- 1726 BILANCIONI, G. Unedited letters of Domenico Cotugno. *Arch. ital. di otol., rinol. e laringol.*, Vol. 26, No. 1, p. 47, Jan., 1915.
- 1727 BILANCIONI, G. Intestinal parasites in oto-laryngology (Importanza degli eminti intestinale in oto-rino-laringoiatria. Le tonsille "da vermi"). *Boll. delle Mal. dell' Orecchio*, p. 230, Vol. 33, No. 11, Nov., 1915.
- 1728 BLAIRE, V. P. Instances of operative correction of mal-relation of the jaws. *Internat. Jour. Orthodon.*, Vol. 1, No. 8, p. 395, Aug., 1915.
- 1729 BORDLEY, J. Arthritis associated with infections of the nose and throat. *South. Med. Jour.*, Oct., 1915.

- 1730 BOSTON, L. N. Hypopituitarism. *N. Y. Med. Jour.*, p. 884, Oct. 30, 1915.
- 1731 BOWERS, E. F. Why are coughs? *Jour. of Outdoor Life*, March, 1915.
- 1732 BRENNAN, J. A. O. Clinical experience with silvol, a new silver preparation. *L'ville Monthly Jour. of Med. and Surg.*, p. 331, April, 1915.
- 1733 BRINKLEY, A. S. Rectal anesthesia. *Va. Med. Semi-Monthly*, p. 419, Dec. 10, 1915.
- 1734 BROWN, P. Desirability of complete immobilization in roentgenization of the head. *Amer. Jour. Roentgenol.*, p. 680, March, 1915.
- 1735 BRUCE, A. N. Paralysis of the spinal accessory nerve following many years after the removal of tuberculous glands from the neck. *Rev. Neurol. and Psychiat.*, p. 51, Feb., 1915.
- 1736 BRUNI, A. C. Origin and development of hypophyseal peduncle (Sul' origine e sullo sviluppo del peduncolo faringoipofisario nell'uomo). *Arch. ital. di Otol., Rinol. e Laringol.*, p. 475, Vol. 26, fasc. 6, Dec., 1915.
- 1737 BRYAN, C. W. G. Plastic operation for closing certain unhealed gunshot wounds of the head. *Lancet*, Sept. 11, 1915.
- 1738 BRYANT, W. S. The dependence of tuberculosis upon respiratory insufficiency of the upper air tract. *Med. Rec.*, p. 270, Aug. 14, 1915.
- 1739 BUNTING, R. C. The deep injection of alcohol for tic douloureux. *Memphis Med. Monthly*, June, 1915.
- 1740 BURDICK, A. F. Physiologic physics in relation to eye and ear. *Mich. State Med. Soc. Jour.*, Jan., 1915.
- 1741 BUTT, W. R. Nose, throat and ear as neighboring organs to the teeth. *Penn. Med. Jour.*, Vol. 18, No. 11, Aug., 1915.
- 1742 BUYS, E. The scope of oto-laryngology (Metodi e scopi della specialita oto-rino-laringologica). *Arch. ital. di otolog.*, p. 417, Vol. 26, No. 5, Oct., 1915.
- 1743 BUYS, E. Stefanini-Gradenigo method of inhalations in therapy (Le inalazioni saline secche in terapia col metodo dei Profs. Stefanini e Gradenigo). *Arch. ital. di Otol., Rinol. e Laringol.*, p. 499, Vol. 26, Fasc. 6, Dec., 1915.
- 1744 BYRNES, C. M. Clinical and experimental studies upon the injection of alcohol into the Gasserian ganglion for the relief of trigeminal neuralgia. *Bull. Johns Hopkins Hosp.*, Jan., 1915.
- 1745 CALDERA, C. Hemostasis in oto-laryngology (I mezzi emostatici in otorino-laringoiatria). *Boll. delle Mal. dell'Orecchio*, p. 217, Vol. 33, No. 10, Oct., 1915.
- 1746 CAMPBELL, W. F. Tuberculous glands of the neck; osteomyelitis. *Med. Times*, p. 118, June, 1915.
- 1747 CARMODY, T. E. and FINNOFF, W. C. A report of a study of coagulation in relation to operative procedures. *Denver Med. Times*, March, 1915.
- 1748 CAPELL, W. L. Catarrh of the air passages. *Med. Summary*, Jan., 1915.

- 1749 CARY, E. H. Malignant tumors in and around the orbit with report of starvation method of treatment. *Jour. A. M. A.*, p. 1540, Oct. 30, 1915.
- 1750 CAVADIE, DR. Treatment of fracture of lower jaw (Méthode directrice rationnelle dans le traitement des fractures du maxillaire inférieure. *Bull. de l'Acad. de Med., Paris*, Aug. 10, 1915.
- 1751 CERQUEIRA, D. Improved technic for staining tubercle bacilli (Sobre um novo methodo de coloracao do bacillo de Koch). *Brazil Medico*, Oct. 30, 1915.
- 1752 CLAPP, A. B. Nasal or oral infection as related to systemic disease. *Med. Cent.*, June, 1915.
- 1753 CLARK, J. S. Some infections of the head and their causes. *Ill. Med. Jour.*, Aug., 1915.
- 1754 COATES, G. M. Vaccine therapy in diseases of the nose, throat and ear. *N. Y. Med. Jour.*, July 10, 1915.
- 1755 COLLIER, C. Tumor of carotid body. Report of case. *Surg. Gynecol. and Obstetr.*, p. 484, April, 1915.
- 1756 COLLIER, M. Deformities of the face and upper and lower jaws due to nasal obstruction. *Intern. Jour. of Orthodontia*, p. 58, Feb., 1915.
- 1757 COOKE, C. T. Original research in ophthalmology and otolaryngology on Pacific coast. *Ophthal.*, p. 237, Jan., 1915.
- 1758 COTT, G. F. Harmless post-operative temperature. *Ann. Otol., Rhinol. and Laryngol.*, March, 1915.
- 1759 COUES, W. P. Gummatous cervical adenitis; its relation to and mistaken diagnosis for tuberculous cervical adenitis. *Bost. Med. and Surg. Jour.*, Nov. 18, 1915.
- 1760 CRANE, C. G. Bacterial vaccines in ear, nose and throat infections. *Long Island Med. Jour.*, Oct., 1915.
- 1761 CRENSHAW, WM. The nose, throat and ear as neighboring organs to the teeth. *Atlanta Jour.-Rec. of Med.*, July, 1915.
- 1762 CROUZON, O. Malformation of skull and face with exophthalmos and divergent strabismus in mother and son (Dysostose cranio-faciale héréditaire). *Arch. de Med. des Enfants*, Oct., 1915.
- 1763 CROUZON, O. Another family with malformation of skull and face in seven members in two generations. *Arch. de Med. des Enfants*, Oct., 1915.
- 1764 CUSHING and GROETSCH. Hibernation and the pituitary body. *Jour. Exp. Med.*, July, 1915.
- 1765 DABNEY, V. The above of surgery in diseases of the ear, nose and throat. *Med. Rev. of Rev.*, p. 651, Nov., 1915.
- 1766 DANDY, W. E. Extripation of pineal body. *Jour. Exp. Med.*, Vol. 22, No. 2, Aug., 1915.
- 1767 DEBIERNE. Localization of foreign bodies by radioscopia. *Presse Med.*, March 4, 1915.
- 1768 DECHERD, H. B. Some results of nose and throat operations in chronic poisoning of heart, lungs, kidneys, joints, etc. *South. Med. Jour.*, May, 1915.

- 1769 DIMITRIADIS, D. S. Wounds of the ears, nose and larynx during the Greek wars of 1912-1913 (Über Verwundungen an den Ohren, der Nase und dem Kehlkopf aus den beiden letzten Kriegen Griechenlands, 1912-1913). *Monatschr. f. Ohrenheilk.*, p. 353, 1915.
- 1770 DIXON, W. E. Nose, anatomically considered with special reference to reflexes and constitutional diseases. *Okla. State Med. Ass'n Jour.*, March, 1915.
- 1771 DOCK, G. Thymus gland. *Ohio State Med. Jour.*, Sept., 1915.
- 1772 DOLMAGE, G. F. Case of hemophilia treated with horse serum. *Iowa State Med. Soc. Jour.*, March, 1915.
- 1773 DONALDSON, H. H., HATAI, S. and KING, H. D. Postnatal growth of brain under several experimental conditions. *Jour. Nervous and Mental Dis.*, Dec., 1915.
- 1774 DONATH, J. Treatment of trifacial neuralgia with alcohol in-
- 1775 DOWNING, D. F. Anesthesia, local and general, in operations on the nose, throat and ear. *Clinique*, July, 1915.
- 1776 DRINKER, K. R. and C. K. Factors affecting coagulation time of blood. Effects of rapid progressive hemorrhage on factors of coagulation. *Am. Jour. of Physiol.*, Feb., 1915.
- 1777 DUNNING, H. S., McWILLIAMS, C. A. and MITCHELL, V. E. Osteomyelitis of the lower jaw. *Surg., Gyn. and Obst.*, p. 306, Vol. 21, No. 3, Sept., 1915.
- 1778 DUPUY, H. Reduction of preventable disasters in ear, nose and throat surgery. *South. Med. Jour.*, Jan., 1915.
- 1779 EARLAM, H. Medical inspection of children. *Volta Rev.*, June, 1915.
- 1780 ENGELHARDT, DE. Prognosis of gunshot wounds of the skull (Zur Prognose der Schädelgeschüsse. *Muench. med Woch.*, p. 1096, No. 32, Aug. 10, 1915.
- 1781 EVANS, J. H. Ether insufflation anesthesia. *N. Y. Med. Jour.*, Jan. 30, 1915.
- 1782 FENTON, R. A. Oxycephalus. *Jour. A. M. A.*, Oct. 30, 1915.
- §1783 FINCKH, L. Early diagnosis of gas-phlegmons by X-ray (Die frühzeitige Erkennung der Gasphlegmone durch das Roentgenbild). *Deut. med Woch.*, May 1, 1915, *THE LARYNGOSCOPE*, p. 137, Feb., 1916.
- 1784 FLEISCH, M. von. Experimental thymus studies. *Beitr. z. klin. Chir.*, Vol. 95, p. 376, 1915.
- 1785 FONIO, O. Comparative study of the blood platelets. *Corresp. Blatt. f. Schweizer Aerzte*, Nos. 48-49, Nov. 27-Dec. 4, 1915.
- 1786 FOOTE, E. M. Partial resection of the lower jaw for cancer. *Amer. Jour. Med Sc.*, p. 1, Vol. 150, No. 1, July, 1915.
- 1787 FORNANI, A. Empirical remedies in diseases of the ear, nose and throat (Dei rimedi empirici usati nelle puglie per la cura delle malattie dell'orecchio del naso e della gola). *Boll. delle Mal. dell'Orecchio*, p. 257, Dec., 1915.
- 1788 FRANK, J. Brain surgery in war. *Military Surgeon*, Nov., 1915.
- 1789 FRANK, E. Essential thrombopenia (Konstitutionelle Purpura—Pseudo-Hämophilie). *Berl. klin. Woch.*, Nos. 18-19, 1915.

- 1790 FRASER, J. S. Tuberculous cervical adenitis in children. *Clin. Jour.*, Vol. 44, p. 41, 1915.
- †1791 FREUNDLICH, D. B. The teeth as a primary factor in diseases of the ear, nose and throat—the diagnostic value of co-operation of the otologist, rhinologist, laryngologist with the dentist. *THE LARYNGOSCOPE*, p. 40, Jan., 1915.
- 1792 FREY, LEON. Corrective surgery in fractures of the jaw and mutilations of the face (Un aperçu la prothèse restauratrice dans les fractures des Machoires et mutilations de la face). *Paris Medical*, p. 282, Aug. 21, 1915.
- 1793 FRIEDLAND, M. O. Dermoid of the frontal bone. *Russky Vrach*, Sept. 26, 1915.
- 1794 FRIEDMAN, H. M. Physical basis for moral and mental deficiencies. *Med. Record*, p. 637, April 17, 1915.
- 1795 FROESCHIELS, E. Re-education after brain injury (Übungsschulen für Gehirnrüppel). *Muench. med. Woch.*; No. 27, July 6, 1915.
- 1796 FUNK, V. A. Congenital cysts of the neck. *Amer. Jour. Surg.*, Vol. 29, No. 8, p. 290, Aug., 1915.
- 1797 GALMOZZI, F. A. Anemia and lymphatism cured by dry inhalations (Ricerche emometriche in alunni anemici e limfatici delle Scuole Elementari di Bergamo curati con le inalazioni di nebbie secche sistema Stefanini-Gradenigo). *Arch. ital. di Otol., Rinol. e Laringol.*, p. 495, Vol. 26, Fasc. 6, Dec., 1915.
- 1798 GIRARD, L. Subcutaneous emphysema in measles. *Med. Rev.*, p. 135, April, 1915.
- 1799 GLOGAN, O. Disease of ear and upper respiratory tract among American factory workers. *N. Y. Med. Jour.*, p. 616, March 27, 1915.
- 1800 GRANT, J. D. Facial plastic surgery, laryngology and stomatology in French military hospitals. *Lancet*, May 1, 1915.
- 1801 GREY, E. G. Fibrin as a hemostat in cerebral surgery. *Surg., Gyn. and Obst.*, p. 452, Vol. 21, No. 4, Oct., 1915.
- 1802 GRIFFITH, STANLEY. Cervical gland tuberculosis. *Lancet*, June 19, 1915.
- 1803 GROSSMAN, J. Enlargement of the thymus. *N. Y. Med. Jour.*, p. 1089, Nov. 27, 1915.
- †1804 GWATHMEY, J. T. Some facts not generally understood about oil-ether colonic anesthesia. *THE LARYNGOSCOPE*, p. 692, Oct., 1915.
- †1805 HALSTEAD, T. H. Somnoform as an anesthetic. *THE LARYNGOSCOPE*, p. 562, Aug., 1915.
- 1806 HALSTEAD, T. H. Report of progress of a case of pituitary tumor reported in 1914. *Trans. Amer. Laryngol. Ass'n*, p. 313, 1915.
- 1807 HAMMOND, R. The role of the nose, throat and accessory sinuses in the etiology of chronic infectious arthritis. *Jour. A. M. A.*, p. 1091, Sept. 25, 1915.
- 1808 HARBITZ, F. Cystic tumor of the maxillae and especially on adamantine denosus. *Dent. Cosmos*, Oct., 1915.
- 1809 HARMAN, N. B. Education of children with defective vision. *Brit. Jour. Child. Dis.*, Vol. 12, No. 141, Sept., 1915.

- 1810 HARMER, W. D. Cases illustrating results obtained by use of Simpson light. *Proc. Royal Soc. Med., Laryngol Section*, Vol. 9, No. 2, Dec., 1915.
- 1811 HASELTINE, B. Modern views of syphilis. *Jour. of Ophth., Otol. and Laryngol.*, p. 243, March, 1915.
- 1812 HAWES, J. B. Treatment of tuberculous adenitis. *Bost. Med. and Surg. Jour.*, Jan. 7, 1915.
- 1813 HECKER, F. The development of the head. *Internat. Jour. Orthodon.*, p. 515, Oct., 1915.
- 1814 HECKER, F. Review of some studies of rachitic skulls of monkeys. *Internat. Jour. Orthodon.*, p. 612, Dec., 1915.
- 1815 HESS, A. F. Protective therapy for mumps. *Amer. Jour. Dis. Child.*, Vol. 10, No. 2, Aug., 1915.
- *1816 HESS, Hemophilla calcipriva. *Johns Hopkins Hosp. Bull.*, Nov., 1915.
- 1817 HILL, R. F. Use of heated ether and oxygen in nose, throat and ear operations. *Jour. Ophth., Otol. and Laryngol.*, p. 56, Jan., 1915.
- 1818 HILL, W. Shell wound of neck. *Proc. Roy. Soc. Med., Laryngol. Sect.*, p. 123, Vol. 8, No. 8, June, 1915.
- 1819 HILL-AITKEN, C. J. A note on injuries to the head. *South African Med. Rec.*, p. 128, Vol. 13, 1915.
- 1820 HIRSCHHELL, G. Obstinate case of trifacial neuralgia cured by injections of alcohol into the Gasserian ganglion. *Muench. med. Woch.*, Bd. 62, H. 5, 1915.
- 1821 HIRSCHFELDER, A. D. Brain lipoid as a hemostatic (Gehirn-lipoid als Hämostatikum). *Berl. klin. Woch.*, No. 37, 1915.
- 1822 HOFFMANN, Memorial to Oskar Brieger. *Ztschr. f. Laryngol.*, Bd. 7, Heft 4, p. 457, 1915.
- 1823 HOLLAND, C. T. Stone in submaxillary gland. *Arch. of Radiol. and Electrotrother*, June, 1915.
- 1824 HOLT, E. E. Economics of eye, ear, nose and throat, together with economics of entire body (physical economics). *Ophthalmology*, Vol. 11, No. 4, July, 1915.
- 1825 HOPKINSON, D. Status lymphaticus. *Wis. Med. Jour.*, p. 311, Jan., 1915.
- 1826 HOUGHTON, H. A. An adult with status thymicolymphaticus of the hetero-sexual chromaffin type. *Med. Rec.*, p. 509, Sept. 25, 1915.
- 1827 HOUGHTON, H. A. Status lymphaticus associated probably with an inferior thyroid lobe. *N. Y. Med. Jour.*, p. 887, Oct. 30, 1915.
- 1828 HOUGHTON, H. A. Hyperplastic thymus. *Jour. A. M. A.*, July 24, 1915.
- 1829 HOWARTH, W. Bullet in pterygoid region of skull. *Proc. Roy. Soc. Med., Laryngol. Sect.*, p. 130, Vol. 8, No. 8, June, 1915.
- 1830 HOWELL, H. L. Indications for use of pituitary extract. *Ill. Med. Jour.*, Jan., 1915.
- 1831 HUBENY, M. J. Roentgenography of the skull. *Illinois Med. Jour* Nov., 1915.
- †1832 HURD, L. M. Case of an excessive bleeder. *THE LARYNGOSCOPE* p. 856, Dec., 1915.

- 1833 HUTCHINSON, J. M. Systemic disease due to focal infections. *Old Dom. Jour.*, Oct., 1915.
- 1834 IBERSHOFF, A. E. Toxic uveitis following vaccination. *Jour. Ophth., Otol. and Laryngol.*, p. 121, Feb., 1915.
- 1835 IVY, R. H. Mesothelial tumors of the jaws. *Jour. A. M. A.*, p. 40, Jan. 2, 1915.
- 1836 JONES, N. Animal extracts in ophthalmology. *Can. Med. Ass'n Jour.*, Aug., 1915.
- 1837 JUDD, A. Use of decanted blood serum after severe hemorrhage and in hemophilia; report of two cases. *Med. Rec.*, No. 15, April 10, 1915.
- 1838 KAEMPFFER, L. G. Blood platelet extract, a physiological hemostatic in nose and throat surgery. A preliminary report. *Amer. Jour. Surg.*, p. 401, Nov., 1915.
- *1839 KAHN, H. and GORDON, L. E. The use of pituitary extract as a coagulant in surgery of the nose and throat. *Jour. A. M. A.*, p. 301, Jan. 23, 1915, and *Ann. Otol., Rhinol. and Laryngol.*, p. 322, Vol. 24, No. 2, June, 1915.
- 1840 KASSEL, K. Rhinology in the eighteenth century (Die Nasenheilkunde des 18 Jahrhunderts). *Ztschr. f. Laryngol. u. Rhinol.*, Bd. 7, H. 5, April, 1915.
- 1841 KASSEL, K. Rhinology in the nineteenth century (Die Nasenheilkunde des 19 Jahrhunderts). *Zeitschr. f. Laryngol.*, p. 695, Vol. 7, H. 6, Sept., 1915.
- 1842 KATZ, L. Localization of metal foreign bodies by roentgen stereogram (Der Salow'sche Tiefenmesser). *Berl. klin. Woch.*, Vol. 52, No. 29, July 19, 1915.
- 1843 KEEN, W. W. Dangers of ether as an anesthetic. *Bost. Med. and Surg. Jour.*, No. 23, Dec. 2, 1915.
- 1844 KINCAID, J. W. Tuberculosis of cervical glands. *Ky. Med. Jour.*, June, 1915.
- 1845 KING, J. J. Diseases of the nose, throat and ear from the practitioner's standpoint. *Internat. Jour. Surg.*, Vol. 38, p. 111, April, 1915.
- 1846 KING, H. D. Epidemiology of mumps. *Atlanta Jour. Rec. of Med.*, April, 1915.
- 1847 KIRKLAND, E. A country practitioner's treatment of colds. *Jour. Ophthalmol., Otol. and Laryngol.*, p. 1016, Dec., 1915.
- 1848 KLINEDINST, J. F. Intrarectal use of sodium salicylate in uveitis. *Penn. Med. Jour.*, p. 28, Oct., 1915.
- 1849 KLINGER, R. Coagulation of the blood. *Corresp. Blatt. f. Schweizer Aerzte*, No. 51, Dec. 18, 1915.
- 1850 KOLB, K. Plastic operation on the face by means of autoplasic transplantation of fat. *Zentrablatt. f. Chir.*, p. 427, Vol. 42, 1915.
- 1851 KONIKOW, M. J. Pituitary extract as a hemostatic in hemorrhages of the lungs. *Bost. Med. and Surg. Jour.*, Sept. 20, 1915.
- 1852 KOERNER, O. The eyebrows in peripheral facial paralysis (Die Stellung der Augenbrauen bei der peripheren Fazialislähmung). *Arch. f. Ohrenheilk.*, Bd. 72, H. 4, May, 1915.

- 1853 KRESS, G. H. Tuberculosis in relation to the eye, ear, nose and throat. *Zeitschr. f. Tuberculose*, Vol. 24, No. 2, June, 1915.
- 1854 KUPFERLE and V. SZITY. Radio therapy in tumors of the pituitary (Über Strahlentherapie bei Hypophysentumoren). *Deutsche med. Woch.*, No. 31, 1915.
- 1855 KUWABARA, K. Facial furuncle or carbuncle. *Sei-I-Kwai Med. Jour.*, Tokyo, Dec., 1915.
- 1856 LABOURAE. Fracture of the inferior maxilla (Fracture du maxillaire inferieur; suture avec la plaque de Lane). *Rev. de Laryngol., d'Otol. et de Rhinol.*, p. 283, Nos. 44-45, Nov. 15, 1915.
- 1857 LACK, L. The influence of nasal obstruction on abnormalities of the jaws. *Brit. Jour. Dent. Science*, Feb. 15, 1915.
- 1858 LASAGNA, F. "Coagulen-Kocher-Fonio" in otolaryngology. *Arch. ital. di Otol., Rinol. e Laringol.*, Vol. 26, April, 1915.
- 1859 LAEWEN, A. Cranial gunshot wounds (Einige Beobachtungen über Schädelschussverletzungen im Feldlazarett). *Muench. med. Woch.*, p. 589, No. 17, April 27, 1915.
- 1860 LEBOUTILLIER, T. Hypertrophy of the thymus and thymus death, with a report of cases. *Arch. Pediat.*, p. 322, May, 1915.
- 1861 LESZYNSKY, W. M. Pituitary tumor with improvement after puncture of the corpus callosum. *N. Y. Med. Jour.*, p. 435, Aug. 28, 1915.
- 1862 LEWISOHN, R. Citrate method of blood transfusion. *Surg., Gyn. and Obst.*, July, 1915.
- 1863 LINNELL, E. H. The use of high frequency currents in special work. *Jour. Ophthal., Otol. and Laryngol.*, p. 847, Oct., 1915.
- 1864 LISCHER, B. E. The classification of dento-facial deformities. *Intern. Jour. Orthodontia*, June, 1915.
- 1865 LLOYD, R. I. Eye and ear surgery and prothesis in 1634. *Jour. Ophthal., Otol. and Laryngol.*, p. 765, Sept., 1915.
- 1866 LOW, W. STUART. Dyspnea, perichondritis, tumor (?) of the mediastinum. *Proc. Roy. Soc. Med., Laryngological Section*, Vol. 8, No. 5, March, 1915.
- 1867 LOW, W. STUART. Bullet wounds of the face and neck. *Proc. Royal Soc. Med., Laryngol. Section*, Vol. 9, No. 2, Dec., 1915.
- 1868 LOW, W. STUART. Soldiers kicked in neck by a horse. *Proc. Roy. Soc. Med., Laryngol. Section*, p. 123, Vol. 8, No. 8, June, 1915.
- 1869 LUEDIN. Mikulicz disease and its treatment by x-rays. *Berl. klin. Woch.*, No. 23, 1915.
- 1870 LUETSCHER, J. A. Bacteriologic and clinical study of non-tuberculous infections of respiratory tract with special reference to sputum cultures as a means of diagnosis. *Arch. Int. Med.*, Oct., 1915.
- 1871 LUMBARD, J. E. Oil-ether colonic anesthesia. *Surg., Gyn. and Obstet.*, May, 1915.
- 1872 LYONS, RANDOLPH. Case of purpura hemorrhagica with accidental complications following subcutaneous use of horse serum. *N. O. Med. and Surg. Jour.*, Aug., 1915.
- 1873 MACFARLAN, D. Relative proportion of cases in a nose and throat clinic, based on 1,000 cases. *Jour. of Ophth., Otol. and Laryngol.*, p. 253, March, 1915.

- 1874 MACKENZIE, G. W. The relationship of dental pathology to eye, ear, nose and throat conditions. *Jour. Ophth., Otol. and Laryngol.*, May, 1915.
- 1875 MACY, MARY S. A possible role for the thymus gland. *Woman's Med. Jour.*, May, 1915.
- 1876 MANCIOLI, T. Prophylaxis of ear, nose and throat diseases in schools. *Policlínico*, May 23, 1915.
- 1877 MARAGLIANO, V. Roentgen localization of foreign bodies. *Rif. Med.*, Vol. 31, No. 26, June 26, 1915.
- 1878 MARAGLIANO, V. Improved roentgen technic for locating foreign bodies. *Gaz. degli Asped. e delle clin.*, Vol. 36, No. 48, June 17, 1915.
- 1879 MARAGLIANO, V. Method of localization of metal foreign bodies. *Policlínico*, Vol. 22, No. 31, Aug. 1, 1915.
- 1880 MARKHAM, H. C. Gleanings from principal European eye and ear clinics. *Kans. Med. Soc. Jour.*, Feb., 1915.
- 1881 MATSUMOTO, T. Syringo-mylitic disease of the cranial nerves with special reference to the ear, nose, larynx and esophagus (Beitrag zur Lehre von der syringo myelitischen Erkrankung der Hirnnerven mit besonderer Berücksichtigung des Ohres, der nase, des Kehlkopfes und des Schlundes). *Passows' Beitrage*, p. 212, Vol. 8, No. 4, Aug. 12, 1915.
- 1882 MCCORD, CAREY P. The pineal gland in relation to somatic, sexual and mental development. Second paper. *Jour. A. M. A.*, p. 517, Aug. 7, 1915.
- 1883 MCLEAN, WM. Experimental investigations concerning intra-ocular pressure. *Jour. Ophthal. and Oto-laryngol.*, p. 471, June, 1915.
- 1884 MEANS, J. H. Studies on basal metabolism in obesity and pituitary disease. *Jour. of Med. Research*, March, 1915.
- 1885 MELCHIOR, DE. Latent infection in its surgical significance (Über den Begriff der ruhenden Infektion in seiner Bedeutung für die Chirurgie). *Berl. klin. Woch.*, No. 5, 1915.
- †1886 METZENBAUM, M. Scopolamin in nose and throat operations. *THE LARYNGOSCOPE*, p. 95, Feb., 1915.
- 1887 MEYER, L. B. The conservative treatment of enlarged tuberculous glands of the neck. *Med. Rec.*, July 10, 1915.
- 1888 MICK, W. H. Radiography of the face and jaw. *West. Med. Rev.*, p. 211, April, 1915.
- 1889 MIEBS, E. M. Common head injuries. *Amer. Jour. Surg.*, p. 399, Nov., 1915.
- 1890 MILLER, C. M. Further report of vaccine therapy in oto-laryngology. *Va. Med. Semi-Monthly*, p. 582, March 12, 1915.
- 1891 MILLER, R. H. The approach to the Gasserian ganglion. *Memphis Med. Monthly*, Aug., 1915.
- §1892 MILLIGAN, W. Dentigerous cyst of lower jaw. *Proc. Laryngol. Section, Royal Soc. Med.*, Vol. 8, No. 6, April, 1915. *THE LARYNGOSCOPE*, p. 123, Feb., 1916.
- 1893 MILLIGAN, W. and WESTMACOTT, F. H. Warfare injuries and neuromes. *Proc. Roy. Soc. Med., Laryngological Section*, p. 109, June, 1915, and *Jour. Laryngol., Rhinol. and Otol.*, p. 297, Vol. 30, No. 8, Aug., 1915.

- 1894 MINOT, G. R. Effect of chloroform on factors of coagulation. *Amer. Jour. Physiology*, Vol. 39, No. 2, Dec., 1915.
- 1895 M'KENZIE, D. Some points of common interest to the rhinologist and the orthodontist. *Intern. Jour. of Orthodontia*; p. 9, Jan., 1915.
- 1896 MOORE, R. Trifacial neuralgia; its successful treatment without major operation. *Arizona Med. Jour.*, Dec., 1915.
- 1897 MORESTIN, H. Vicious scars following severe burn of the face by acid; autoplasmic operation using skin from thorax and mammary gland. *Bull. et Mem. Soc. de chir. de Paris*, p. 1381, Vol. 41, 1915.
- 1898 MORSE, D. H. Color changes and the margin of safety in nitrous oxide anesthesia. *Amer. Jour. Surg., Anesthesia Supp.*, p. 133, Oct., 1915.
- 1899 MOSER, H. and ARNSTEIN, A. Mumps in men over thirty; seventeen cases in one troop. *Wien. klin. Woch.*, Vol. 28, No. 24, June 17, 1915.
- 1900 MOSKOWITZ, S. The newer therapeutics in otology. *N. Y. Med. Jour.*, p. 354, Aug. 14, 1915.
- 1901 MOSS, C. A. Head colds; parts involved and some of the results. *Kentucky Med. Jour.*, Dec., 1915.
- 1902 MOURE, E. J. Oto-rhino-laryngology in war time. *Presse Med.*, Feb., 18, 1915, and *Rev. de Laryngol., d'Otol. et de Rhinol.*, p. 281, Nos. 44-45, Nov. 15, 1915.
- 1903 MOURE, E. J. Wounds of ear, nose and throat. *Bull. de l'Acad. de Med.*, Feb. 16, 1915.
- 1904 MUELLER, P. Flap from the sternum to close gap in skull; two cases. *Zentralb. f. Chir.*, Bd. 42, No. 23, June 5, 1915.
- 1905 MUELLER, S. Treatment of post-operative parotitis. *Hospitalstundende*, Oct. 6, 1915.
- 1906 MUELLER, W. B. Operative treatment of gunshot wounds of the brain (Verletzungen des Gehirns). *Arch. f. klin. Chir.*, Vol. 107, No. 1, 1915.
- 1907 MYERS, D. W. Relationship between rhinology and orthodontia. *Intern. Jour. of Orthodontia*, p. 86, Feb., 1915.
- 1908 MYERS, D. W. Streptococcus question. *Jour. Ophth., Otol. and Laryngol.*, p. 283, April, 1915.
- 1909 NORDMANN. The thymus gland. (Experimentelles und Klinisches über die Thymusdrüse). *Arch. f. klin. Chir.*, Bd. 106, H. 1, 1915.
- 1910 O'MALLEY, J. F. Bullet wound in the head. *Proc. Roy. Soc. Med., Laryngol. Sect.*, p. 128, Vol. 8, No. 8, June, 1915.
- 1911 PARKINSON, DR. Suppurative parotitis following pneumonia. *Brit. Jour. Child. Dis.*, May, 1915.
- 1912 PARRISH, H. Relation of diseases of nose and throat to general practice of medicine. *N. Y. Med. Jour.*, May 15, 1915.
- 1913 PETERS, DR. Wounds of the neck (Zwei interessante Halsverletzungen). *Muench. med. Woch.*, p. 1134, No. 33, Aug. 17, 1915.
- 1914 PETERSON, W. A. The special sense organs in life insurance. *Lan.-Clin.*, p. 45, July 10, 1915.

- 1915 PFINGST, A. O. Relation of the lymphatic glands to diseases of the eye, ear, nose and throat. *L'ville Med and Surg. Jour.*, p. 228, Jan., 1915.
- 1916 PFISTER, F. Infections of the nose, throat and accessory sinuses. Their importance to the general practitioner. *Wis. Med. Jour.*, p. 391, March, 1915.
- 1917 PHILIBERT, O. Relation between nose breathing and pulmonary tuberculosis. *Sem. Med., Buenos Aires*, April 1, 1915.
- 1918 PHILLIPS, W. H. Colds: their etiology and treatment. *Jour. Ophthalmol., Otol. and Laryngol.*, p. 1012, Dec., 1915.
- 1919 PIFFL, O. Diseases of the ear and nose in war. *Wien. klin. Woch.*, No. 13, 1915.
- 1920 PODTLAGINA, V. Treatment of fractured jaw. *Russky Vrach*, Vol. 14, No. 20, 1915.
- 1921 POLLOCK, L. J. Hypo-pituitarism in chronic hydrocephalus. *Jour. A. M. A.*, p. 395, Jan. 30, 1915.
- 1922 PONZO, M. Graphic registration of respiration (Nuovi dispositioi per la registrazione grafica del respiro). *Arch. ital. di otol., rinol. e laringol.*, p. 302, Vol. 26, No. 4, Aug., 1915.
- 1923 PONZO, M. Modifications of respiration during mental reading and their significance (Modificazioni del respiro durante la lettura mentale e loro significato). *Arch. ital. di otol., rinol. e laringol.*, p. 306, Vol. 26, No. 4, Aug., 1915.
- 1924 PORTEUS, R. W. Nose and throat in etiology of general diseases. *Wisconsin Med. Jour.*, Oct., 1915.
- 1925 POTTS, W. A. What tests in childhood are best calculated to throw light upon the capacities of mental defectives for future work? *Lancet*, July 17, 1915.
- 1926 PUSATERI, S. Statistics of the ear, nose and throat clinic for 1914 (Resoconto statistico dal 1 Gennaio al 31 Dicembre, 1914). *Arch. ital. di otol., rinol. e laringol.*, Vol. 26, No. 3, p. 213, June, 1915.
- 1927 PUSATERI, S. Pappillomata in oto-rhino-laryngology (I papillomi nel campo della oto-rino-laringologia). *Arch. ital. di otol., rinol. e laringol.*, Vol. 26, No. 3, p. 210, June, 1915.
- *1928 RAMOND, F. and GOUBERT, G. Some ocular complications of mumps. *Bull. et mem. Soc. med. des Hop. de Paris*, July 29, 1915.
- 1929 RANSHOFF, J. The status of cerebral surgery. *Lancet-Clinic*, May 15, 1915.
- 1930 RANZI, DR. Successful operation on tumor of hypophysis. *Wiener klin. Woch.*, No. 5, 1915.
- 1931 REBER, W. Medical treatment of pituitary disease. *N. Y. Med. Jour.*, p. 392, Feb. 27, 1915.
- 1932 REED, W. M. Chronic focal infection of nose, throat and ear as cause of general infection. *Mo. State Med. Ass'n Jour.*, No. 5, May, 1915.
- 1933 REEDE, E. H. Advances in the knowledge of the thymus gland. *Wash., Med. Ann.*, No. 4, July, 1915.
- 1934 REZNICEK, R. Unilateral multiple injuries of cerebral nerves. *Neurol. Zentralbl.*, p. 370, Vol. 34, 1915.
- 1935 RHEIN, J. H. W. Symptoms of disease of the pituitary body. *N. Y. Med. Jour.*, p. 438, March 6, 1915.

- 1936 RICE, P. A few plain truths. *Jour. Ophth., Otol. and Laryngol.*, p. 326, April, 1915.
- 1937 ROBERTS, J. E. H. Treatment of gunshot wounds of the head, with special reference to apparently minor injuries. *Brit. Med. Jour.*, p. 498, Oct. 2, 1915.
- 1938 ROBERTSON, T. B. and BURNETT, T. C. Influence of anterior lobe of pituitary on growth of carcinomata. *Jour. of Exper. Med.*, March, 1915.
- 1939 ROCCA VILLA, A. Diagnosis of tumors of the thymus (Alcune forme di neoplasia timica). *Riforma Med.*, Nos. 13-14, April, 1914.
- 1940 RODGER, T. R. Reports for the year 1914 from the ear and throat department of the Royal Infirmary, Edinburgh. *Jour. Laryngol., Rhinol. and Otol.*, April, 1915.
- 1941 ROEMHELD, L. Hemiplegia after head lacerations (Über homolaterale Hemiplegien nach Kopfverletzungen). *Muench. med. Woch.*, p. 600, No. 17, April 27, 1915.
- 1942 ROWLETTE, R. J. Vaccines in the treatment of respiratory diseases. *Lancet*, June, 26, 1915, and *Dublin Jour. Med. Sc.*, Aug., 1915.
- 1943 ROY, DUNBAR. Some statistical observations on oto-laryngological diseases among negroes based on fifteen hundred cases. *Trans. Amer. Laryngol. Ass'n*, p. 306, 1915.
- 1944 ROYCE, G. Headache in diseases of the ear, nose and throat. *Can. Practitioner and Rev.*, Jan., 1915.
- *1945 RYLAND, A. Oto-laryngological case from a military hospital. *Jour. Laryngol., Rhinol. and Otol.*, p. 304, Vol. 30, No. 8, Aug., 1915.
- 1946 RUTTIN, E. Facial paralysis (Über Parese des Mundfozialis). *Zeitschr. f. Ohrenheilk.*, p. 243, Bd. 73, H. 3, Nov., 1915.
- 1947 SACHS, E. Factors that make for better results in cranial surgery. *Jour. Mo. State Med. Ass'n*, p. 517, Dec., 1915.
- 1948 SAENGER, N. Influenza and its complications. *Med. Fortnightly*, p. 84, April 15, 1915.
- 1949 SCARLETT, R. B. Nose, throat and ear conditions, their diagnosis and treatment in general practice. *N. Y. Med. Jour.*, p. 246, Feb. 6, 1915.
- 1950 SCHEPELMANN, E. Plastic operations on the cheek. *Deut. Zeitsch. f. Chir.*, p. 270, Vol. 133, 1915.
- 1951 SCHEPELMANN, E. Tumors of the hypophysis. *Deut. Zeitsch. f. Chir.*, p. 390, Vol. 133, 1915.
- 1952 SELIKOVITCH, S. The teeth and their relation to health. *N. Y. Med. Jour.*, p. 277, Aug. 7, 1915.
- 1953 SELLARS, R. B. Treatment of trifacial neuralgia by deep alcoholic injections. *Tex. State Jour. Med.*, March, 1915.
- 1954 SENGER, W. Congenital cysts of the neck. *Colorado Med.*, May, 1915.
- 1955 SEWALL, E. C. Operation on the hypophysis according to Hirsch's method. Presentation of a case with brain, eye and other symptoms. *Jour. A. M. A.*, p. 681, Aug. 21, 1915.
- 1956 SHANKLIN, E. M. Oto-laryngology. *Ind. State Med. Ass'n Jour.* Jan., 1915.

- 1957 SHARPE, N. Herpes zoster of the cephalic extremity with special reference to the geniculate, auditory, glossopharyngeal and vocal syndromes. *Amer. Jour. Med. Sc.*, May, 1915.
- 1958 SHARPE, WM. Cranial decompression. *Amer. Jour. Med. Sc.*, p. 563, April, 1915.
- 1959 SHAW, H. L. K. and WILLIAMS, F. J. Blood coagulation in infancy. *Albany Med. Ann.*, p. 571, Dec., 1915.
- 1960 SHAXBY, J. H. Localization of foreign bodies by x-rays. *Brit. Med. Jour.*, July 3, 1915.
- 1961 SJOELANDER, A. and STRANDBERG, A. Innervation of the thymus. *Uppsala Laekarefoer. Foerhand.*, Feb. 20, 1915.
- 1962 SKINNER, E. H. Roentgenotherapy for angiomas. *Jour. Mo. State Med. Ass'n*, p. 411, Sept., 1915.
- 1963 SMITH, O. C. Differential diagnosis and indications for treatment of tumors of the neck. *Boston Med. and Surg. Jour.*, Feb. 11, 1915.
- 1964 SMITH, WM. H. High frequency in special practice. *Jour. Ophthal., Otol. and Laryngol.*, p. 850, Oct., 1915.
- 1965 SPENSE, T. B. Tuberculosis of the cervical lymph nodes. *L. I. Med. Jour.*, Jan. 15, 1915.
- 1966 STEINER, R. Inhalations (Über Inhalationstherapie). *Prag. Med. Woch.*, No. 25, 1915.
- 1967 STRANDBERG, O. The use of the universal light bath in rhinolaryngology (Über die Anwendung des universellen Lichtbades in der Rhino-Laryngologie). *Ztschr. f. Laryngol. u. Rhinol.* Bd. 7, H. 5, April, 1915.
- 1968 STREETER, G. L. Development of the venous sinuses of the dura mater in the human embryo. *Amer. Jour. Anatomy*, p. 145, Vol. 18, No. 2, Sept. 15, 1915.
- 1969 STUCKY, J. A. Relation of pathologic conditions in oto-rhinology to general medicine and surgery. *South. Med. Jour.*, Feb., 1915, and *Ky. Med. Jour.*, March 1, 1915.
- 1970 SYMINGTON, J. Relation of the inner surface of the cranium to the cranial aspect of the brain. *Edin. Med. Jour.*, p. 85, Vol. 14, 1915.
- 1971 SYRING, DR. Treatment of cranial gunshot wounds (Zur Behandlung der Schädelchüsse im Felde). *Muench. med. Woch.*, p. 592, No. 17, April 27, 1915.
- 1972 TABUTEAU, G. G. Treatment of gunshot wounds of the head based on a series of ninety-five cases. *Brit. Med. Jour.*, p. 501, Oct. 2, 1915.
- 1973 TAMEMORI, Y. Study of the thymus gland during involution age. *Virchows' Arch.*, Vol. 1207, p. 255, 1915.
- 1974 TAYLOR, E. W. Progressive vagus-gloss-pharyngeal paralysis with ptosis. *Jour. Nerv. and Mental Dis.*, p. 129, March, 1915.
- *1975 TELFORD, E. D. Acute actinomycosis of the parotid gland. *Brit. Med. Jour.*, p. 534, Oct. 9, 1915.
- 1976 THORPE, J. H. Blood pressure in diseases of eye, ear, nose and throat. *Kentucky Med. Jour.*, Nov., 1915.
- 1977 THOST, DR. Gunshot wounds of the neck (Über Halsschüsse). *Zeitschr. f. Ohrenheilk.*, p. 207, Bd. 73, H. 3, Nov., 1915.

- 1978 THURSFIELD, H. Status lymphaticus. *Clin. Jour.*, June 16, 1915.
- 1979 TODD, F. C. Higher preparation in ophthalmology and oto-laryngology. *Ophthalmology*, Vol. 11, No. 4, July, 1915.
- 1980 TROEMNER. Diagonal gunshot wound through the neck. *Berl. Klin. Woch.*, p. 361, No. 14, 1915.
- 1981 TYKOCINER. Parotitis in the course of typhoid. *Deut. med. Woch.*, p. 238, No. 8, 1915.
- 1982 VAGLIO, R. Insufficiency of the ductless glands common with inherited syphilis. *Pediatrics*, p. 1, Jan., 1915.
- 1983 VANDEN BERGH, A. A. H. Mikulicz's disease. *Nederl. Tijdschr. v. Geneesk.*, Jan. 9, 1915.
- 1984 VINCENT, S. Functions of pituitary body. *Practitioner*, Jan., 1915.
- 1985 VIRDEN, J. E. Are cocain solutions injured by boiling? *Amer. Jour. Surg.*, Vol. 29, No. 8, p. 288, Aug., 1915.
- 1986 VOLKMANN, J. Wounds of the lungs. *Deutsche Zeitschr. f. Chir.*, July, 1915.
- 1987 VON BRUNN. Wounds of the head (Kopfschüsse). *Deut. mediz. Woch.*, Nov. 11, 1915.
- 1988 WASHBURN, J. L. Tumors of the lachrymal gland. Report of a case with operation. *Ohio State Med. Jour.*, Nov. 15, 1915.
- 1989 WATERS, C. A. Roentgenization of the thymus gland. *Jour. A. M. A.*, April 24, 1915.
- 1990 WEIL, R. Treatment of parotid tumors by radium. *Jour. A. M. A.*, p. 2138, Dec. 18, 1915.
- 1991 WEINGAERTNER, M. Roentgenstereoscopy and its application in rhino-laryngology. (Die Röntgenstereoskopie und ihre Anwendung in der Rhino-Laryngologie). *Arch. f. Laryngol. u. Rhinol.*, p. 297, Vol. 29, No. 3, 1915.
- 1992 WEISCHER, DR. Levy-Dorn roentgen localization of foreign bodies (Zur Lokalisation der Fremdkörper nach Levy-Dorn). *Zentralblatt f. Chir.*, Vol. 42, No. 27, July 3, 1915.
- 1993 WERSHOFF, A. E. Carious teeth as a factor in ocular disease. *Jour. Ophthal. and Oto-laryngol.*, p. 141, May, 1915.
- 1994 WESKI, O. Localization of foreign bodies by radioscopy. *Muench. med. Wchnschr.*, Feb. 16, 1915.
- 1995 WHITE, W. C. Silvol. *Jur. Ophthal. and Oto-laryngol.*, p. 380, Dec., 1915.
- 1996 WIDEROE, S. Plastic operation for adamantoma of lower jaw. *Norsk. Mag. for Laegevid.*, Vol. 76, No. 7, July, 1915.
- 1997 WILLIAMS, F. Utility of catarrhal vaccines. *Calif. State Jour. Med.*, Jan. 15, 1915.
- 1998 WILLIAMSON, W. D. Treatment of facial neuralgia with deep injection of alcohol. *Maine Med. Ass'n Jour.*, Dec., 1915.
- 1999 WISHART, D. J. G. Animal extracts in rhinology and laryngology. *Can. Med. Ass'n Jour.*, Aug., 1915.
- 2000 WOOD, C. A. The eyelids and lachrymal apparatus of birds. *Ophthalmology*, July, 1915.
- †2001 WRIGHT, J. D. The value of discipline in the care of the sick child. *THE LARYNGOSCOPE*, March, 1915.

- 2002 WURDEMAN, H. V. Eye symptoms of hypophyseal diseases. *Ophthal.*, April, 1915.
- 2003 WYLIE, A. Bullet wound of face. *Proc. Royal Soc. Med., Laryngol. Section*, Vol. 9, No. 2, Dec., 1915.
- 2004 YALE, A. W. Static electricity in the treatment of the eye, ear, nose and throat. *Jour. Ophthal., Otol. and Laryngol.*, p. 877, Oct., 1915.
- 2005 YEN, F. C. Case of fibroma of the parotid. *Jour. A. M. A.*, p. 1238, April 10, 1915.
- 2006 YORK, C. Recent progress in rhinology, laryngology and otology. *Liverpool Med.-Chir. Jour.*, Jan., 1915.
- 2007 ZESAS, D. G. Injury of vagus nerve in the neck (Klinik und Therapie der Vagusverletzungen am Halse). *Centralbl. f. d. Grenzgeb. der Med. u. Chir.*, No. 6, pp. 507-636, March, 1915.

Books.

- 2008 BASS, C. C. and JOHNS, F. M. Alveodental pyorrhea. W. B. Saunders Company, Philadelphia and London, 1915.
- 2009 BILANCIONI, G. Manuale di oto-rino-laringo-jatria. Il Policlinico, Rome, 1915.
- §2010 BLAU, ALBERT. Die nasal Reflex neurosen und ihre Behandlung. A Marcus und E. Weber's Verlag, Bonn., 1915. THE LARYNGOSCOPE, p. 801, Nov., 1915.
- *2011 BROPHY, TRUMAN, W. Oral surgery. A treatise on the diseases, injuries and malformations of the mouth and associated parts. P. Blakiston's Son & Co., Philadelphia, 1915.
- §2012 BROWNING, WILLIAM. The etiology of stammering and methods for its treatment. Albert T. Huntington, Brooklyn, N. Y., 1915. THE LARYNGOSCOPE, p. 801, Nov., 1915.
- 2013 BRUHN, CHRISTIAN. Die gegenwartigen Behandlungswege der Kieferschussverletzungen. Wiesbaden, 1915.
- §2014 BRUHN, MARTHA E. The Müller-Walle method of lip-reading for the deaf. Thos. P. Nichols & Son Co., Lynn, Mass., 1915. THE LARYNGOSCOPE, p. 800, Nov., 1915.
- 2015 BRUENINGS, W. Translated and edited by Dr. W. G. Howarth. Direct laryngoscopy, bronchoscopy and esophagoscopy. William Wood and Co., New York, 1915.
- 2016 BRUZZONE, CARLO. La suppurazione dell'orecchio nell'infanzia. Unione tipografico-editrice. Torino, 1915.
- *2017 BUXTON, DUDLEY WILMOT. Anesthetics; their uses and administration. P. Blakiston's Son & Company, Philadelphia, 1915.
- §2018 Committee on nurses of the Manhattan eye, ear and throat Hospital, New York. Nursing in diseases of the eye, ear, nose and throat. W. B. Saunders Co., Philadelphia and London, 1915. THE LARYNGOSCOPE, p. 865, Dec., 1915.
- *2019 COOLIDGE, ALGERNON. Diseases of the nose and throat. W. B. Saunders Company, Philadelphia and London, 1915.
- 2020 DENKER and BRUENINGS. Lehrbuch der Krankheiten des Ohres und der Luftwege einschliesslich der Mundkrankheiten. Gustav Fischer, Jena, 1915.

- *2021 GILE, BEN CLARK. The nose, throat and ear; their functions and diseases. A treatise upon the breath-road, food-road and accessory organs. P. Blakiston's Son & Co., Philadelphia, 1915.
- §2022 GLEASON, E. B. A manual of diseases of the nose, throat and ear. W. B. Saunders Co., Philadelphia and London, 1915. THE LARYNGOSCOPE, March, 1916.
- 2033 GRAZIANI, V. Polipo congenito del faringe. Tipografico della casa editrice, Pietròcola, Napoli, 1915.
- 2024 HAJEK, M. Pathologie und therapie der entzündlichen Erkrankungen der Nebenhöhlen der Nase. Franz Deuticke, Leipzig und Wien, 1915.
- §2025 JACKSON, CHEVALIER. Peroral endoscopy and laryngeal surgery. THE LARYNGOSCOPE, St. Louis, 1915. THE LARYNGOSCOPE, March, 1916.
- §2026 JOHNSTON, R. H. The straight methods of direct laryngoscopy, bronchoscopy and esophagoscopy. *American Jour. of Surgery*, New York, 1915. THE LARYNGOSCOPE, p. 801, Nov., 1915.
- *2027 KELSON, WILLIAM H. Diseases of throat, nose and ear. Oxford University Press, New York, 1915.
- 2028 KELYNACK, T. N. Defective children. Bale, Sons and Danielson, London, 1915.
- §2029 KNOX, ROBERT. Radiography, x-ray therapeutics and radium therapy. O. and C. Black, Ltd., London; The Macmillan Company, New York, 1915. THE LARYNGOSCOPE, March, 1916.
- 2030 KOLMER, JOHN, A. A practical text-book of infection, immunity and specific therapy, with special reference to immunologic technique. W. B. Saunders, Co., Philadelphia and London, 1915.
- 2031 KYLE, D. B. Text-book of diseases of the nose and throat. W. B. Saunders Co., Philadelphia and London, 1915.
- 2032 LONGO, FRANCESCO PAOLO. Complicazioni intracraniche delle suppurazioni dell'epitympano. T. Manuzio, Roma, 1915.
- 2033 LORD, FREDERICK, T. Diseases of the bronchi, lungs and pleura. Lea and Febiger, Philadelphia and New York, 1915.
- 2034 PARK, DAVIS & Co. Collected papers from the research laboratory. Detroit, 1915.
- *2035 RAMSAY, A. M., GRANT, J. D., WHALE, H. L. and WEST, C. E. Injuries of the eyes, nose, throat and ears. Oxford University Press, New York, 1915.
- §2036 SCHAEFER, KARL L. Einführung in die Musikwissenschaft auf physikalischer, physiologischer und psychologischer Grundlage. Breitkopf und Härtel, Leipzig, 1915. THE LARYNGOSCOPE, p. 865, Dec., 1915.
- 2037 SCHEIBLE, A. F. Bezold's Sektionsberichte über 73 Fälle von Mittelohreiterung. Curt Kabitzsch, Würzburg, 1915.
- 2038 SHEFFIELD, HERMAN, B. The backward baby. A treatise on idiocy and allied mental deficiencies in infancy and early childhood. Rebman Company, New York, 1915.
- 2039 TERMAN, L. M. The hygiene of the school child. Houghton, Mifflin & Co., Boston, New York and Chicago, 1915.

- 2040 Transactions of the American Otological Society. Forty-eighth annual meeting held at Niagara Falls, Ont., Can., June 3 and 4, 1915. Vol. 13, part 3. Published by the society.
- 2041 Transactions of the thirty-seventh annual meeting of the American Laryngological Association, held at Niagara Falls, Canada, June 1, 2 and 3, 1915.
- 2042 Transactions of the section on laryngology, otology and rhinology of the American Medical Association, 1915.
- 2043 WALLIN, J. E. W. The mental health of the school child. Yale University Press, New Haven, 1915.
- 2044 WOOD, ANDREWS and BALLENGER. Eye, ear, nose and throat. Practical medicine series. Year Book Publishers, Chicago, 1915.
- *2045 WORTH, CLAUD. Squint; its causes, pathology and treatment. Bale, Sons and Danielson, Ltd., London, 1915.
- *2046 WRIGHT, J. D. What the mother of a deaf child ought to know. Frederick A. Stokes Co., New York, 1915.

DIGEST OF OTO-LARYNGOLOGY.

12

Normal Nasal Septum and the Pathology of Deflections. F. O. LEWIS,
New York Med. Jour., April 10, 1915.

There are three forms of deflections that the author calls attention to because of their difficulty of correction, and their local and constitutional symptoms: (1) deviation of the quadrilateral cartilage resulting from trauma in such a manner that the convexity of the deformity obstructs one side of the nose, and the free edge, by projecting into the opposite nostril, partially or completely obstructs the other side (with compensatory hypertrophy of the inferior turbinate); (2) deflection of both of the cartilaginous and bony septum extending from within the vestibule almost to the posterior nares, impinging against the inferior turbinate, interfering with drainage and ventilation and associated with Eustachian catarrh and middle-ear deafness; (3) deflection of the vertical plate of the ethmoid, impinging against the middle turbinate giving rise to reflex symptoms (headache, neuralgic pains, cough, asthma). P. F.

13

Transplantation of Cartilage in the Correction of Septal Deformities.
F. O. LEWIS, *Ann. Otol., Rhinol., and Laryngol.*, Sept., 1915.

The indications for cartilage transplantation are: (1) inevitable perforation during the submucous operation; (2) small septal perforations regardless of the cause; (3) deviation with displacement of the cartilaginous septum requiring complete removal of the free end of the quadrilateral cartilage.

To avoid the objectionable symptoms following perforation the following method is recommendable by the author: In performing a submucous resection a portion of the quadrilateral cartilage which has been found necessary to remove is taken out in as large a piece as possible. This is washed and placed in sterile salt solution and after the completion of the operation if it is found that both sides of the mucous membrane have been penetrated or injured by removing pieces of bone, a piece of the cartilage is trimmed by means of scissors to the proper size and introduced between the two surfaces of mucous membrane. Both sides of the nostril are then packed tightly with gauze, being careful to see that the cartilage is in its proper position and the mucous membrane not curled upon itself. In about twelve hours the packing is carefully removed. The patient should be seen daily for four or five days in order that the position of the cartilage may not be altered and that the surfaces of the mucous membrane are kept clean because if infection occurs the cartilage is rapidly destroyed. Therefore, if infection is suspected free drainage is necessary and if instituted promptly the life of the cartilage is not impaired. The patient should not blow his nose but should snuff his secretions through his mouth. No washes or douches need be prescribed.

The method has been used by the author in twelve cases, all of which have entirely healed with no signs of perforation remaining. P. F.

24

The Inferior Turbinate. J. O. CAVANAUGH, *Ann. of Otol., Rhinol. and Laryngol.*, Sept., 1915.

The author details the anatomy, physiology and histology of the inferior turbinates, mentioning some original observations and measurements, and states that these bodies are deserving of more recognition because they mean more to the human economy than most rhinologists would have us believe. In the author's opinion there are three types of turbinates which attract attention, excluding the malignant and non-malignant tumor types. These are (1) the intumescent, (2) hypertrophy of the mucous membrane, and (3) hypertrophy of the bony part, the mucous membrane being practically normal. It is only exceptionally that an inferior turbinate should be totally removed and the author does not agree with those who advocate its removal and preserving the septum. A turbinate, even though it may look pathologic, will always have some function and should be left untouched. The septum should be the point of attack if by so doing we can preserve a turbinate and accomplish our purpose. Where operation on the turbinate is indicated, the author recommends an operation similar to those of Freer and Yankauer. If the entire membrane is at fault, whether it be intumescence or hypertrophy, cocaine and adrenalin (10 per cent cocaine; 1:1000 adrenalin) and make an incision from behind forward at the lower and internal surface down to the bone, then a corresponding incision on the lower and outer side. With a Freer sharp elevator elevate the mucous membrane from the bone on the outer and inner side for one-eighth of an inch from the anterior surface all along the long axis of the turbinate; then, with a Myles alligator forceps, bite out a part of the bone with the triangular mucous membrane area. Now draw the cut edges together, but there should be no tension on the sutures, and suture with silk. Place a piece of cotton or gauze on the under and inner side to aid the tissues in remaining in opposition. Remove in twenty-four hours.

P. F.

29

Inferior Turbinate Operations and the Rational Cure of Hypertrophic Rhinitis. GIUSEPPE LEALE, *Boll. d. Mat. dell'Orecchio*, V. 33, No. 9, Sept., 1915.

The author is of the opinion that in all cases in which the hypertrophic rhinitis is of moderate grade and of the vasomotor type simple cauterization of the mucosa is effective; but where the hypertrophic rhinitis is advanced and gives rise to symptoms of obstruction to breathing operative interference is indicated.

P. F.

41

Adenoids in General Practice. DR. J. D. PAGE, *Bulletin Medical*, Nov., 1915.

The writer aims at indicating to the general practitioner the points which should guide him in deciding when the proper moment has arrived for the intervention of the specialist. He notes the large number of cases demonstrated at the anti-tuberculosis exposition, where

adenoid vegetations were present, and where the nose had become for the moment, a useless ornament, and the necessity which this observation imposes for the regular practitioner to school children. The pathology, symptoms, and diagnosis of adenoid vegetations are briefly sketched.

Under the head of treatment, the writer points out that many fewer cases would come to operation if the general practitioner would make an early diagnosis, and proceed, by establishing better hygiene, and a maximum of nutrition combined with the administration of syrup of the iodide of iron and codliver oil.

In addition, an endeavor should be made to keep the nose free by warm vapors and, if necessary, by irrigations. Again, the habit of nose breathing should be stimulated by training the child to keep the mouth shut, and, if necessary, by applying a bandage around the chin during the night. Operation should be advised when it has become impossible to secure nasal breathing, or when a complication such as acute otitis has supervened.

WISZART.

49

Pollen Therapy in Hay Fever. J. L. GOODALE, *Ann. of Otol., Rhinol. and Laryngol.*, June, 1915.

In the first period 13 patients gave positive reactions to one or more of the following plants: coltsfoot, dandelion, white maple, willow, alder, birch, hawthorne, apple, lilac, oak, tulip, lily of the valley. In the second period (36 patients) the symptoms were ushered in with the flowering of the grasses. Disturbances from many garden flowers are of minor importance. The third period, Goodale has shown to be due in a great measure to the mid-season compositure, such as field daisy, hawkweed, yarrow, etc. The fourth period begins with the general flowering of ragweed, goldenrod, aster, etc., and lasts until frost.

The dosage is determined as follows: The special exciting pollen is first determined by the skin test. A second series of scratches are made at a distance from the first and different dilutions of the pollen extract in question are applied. It is important not to have this test applied near the skin reddened by the first tests because of the increased excitability of this region.

The dilutions may be made by adding some of the stock solution to alcohol of the same strength and a 25 per cent, 10 per cent, 1 per cent, or weaker dilution of the original extract are applied to a second series of scratches. The initial dose is determined by the dilution which does not excite a definite skin reaction. Not more than five or ten drops should be injected. The injections should be made at intervals of two days to a week, gradually increasing the dose a few drops at a time, and in the strength of the percentage.

P. F.

53

Hay Fever; Its Treatment by Injections of a Solution of Ragweed Pollen. E. T. MANNING, *Jour. A. M. A.*, Feb. 20, 1915.

The definition that the author gives of hay fever, in accordance with our present understanding, is that it is an exudative catarrh of the

conjunctival, nasal and tracheobronchial mucous membranes, as the result, in sensitive individuals, of the sensitizing anaphylactic action of the pollen of certain plants. The best explanation of the action of the pollen in hay fever is, in the author's opinion, that of Koessler who attributes the disorder to conditions of the mucous membrane which interfere with its normal digestive capacity of the foreign protein which, therefore, enters the system and sensitizes the local tissue. Manning states that there are two ways in which to treat the disease on the basis of this theory. One is to combat the disease by adding some substance to the organism capable of neutralizing the poisonous radical of the protein (passive immunization). The other is to develop within the organism a substance which will neutralize the action of the poisonous radical (active immunization). Manning employed, in his experiments, a modified Koessler technic. Ten milligrams of pollen from each variety is triturated with sterile silicon in an agate mortar. Twenty cubic centimeters of a sterile salt solution ten times stronger than a physiologic is then gradually added and thoroughly shaken. The suspension is placed in the incubator for twenty-four hours and again shaken. The mixture is then centrifugalized and the supernatant fluid pipetted off. This solution is a dilution of 1:1000 and all others are prepared from it.

The unit of pollen toxin used was the amount of portein contained in 1/100000 grams of pollen.

The experiments were made on twenty-one cases. Fourteen of them were objectively and subjectively relieved. In the other seven the treatment was incomplete, but four of these reported that their attacks were considerably lighter.

The immunity conferred is, however, of brief duration and while it is difficult to say what constitutes a cure, the impression that the author has obtained from his work is that the distressing symptoms were decidedly ameliorated.

P. F.

56

Pollinosis (Hay Fever); Consideration of Its Treatment by Active Immunization. S. OFFENHEIM and M. J. GOTTLIEB, *New York State Jour. Medicine*, May, 1915.

The antigen which the authors have developed for curative purposes was prepared as follows: The pollen was ground up for several days with sand and a sufficient amount of 5 per cent sodium chlorid solution and 0.5 per cent phenol added to prevent bacterial growth. The mixture was placed in the incubator for 72 hours and then filtered by suction. The filtered extract was then precipitated with absolute alcohol (eight parts) and filtered (in a Buchner filter). The precipitate was dried and weighed. This precipitate did not give the biuret or ninhydrin reactions. It is partly soluble in 0.85 per cent NaCl solution and physiologically active in very weak solutions. The dried precipitate was dissolved in 0.85 per cent NaCl solution, 0.25 per cent phenol added, and serial dilutions made.

Of eleven patients treated five were cured for the season. In four there was marked improvement. Of the two patients who were not at all benefited one had polypoid degeneration of the middle turbinate

with bone necrosis and the other patient (a physician) reacted both to ragweed and goldenrod pollen. He received alternating injections of the ragweed and goldenrod extracts but he could not develop a tolerance.

The authors recommend that treatment had best be undertaken about ten weeks before the first attack is expected to occur and regularity of attendance at weekly intervals is important. Beginning with from one to five units of pollen extract the dose is gradually increased until a local reaction appears at the site of the injection. This dose is continued until the patient no longer shows any reaction. Then the dose is gradually increased again. A unit of pollen extract is the amount of antigen dissolved in one cubic centimeter of extract at a dilution of from 1 to 20 millions.

P. F.

73

Changes in the Pituitary in a Case of Lympho-sarcoma of the Nasopharynx.

G. BASILI, *Zeitschr. f. Laryngol.*, Bd. 7, H. 6, Sept., 1915.

The lympho-sarcoma was of the small, round cell variety. The patient was a man, 36 years old, who presented the psychic syndrome described by Citelli. At autopsy the pituitary was found greatly enlarged. Microscopic examination of the pituitary showed a marked increase in size and number in all the cells. Capillaries were dilated and filled with blood and celloid substance. Treating the preparation by Ciaccio's method and staining it with Sudan III, a large quantity of cellular liquid could be demonstrated in the pituitary. Both the increase in eosinophil cells and cellular lipid, Basil regards as an expression of hyperfunction. The findings are identical with those obtained by Citelli in studying the effect of adenoids on the pituitary gland.

P. F.

99

The Treatment of Scleroma of the Upper Air Tract with Auto-vaccines.

J. BRUNNER and C. JAKUBOWSKI, *Arch. f. Laryngol.*, Bd. 29, H. 2, 1915.

The authors give a short, historical account of the advancements regarding the question of scleroma of the upper air passages and their own observations on the anatomical pathology of the disease. The auto-vaccine prepared by the authors was injected subcutaneously. No serious general reactions were noted. Very good results were obtained in a number of cases by means of the vaccine, provided the disease was not of too long standing. The disease was arrested and the infiltrations either disappeared altogether or became greatly reduced in size. The author urges that the disease be regarded as an infectious disease and as such it should be under state regulation.

P. F.

137

The Etiology and Treatment of Ozena. HENRY HORN, *Jour. A. M. A.*, Aug. 28, 1915.

Horn reports six cases treated with the Perez vaccine and the results obtained tend to show, he believes, although they do not prove, the great value of the remedy in the treatment of ozena. He describes the method of preparing the vaccine and he states, also, that several of

the stock vaccines on the market have been used with advantage. These vaccines usually contained the Friedländer bacillus. The micrococcus catarrhalis, the pneumococcus and streptococcus, and this may raise the question whether there are not types of ozena closely resembling that due to the Perez bacillus, but caused by other organisms, or whether ozena is not a mixed infection. His conclusions are: (1) The coccobacillus foetidus ozenae, Perez, as isolated by Hofer, has fulfilled all the bacteriologic requirements necessary to establish its identity as the etiologic factor in ozena; (2) the isolation of this organism is attended with considerable difficulty; (3) the production of agglutinating serum in rabbits is an exceedingly difficult task; (4) the preparation of autogenous vaccines in every case is very difficult, if not impossible; (5) at present mixed vaccines made from various strains of Perez bacillus is the most practical method of treatment now available; (6) it may be necessary to precede or combine with the treatment the vaccines made from the organisms which are usually present in combination with the Perez bacillus; (7) it is possible that there may be two or more types of ozena bacteriologically different but clinically identical.

P. F.

139

Complement Fixation in Acute Rhinitis. K. HOWELL, *Jour. Infect. Dis.*, May, 1915.

Using the bacillus rhinitis as antigen, Howell obtained complement fixation with the serums of persons affected with rhinitis and of persons injected with the bacillus after it has been killed by heat. The period of fixation is short but it is most marked a few days after the onset of the infection. Control tests with serums of normal persons and of patients with other infections were negative with the bacillus rhinitis. Further proof of the specific relationship that the bacillus bears to acute rhinitis is shown by the fact that complement fixation is obtained when the serum is tested with suspensions of other bacteria presumed to be closely associated with rhinitis if not the actual cause of the condition.

P. F.

143

Total Rhinoplasty; Report of a Case. R. H. JOHNSTON, *Amer. Jour. Surg.*, April, 1915.

The patient had his nose entirely severed by being drawn up against a revolving saw. The lower lip was cut off to the lower border of the maxilla and the upper lip was also cut through. The operation performed was as follows: A piece of cartilage from the left eighth rib was removed and slipped beneath the periosteum according to Carter's method, a little above the center of the left forehead. Three months later the skin on the two sides of the remains of the nose was dissected up and the flaps, skin surface down, were turned into the facial opening and sutured in the middle line. The flap for the formation of the nose began at the inner end of the eyebrow and continued up to the hair line and across the forehead to the end of the transplanted cartilage. From above the left eyebrow to the lobe of the nose

it was bent downward and inward. The next step in the operation consisted in dissecting away the skin from the periosteum up to the cartilage which was also removed from the bone with its strip of periosteum. The flap was then turned down raw surface below, and the upper end of the cartilage was stitched to hold it firm. The lower end was sutured into an incision at the upper lip so that the nose would project from the face. P. F.

154

The Early Recognition of Cancer of the Upper Air Passages.

EMIL

MAYER, *Amer. Jour. Surg.*, July, 1915.

The author points out the necessity of viewing with suspicion the so-called innocent intranasal growths. These have a tendency to undergo malignant metamorphosis. In the nose, cancer often springs from one of the turbinates; but in most instances the neoplasm is deep-seated with its origin in one of the sinuses, especially the antrum of Highmore. In the mouth, the most frequent site of malignant growth is on the tongue and as leucoplakia buccalis is often a forerunner of malignancy any induration of the tongue should be viewed with suspicion. The author calls attention to the advantages of the naso-pharyngoscope in the determination of the location and size of post-nasal new growths. Cancer of the larynx has its favorite site on one vocal or false cord. In the early stages it is limited to one-half of the larynx and its early recognition is imperative because its complete removal is possible and the results are gratifying. In a case of cancer of the epiglottis, mentioned by Mayer, there has been no discomfort following removal of the epiglottis and no recurrence in two and one-half years. P. F.

161

Rhinophyma; Its Etiology, Pathology and Treatment. W. MILLIGAN, *Lancet*, Sept. 18, 1915.

The author reports a case successfully treated by dissecting away the hypertrophied tissue followed by skin grafting. He does not believe that alcoholism or traumatism are etiological factors in rhinophyma. Histological examination of the diseased tissue shows marked thickening of the connective tissue of the corium with hyperplasia and cystic dilatation of its sebaceous glands. There also occurs deposit of fat between the various hypertrophied areas. As a result of these changes there is an elevation of the papillary layer of the true skin into projections of various sizes and shapes. A very marked feature of the histologic picture is the cystic dilatation of one or more of the sebaceous glands associated also with marked thickening and tortuosity of the arteries, veins and capillaries. P. F.

172

Headache Associated with Intranasal Disorders. L. HEMINGTON PEGLER, *Lancet*, Feb. 27, 1915.

Headache due to nasal causes may be classified into three groups; (1) simple or non-inflammatory group in which the headache is due to pressure within the nasal cavities by one piece of bone or cartilage

against another, the mucous membrane intervening. For example, a septal spine projecting against a middle or inferior turbinate and concealed by an anterior deflection of the triangular cartilage; (2) chronic inflammatory group in which there is simple degeneration and softening of bone in turbinate disease, polypi or polypoid degeneration with or without associated involvement of the mucosa of the accessory sinuses; (3) inflammatory group, in which the headache is associated with various forms of rhinitis.

P. F.

205

Gonorrheal Infection of the Upper Respiratory Tract in Adults. JOHANNES ZANGE, *Zeitschrift f. Ohrenheilkunde*, Bd. 73, H. 3, Nov., 1915.

A noted surgeon, 40 years old, while operating on a patient for gonorrheal phlegmon was struck in the face by some of the pus. The surgeon merely wiped his face and continued to operate. Three days later he developed a left conjunctivitis and the following day a right conjunctivitis, which yielded to treatment with 10 per cent solution of protargol. Microscopic examination showed a pure culture of gonococci. Three days after the appearance of the conjunctivitis the patient complained of difficulty in swallowing and pain in the pharynx. A swab from the pharynx showed numerous, and many intracellular, gonococci. Nasal examination the same day showed pus, more profuse on the right side than on the left, issuing from the naso-lachrymal duct. Throat examination showed a thin layer of pus covering the fauces and the posterior pharyngeal wall. Abundant gonococci were present in the pus from the nose and in the pharynx. No other organisms could be demonstrated.

In the course of the next six days the symptoms increased in severity, the infection involving the larynx and the Eustachian tube. Shrapnel's membrane and the malleus on the left side were reddened and laryngeal examination showed the whole hypopharynx, the epiglottis and the laryngeal ventricle deeply red and covered in places with pus. The vocal cords were deeply injected and edematous.

In none of the affected parts did erosion or ulceration occur. The infection gradually yielded after about fourteen days to local treatment four or five times a day with 5 per cent solution silver nitrate and 2 per cent protargol solution.

The hoarseness lasted eight days. Otherwise there were no after effects.

P. F.

212

The Late Results of Cleft-palate Operations. T. W. BROPHY, *Surg., Gyn. and Obst.*, Jan., 1915.

The operation for cleft-palate, when performed in early infancy, brings into action the muscles of the palate and the muscles develop instead of becoming atrophied for want of use. If the operation is delayed the muscles cannot as surely be made to subserve the same purpose as the tissues which develop through natural development and the development of the muscles is proportional to that of other tissues. Again, if the operation is performed early there is much less deformity in the bony as well as in the soft tissues.

When the operation is performed in late infancy, when the child is from six to twenty-four months of age the conditions are a little different, because the tuberosities of the maxillary bones still remain widely separated and the posterior part of the cleft consequently will also be widely separated. As the patient grows older the difficulties increase. The arch is broader and remains broader. The wider the cleft the shorter the velum and, therefore, the soft palate does not approach the posterior pharyngeal wall as it normally should. Phonation is not as clear unless the pilato-pharyngeal muscles are in part utilized to lengthen the velum.

P. F.

232

Appendicitis as a Sequela of Tonsillitis. H. B. ANDERSON, *Amer. Jour. of the Med. Sci.*, Vol. CL., No. 4, p. 541, Oct., 1915.

After referring to the importance of the tonsils as portals of entrance for numerous and varied infections, the author credits Kelynack with first directing attention to the occurrence of appendicitis secondary to tonsillitis. A long study of the subsequent references to the subject follows. The author reports one case under his own observation and gives an account of several others from personal communications.

PACKARD.

238

The Proper Position of Tonsillectomy in Pediatrics. SANFORD BLUM, *Archives of Pediatrics*, Nov., 1915.

Thinks tonsillectomy done too frequently and is of more benefit in older than younger children. Discusses conditions it is most often done for.

1. Rheumatism. Not certain as to its exact cause and nature and why should essential organs be removed for theory?
2. Endocarditis. Rare condition in children anyway. "In no single case of endocarditis has it, as far as I am aware, been positively proved as emanating from the tonsil."
3. Chorea. Etiology unknown. Knows of two previously healthy children who subsequently developed chorea after removal of the tonsils.
4. Enlarged cervical glands. The author's experience has seen no benefit from tonsillectomy in these cases. Thinks that the tonsil is an excretory organ for the cervical glands and the tubercle bacilli found in them are not in them as infective agents but rather as waste products from the cervical glands.

In the author's opinion, if a child can be gotten safely over his sixth or seventh year with conservative treatment, he will not probably ever require tonsillectomy. Children over eight years who have their tonsils are heavier and healthier as a rule than those who do not.

The author thinks tonsils should be conserved for the following reasons:

1. Not superfluous if nature put them there.
2. Combat infection.
3. Possible internal secretion.
4. Eliminative organs for waste products of dentition.
5. Eliminative organs for systemic diseases.
6. Modulate the voice.
7. Moisten the food and possibly throw out a digestive ferment.
8. Take up foreign matter from nasal mucous membrane.
9. Author thinks they are excretory organs for cervical glands.

PACKARD.

239

Tonsil, Excretory Organs for Cervical Glands. SANFORD BLUM, *Archives of Pediatrics*, Nov., 1915.

Injected chemicals into cervical glands not found normally in cervical glands or tonsils, and recovered them in the tonsils and in the mouth.
PACKARD.

258

Effects of Tonsil Extract on the Blood Picture. C. B. FARMACHIDIS and A. VATTUONE, *Policlinico*, No. 3, March, 1915.

Tonsil extract increases the number of both the red and white blood corpuscles. It possesses a glycolytic action, and is also able to counteract the fatal toxic action of epinephrin.
P. F.

263

The Tonsils and Cervical Adenitis. H. GARDINER, *Lancet*, Oct. 2, 1915.

The author's investigation was carried out to determine what evidence there was bacteriologically of the infectivity of the tonsils in cervical adenitis, in the absence of any other possible source of infection. Organisms were found in 80 per cent of the cases, as follows: *Micrococcus catarrhalis*, 7 cases; *staphylococcus*, 5 cases; *pneumococcus*, 7 cases; *streptococcus*, 7 cases; *bacillus coli communis*, 2 cases; *bacillus tuberculosis*, 4 cases; *bacillus of Friedländer*, 1 case. The conclusions reached by the author are: (1) In the majority of cases (80 per cent) of chronic cervical adenitis where no obvious source of infection is present the tonsils are infected; (2) the size of the tonsils makes no difference as to their infectivity, except that the small fibrous tonsil is likely to be more dangerous than the large; (3) the number of cases showing the presence of tubercle bacillus is relatively small but is larger than in simple cases of enlarged tonsils; (4) the frequent presence of other organisms suggests that a large proportion of so-called chronic tuberculous glands are in reality chronic septic glands; (5) as the organisms are present in the deepest parts of the glands they are removable only by complete enucleation of the tonsils. P. F.

274

Tonsillar Spots in Measles. CHARLES HERMAN, *American Journal of Diseases of Children*, Oct., 1915.

The author describes tonsillar spots in measles.—The patients must be seen early and the spots must be looked for. A good view of the tonsils is necessary. Spots vary in number from 2 to 30, in size from point to head of pin, in shape from regular or irregular streaks, 2 to 3 mm. in length, to round spots, in color from bluish grey to white. They last from one to three days. They are less valuable for diagnosis than the Koplick spots since they are present in only about 40 per cent of cases and because their appearance is not always characteristic sometimes hard to diagnose from follicular tonsillitis. They may be valuable in making an early diagnosis in an epidemic for purposes of isolation. The spots are frequently present when the patient has no other objective symptoms except rise in temperature.
T. H. HALSTED.

277

Tonsillectomy by Blunt Dissection Under Local Anesthesia. JAMES B.

HORGAN, *Jour. Laryngol., Rhinol. and Otol.*, Dec., 1915.

The following is Dr. Horgan's technique: Fifteen minutes before the operation the patient is given a hypodermic of morphine sulphate 1/4 and atrophine sulphate 1/120. The fauces are first anesthetized superficially by the application of 10 per cent cocain together with a few drops of adrenalin. By means of a sufficiently long syringe up to 15 c.c. of novocain solution are injected around each tonsil (two tablets of novocain to 30 c.c. sterile normal salt solution. After the pillar has been drawn aside by any suitable forceps the tonsil is grasped with a tonsil volsellum forceps and by maintaining firm inward tension the plica will be put on the stretch and little difficulty will be experienced in splitting it with the dissector. The dissector should first be worked in an antero-posterior direction, after which by a strong upward sweep of the instrument the upper pole of the tonsil is cleanly and totally evulsed. The lingual pole of the tonsil may be divided by a suitable scissors or a snare. As soon as it is ascertained that all hemorrhage has ceased a small quantity of equal parts of iodoform and boric acid is insufflated on the raw surface. After the operation the patient should be propped up in bed for six hours. Post-operative treatment consists of a peroxide spray or gargle.

P. F.

289

The Post-operative Antiseptic Treatment of the Tonsillar Fossae. G. P.

MARQUIS, *Jour. A. M. A.*, July 10, 1915.

This consists in applying 50 per cent tincture of iodine to the tonsillar fossae immediately after removal of the tonsils.

P. F.

311

The Relation of the Tonsil to the Thyroid. D. ROMAN, *Jour. Ophthal.*

Otol. and Laryngol., July, 1915.

As distinguished from hypertrophy and hyperplasia there is a form of enlargement of the thyroid resembling simple goitre histologically in which the cellular proliferation is more of a type of inflammatory infiltration. The enlargement is symmetrical. The basis upon which the tonsils may be accused of being an etiological factor in which form of enlargement, which may be induced by direct infection, is the pharyngolinguinal origin of the embryonic thyroid from the fore-gut ventrally in the median line to the bronchial arches and the persistence of the ductus thyroglossus or the foramen caecum. The infection from the tonsil and peritonsillar region can be carried to the thyroid either by the blood stream or the lymphatics. However, no germ, in any of the cases studied by the author, could be isolated from the thyroid. This negative result might be due to one of two causes: (1) either by a chemical destruction of the bacteria by the thyroid secretion, or (2) that no actual bacterial invasion occurred, but only a toxic irritation or chemical stimulation to cellular proliferation. Out of 2,236 cases of thyroid disease the author observed this form of thyroid enlargement in 187 cases and nearly 80 per cent of these were in young people

between two and twenty years of age. Surgical treatment of the tonsils and adenoids produced rapid subsidence and resolution of the condition.

P. F.

319

Primary Syphilis of the Tonsil. C. MORTON SMITH, *Journal of Cutaneous Diseases*, Oct., 1915.

Syphilis is more than a venereal disease as shown by the large number of accidental extragenital infections. Among accidental infections, chancre of the tonsil ranks, with different observers, from first to fifth in genital inoculation.

Chancre of the tonsil is usually unilateral and more frequently on the right side; males more than females. Kissing plays a most important part in primary syphilis of the tonsil. Drinking glasses, eating utensils, pipes, mouth pieces of wind instruments, nursing bottle nipples, cigars, cigarette stubs, tonsillotomes, Eustachian catheters, mouth to mouth insufflation of new born, have been reported as the mode of infection. Literature and Osler in his latest edition give prominence to "improper practices." Such, however, does not appear to be the author's opinion.

Symptoms.—Sore throat with swelling of tonsil followed by stinging pain on swallowing are first symptoms, a hard, usually painless, non-inflammatory enlargement of the gland under the angle of the jaw on the same side appears in from one to three weeks. On examination, tonsil is enlarged, of deep red color, pillars red and swollen, varying sized ulcer or erosion with dirty gray secretion or false membrane is seen on the tonsil. The crypts and follicles are ideal incubators for spirochetæ. Chancre of the tonsil becomes decidedly indurated early in its course. Clinical findings should be verified by dark field and the Wassermann reaction.

Chancre of the tonsil should be considered in all sore throats, especially if unilateral, lasting over two weeks and failing to respond to treatment for simple angina. It must be differentiated from (a) Acute tonsillitis, (b) Peritonsillar abscess, (c) Vincent's angina, (d) Diphtheria, (e) Tuberculous ulcer of tonsil, (f) Malignant disease, (g) Mucous patch (secondary), (h) Gumma of tonsil (tertiary).

In all of these the behavior of the near-by lymph glands is most important in making a correct diagnosis:

In Peritonsillar—Difficulty in opening the mouth, uvula and median raphe are pushed to opposite side, fever, chill, foul breath and if glands are enlarged they are tender. Motion of jaws not limited in chancre. In Vincent's angina smears show short, thick spirochetæ and fusiform bacilli. Diphtheria—Smears and culture verify—glands tender and painful. Malignant disease and tuberculosis—slow development, glands enlarge late. Gumma—No glandular enlargement.

T. H. HALSTED.

337

Spirochetal Ulceration of Tonsils in Soldiers. W. WINGRAVE, *Lancet*, July 24, 1915.

The disease described by Wingrave, probably identical with Vincent's angina, may be divided into two groups: an acute, in which the disease lasts for about seven days, and a subacute, which lasts two or three

weeks. Vincent's spirochete (*spirochete fetida*) is the exciting cause. The condition is characterized by three symptoms: a deep, sloughing ulcer on one tonsil, fetid breath, the presence of spirochetes and fusiform bodies. Treatment consists in applying a 5 per cent solution of trikresol in alcohol to the ulcer accompanied by mouth washes of weaker solutions of the same antiseptic.

P. F.

338

Vincent's Angina; A Review of the Present Position. W. WINGRAVE,
Jour. Laryngol., Rhinol. and Otol., December, 1915.

In practice, two distinct forms of Vincent's angina are met with: One which runs its course in a few days, another which persists for many weeks. Either form may present one or three local types: (1) Small multiple plaques which correspond with the lacunae of both tonsils; (2) solitary, unilateral, circumscribed patches showing slight edema and shallow ulceration; (3) solitary large, deep, irregular ulceration.

With but slight elevation of temperature and great depression, Vincent's angina may at first be mistaken for diphtheria, especially when the patches are bilateral. A culture will differentiate the conditions. Clinically, a patch in Vincent's angina does not show much increase in size and, if bilateral, never extends across the middle line and it remains fairly circumscribed. Vincent's patches are generally single and large, which differentiates them from ordinary lacunar tonsillitis.

The temperature and constitutional disturbances are markedly affected when the condition is associated with a pyogen such as the streptococcus. Glandular enlargement is rare except when there is supplementary pyogenic infection.

Two organisms, a spirochete and a fusiform body are accepted as essential to the diagnosis of Vincent's angina. The spirochete is larger and coarser than the pallida. The fusiform rod is sluggishly motile and appears as a straight or curved rod with tapering ends. At the equator there is an unstrained cleft. It is always thicker at the equator and it may occur singly or in pairs end to end. Both organisms are Gram negative.

The spirochetes are situated superficially, being confined either to the surface of the tonsil or to its crypts. They are not confined to the throat but may occur in the nose, the accessory sinuses, gums, decayed teeth, middle-ear, meningeal and cerebral abscesses, etc. They simply follow the death of tissue rather than precede or actually cause necrosis.

To demonstrate the organisms: (1) Fix the film by heat; (2) wash with acid alcohol; (3) stain with aniline gentian violet; (4) treat with iodine solution; (5) wash with acidulated water. Or, (1) fix film by heat; (2) quickly flood the slide with 5 per cent collargol solution. Drain the slide by standing it upright and then dry by placing it in an incubator for five minutes. Do not place over a flame as fissures will be formed, resembling spirochetes. The spirochetes are seen clear and unstained on a yellowish background.

P. F.

341

Tonsillectomy as a Therapeutic Measure in the Treatment of Chorea and Endocarditis. J. H. YOUNG, *Boston Med. and Surg. Jour.*, Sept. 2, 1915.

In twelve out of twenty-one cases chorea occurred after tonsillectomy, which suggests that removal of the tonsils does not protect against this disease, and the always present possibility of endocarditis. P. F.

360

Paralysis of the Tongue from Traumatic Division of the Hypoglossal Nerves. MORESTIN. *Societe de Chirurgie de Paris*, June 23, 1915.

A soldier was wounded by a bullet which passed from one side to the other in the intrahyoid region with resulting complete paralysis of the tongue. There are no cocatrices and no induration and no disturbance of taste or tactile sensibility of the organ. The tongue, however, takes no part in the act of swallowing solids and the patient has to throw his head back and push the bolus backward with his finger. Speech is interfered with but the patient can be understood fairly well. P. F.

364

Leukoplakia of the Tongue and Its Treatment by Radium. D. L. R. F. SIERRA, *Rev. Espan. de Urologia y Dermatologia*, Aug., 1915.

The author considers the present standpoint of the etiology, symptomatology and histopathology of leukoplakia and reports the case of a thirty-five-year-old man in whom radium affected a complete cure. The radium, in the form of the bromide, was applied in a dose of 0.6 gm. for forty minutes at the first seance and for ten minutes during the subsequent seances. The treatment lasted forty days, after which time, cicatrization of the lesion was complete, the plaque and induration had entirely disappeared and the flexibility of the tongue returned. P. F.

369

Pathogenesis, Treatment and Prophylaxis of Mercurial Stomatitis. J. ALMEVIST, *Hygiea*, Vol. 77, No. 16, 1915.

The author has carried out extensive investigations on the origin of mercurial stomatitis and is of the opinion that the *modus operandi* of the causation of this condition is as follows: Local processes of decomposition within the mouth set free hydrogen sulphid; the hydrogen sulphid is absorbed by the blood in the superficial capillaries where it combines with mercury in the blood forming mercurous sulphid; the mercurous sulphid is deposited in the endothelium of the capillaries in the form of a granular deposit; as a result of this deposit the epidermis becomes degenerated, thus making a suitable medium for the bacteria present in the mouth, especially the *bacillus fusiformis* and the *spirochete dentium*, which bring about ulceration of the tissue. The sites of predilection are primarily the gums, the angle behind the last upper molar tooth and the tonsils. Secondarily affected are the tongue, the mucous membrane of the mouth and lips. Extension of the process downwards leads to loosening of the teeth from periostitis.

Treatment consists in the use of bichlorid and cyanid of mercury as antiseptics and of potassium permanganate and chlorate and hydrogen peroxid as oxidizing agents. P. F.

371

Leucoplakia; Its Pathogenesis and Treatment with Salvarsan. T. BAER, *Dermatologische Zeitschrift*, March, 1915.

This paper gives a very complete review of the literature of leucoplakia and a discussion of its causation, the author leaning more towards the syphilitic etiology of the disease. He reports in detail a case in which salvarsan proved of marked benefit.

P. F.

386

Oral Tuberculosis. T. E. CARMODY, *Ann. of Otol., Rhinol. and Laryngol.*, June, 1915.

Carmody's investigations show that the parts are affected in the following order of frequency: (1) Tip of tongue, (2) border of tongue and floor of mouth, (3) soft palate, anterior pillar and uvula, (4) dorsum and base of tongue, (5) lower jaw and gums, (6) upper jaw and gums, (7) lips, (8) hard palate, (9) salivary glands.

Ulcers may appear as fissures which, in the early stages, may be entirely overlooked, resulting from infection taking place either from the sputum or through the blood or lymph stream. These fissures may extend into the muscular substance of the tongue and yet show no surface extension except at the corner of the mouth. Ulcers appearing upon the lips, cheeks, surface, edge or tip of the tongue, usually exhibit a reddish base covered with granulation tissue upon which small whitish tubercles seem to be sown. The edges are irregular and may show some induration. A sharp-cut appearance of the edge is not usually found.

Ulcers of the gums begin at the gingival margin, usually around a decayed tooth or one having an artificial crown.

The soft palate is involved often when no other oral structure is affected. This occurs mainly by extension from pharyngeal and tonsillar tuberculosis and usually late in the disease.

The author suggests that there may be a difference in the mode of infection between lupus and tuberculosis and possibly a difference in the organism not shown in the tissues or in cultural characteristics by present laboratory methods. Lupus has little effect on the general nutrition and as a rule causes little pain.

Primary tuberculosis of the mouth must be differentiated from syphilis. Here the Wassermann reaction is of service and also the v. Pirquet test. The appearance of a chancre is that of a punched out ulcer with indurated base and showing no tubercles.

Epithelioma is differentiated from tuberculosis by induration of the edges as compared with the generally soft edges of the tuberculous ulcer. There may, however, be tuberculosis engrafted from epithelioma. Microscopic examination is the only absolute and reliable test.

P. F.

396

Vincent's Angina. A. G. DE SANCTIS, *N. Y. Med. Jour.*, Nov. 6, 1915.

The author reports the case of a little girl ten years old who developed Vincent's angina after scarlet fever. The tonsil became covered with a

greenish white membrane which, when stripped off, left a raw, bleeding surface. Diphtheria was diagnosed and 10,000 units of antitoxin administered. This produced no change and smears made directly from the lesion showed the presence of the characteristic bacilli and spirocheta. Local application of tincture of iodine failed to improve the condition. Accordingly, 0.6 gm. neosalvarsan was given intravenously and from this time on the lesions became smaller and smaller until they finally disappeared.

Another case, in a man 42 years old, cleared up under potassium chlorate, three grains every four hours, in conjunction with mouth washes of the same drug. Diagnosis was corroborated by microscopic examination.

P. F.

420

A Study of 250 Stained Blood Films in Pyorrhea Alveolaris. F. HECKER, *American Jour. Med. Sciences*, June, 1915.

Wright's stain was used in staining the films. Differential count showed a diminution in the small lymphocytes and polymorphonuclear neutrophils and an increase in the large lymphocytes and the irritation forms of Ehrlich. The large lymphocytes showed wide variation in the intensity of nuclear staining and the nuclei showed granules of various sizes and shapes not subject, however, to any definite arrangement. The cytoplasm also showed variations in intensity of staining. The same variations are true also for the polynuclears.

P. F.

453

Vincent's Angina. F. MASSEI, *Arch. ital. di Laringol.*, Oct. 5, 1915.

Massei takes up the consideration of the various remedies suggested for the treatment of Vincent's angina and especially the use of salvarsan and neosalvarsan, locally, as suggested by Rolleston. He has had occasion to treat three cases by this method, the salvarsan being applied once a day for three days. One of the three patients was suffering from active syphilis. Improvement was marked in all three cases. The beneficial action of salvarsan on the disease is probably due to its being caused by the various spirochetes, including the pallida, and the spirilli, that are found in the mouth.

P. F.

472

Unusual Course and Complications of Vincent's Angina. F. REICHE, *Munchener med. Woch.*, Feb. 16, 1915.

The author reports a case in which the constitutional symptoms were very severe, headache being the most prominent symptom. There was an enlarged spleen; unilateral paralysis of the abducens nerve and blood examination showed a leucopenia. The blood also showed an eosinophilia of over 16 per cent which was decidedly against typhoid leucopenia. There was considerable ulceration of the mouth. Wassermann negative (blood and spinal fluid). No diphtheria bacilli could be demonstrated in the mouth, but the Plaut-Vincent organism was present in large numbers.

P. F.

488

Fifteen Cases of Cancer of the Mouth Successfully Treated with Radium.

A STICKER, *Berliner klin. Woch.*, Vol. 52, p. 1040, 1915.

Of the fifteen cases, three were cancer of the tongue, seven of the lower jaw and five of the upper jaw. The radium was applied for twelve hours each night for a total of from three up to as many as fourteen nights. In cancer of the mouth where disfigurement follows radical surgical treatment Sticker believes that radium treatment is superior and is to be preferred. His views are quite radical in this respect and would hardly meet with general approbation as he believes that in cancer, general surgery tends to favor recurrence. Radium treatment, he holds, tends to produce immunity, the destroyed cancer cells being taken up in the body fluids and producing antibodies which immunize the body against cancer.

P. F.

506

Frontal Sinus Suppuration with Results of a New Operative Procedure.

H. A. LOTHROP, *Jour. A. M. A.*, July 10, 1915.

The patient is etherized and placed in a semi-supine position. A single curved one-inch incision is made in the inner portion of the eyebrow, limited externally by the supra-orbital arch in order to avoid the nerve. The bone over the sinus is bared of periosteum and the sinus entered by means of a chisel. It is then enlarged with a rongeur forceps to make an oval opening. Any pus, polypi or granulations are removed. A probe is passed through the ostium into the nose to serve as a guide. The walls on the floor of the sinus are broken down with small curved curettes, the posterior angle of the sinus being avoided. By means of the author's rasps and the burr drills devised by Tilley and Ballinger the interfrontal septum should be perforated and burred away for the purpose of exploring the other sinus. The perpendicular plate of the ethmoid should also be removed by the burr and the agger-nasi cells and other neighboring ethmoid cells broken up with a large burr. The skin incision is closed without drain and all tampons removed.

P. F.

518

Tuberculosis of the Frontal Sinus; Report of Two Cases. J. B. THOMAS,

Jour. A. M. A., July 24, 1915.

In the first case the outer table became perforated and the patient recovered after the operation. The patient, a female, had attended her mother for a number of years of chronic pulmonary tuberculosis and had herself suffered from Pott's disease which had left her somewhat deformed. In the second case the frontal sinus tuberculosis was complicated by osteomyelitis, epidural, subdural and cerebral abscesses. Thomas believes that the high position of the sinus and its better drainage, the bactericidal action of the mucousa, cilia and tears serve to protect the frontal sinus from infection.

P. F.

519

The Pernal Operation for Frontal Sinus Suppuration. P. WATSON.

WILLIAMS, *Lancet*, p. 362, and *Bristol Med.-Chir. Jour.*, p. 24, 1915.

The technic of the operation is the following: The anterior margin of the middle turbinate is engaged with a small angular ethmoidal forceps at its point of attachment to the outer nasal wall and cut through. Keeping to the outer side of the vertical plate of the ethmoid all the agger cells and anteconchal cells are clipped away right up to the crista nasalis. The anterior ethmoidal cells including the bulla ethmoidalis are removed by forceps and the thicker projecting partitions of the cells are laid open and punched away. The size of the sinus is gauged by means of bougies. If a No. 17 bougie enters the sinus the bony boss can be burred away with a 4 mm. wide burr until it enters the sinus. The operation ought to be controlled by a skiagram and if the frontal sinus opening lies well to the outer side and tends to guide entering probes towards the orbital roof it is well to draw the sliding forceps or burr towards the front so as to enlarge the frontal ostium to the front and inward rather than toward the orbital roof outward. With the small forceps any projecting ethmoidal cells may be clipped away. After treatment consists of lavage of the sinus with mild antiseptics or colloidal silver preparations, the passage of the largest bougie that the canal will take in order to prevent adhesions and keep the passage free. In streptococcal infections the author recommends giving about 30 c.c. of polyvalent antistreptococcal serum before operation, and employing sensitized vaccines after operation. P. F.

527

Pathology and Therapy of the Spheno-ethmoid Recess. P. J. MINK,

Arch. f. Laryngol. und Rhinol., Bd. 29, H. 2, 1915.

Mink states that heretofore the spheno-palatine recess has been reached surgically by first removing the posterior end of the middle turbinate. This recess bears the same relation to the posterior sinuses as the hiatus bears to the anterior sinuses. Pathological conditions in this region are of importance owing to the close relationship between the mucous membrane and the spheno-palatine ganglion. The spheno-palatine recess should be considered a part of the respiratory tract; the air stream at first follows the passage between the middle turbinate and the septum and rises higher according to the depth of the inspiration. Although the location of the recess forms a relative protection against any harmful influence of the air, the upper posterior part of the fossa, through the suction exerted at the close of inspiration, has a tendency to draw secretions and foreign bodies into the sinuses. In acute inflammations of the nasal mucosa the posterior upper part of the nasal fossa is shut off from the rest of the nose by the swelling of the rima olfactoria, so that the secretions have to drain through the spheno-ethmoidal recess. Treatment of inflammatory conditions of the upper posterior part of the nasal fossa requires the application of vapors by means of a thin catheter which can be pushed up into the swollen rima olfactoria. A vapor of a saturated solution of menthol in oil of turpentine is applied through the catheter. Adrenalin should be ap-

plied to the rima olfactoria in order to keep it open; for this purpose Mink has devised a thin double-tubed apparatus. In chronic catarrhal inflammation the nasal tissues should be shrunk with 10 per cent cocain followed by silver and iodine applications (Mink describes a thin 1 mm. silver wire to facilitate these applications). P. F.

552

A Modification of Skillern's Preturbinal Operation on the Maxillary Sinus. J. C. G. MACNAB, *Jour. Laryngol., Rhinol. and Otol.*, p. 333, Sept., 1915.

The disadvantages encountered by Macnab with the Skillern operation: (1) The dressings were always painful; (2) the opening had a great tendency to close up; (3) it was impossible to know when to leave out the packing.

The modification of the operation by Macnab makes it partake more or less of a radical operation. Having completed the operation as described by Skillern, the antrum is filled with the French embalming lotion (iodoform, menthol, eucalyptol, balsam Peru, ether) which is allowed to remain five minutes. A small curette is now employed to freshen the mucous lining of the floor of the antrum for about a centimeter adjacent to the nasal floor. The ridge of bone having been carefully polished down, with a scissors the lining of mucous membrane of the outer nasal wall is cut through close to the insertion of the inferior concha and folded neatly on the antral floor, being kept in place by packing the antrum with one-inch bismuth gauze which has been previously boiled in parolein. This remains forty-eight hours. Should the mucous flap show any tendency to curl up, the antrum is repacked.

The patient now has a common antral and nasal cavity all under the inferior turbinate as no part of that bone has been sacrificed. The cavity can be washed out without any discomfort.

The patient need be seen about once a week. All the cases operated on by this modification were well within five weeks. P. F.

554

Intranasal Antrum Operations, with Report of Ninety Cases. H. McNAUGHT, *Jour. A. M. A.*, Sept. 4, 1915.

From an experience covering ninety cases, McNaught does not believe that antrum infections arise most often from the teeth. In his series very few cases were of this etiology. The author prefers the Krause operation modified by himself with the use of the Reeves punch which obtains better drainage and facilitates intranasal applications.

P. F.

568

The Importance of the Paranasal Sinuses in the Explanation of Pain in the Face, Head, Neck and Shoulders. M. A. BLISS, *American Journal of the Medical Sciences*, February, 1915.

It is the author's view that inflammation of the nasal sinuses or in the nasal cavity proper may involve the sphenopalatine ganglion and give rise to pain at the root of the nose, around the eyes, in the jaws and teeth extending to the zygoma, ear, mastoid, occiput, neck, scapula,

breast, and in severe cases, to the arm, forearm, hand and fingers. Associated with the pain may be disturbances of taste and diminished sensibility of the soft palate, pharynx, tonsils and nasal mucous membrane. The author has recently met with cases exhibiting all the features mentioned.

P. F.

577

Diagnosis and Treatment of Suppuration of the Nasal Accessory Sinuses.

F. DIEBOLD, *Correspond. Blatt für Schweizer Aerzte*, Aug. 14, 1915.

For locating the source of pus, Diebold uses hexaethyl violet (hexaethyl rosanillin chlorid) as the dye best adapted for this purpose. Furthermore, the stain may be used as a therapeutic agent because it softens up the thick secretion in the sinus and the fetid odor is notably reduced even by one application.

Its employment is as follows: a small piece of the substance (it comes in solid form) is melted on the end of a probe and introduced into the cavity to be examined. A single grain will stain all the secretions accumulating there. The tint, however, deepens gradually and becomes most intense from the third to the sixth day, but traces of it may even be found as long as six weeks after the application. The thicker the secretion the longer it takes for the crystals to dissolve and as the stain is not readily soluble in salt solution or blood, there is little danger of any toxic effect from it.

P. F.

587

Three Cases Showing the Shoulder-Arm-Hand Syndrome of Paranasal Disease Cured by Operation. J. C. G. MACNAB, *Jour. Laryngol., Rhinol. and Otol.*, Sept., 1915.

The three patients were males between twenty and forty years of age. All showed sufficient nasal abnormality to warrant an exploration of the posterior ethmoids and sphenoids. Examination showed that the principal pathologic condition was enlarged cystic cells easily broken down and lined with mushy mucous membrane. As complete an exenteration as possible of the posterior cells was carried out and the entire anterior wall of the sphenoid was removed. The symptoms rapidly disappeared, followed by complete cure.

P. F.

601-

Sudden Blindness Due to Suppuration of the Accessory Nasal Sinuses, with Report of Three Cases. H. H. STARK, *Journal of the American Medical Association*, October 30, 1915.

The author has collected some 88 cases from literature, and the three of his own. In two of his cases, the patient became suddenly blind, the other one gradually. Ophthalmoscopic examination practically negative. The cases regained sight after removal of anterior portion middle turbinate and draining the ethmoid cells. Draws attention to the importance of recognizing this condition.

PACKARD.

605

The Treatment of the Various Forms of Inflammation of the Mucous Membranes of the Nasal Accessory Sinuses. W. UFFENORDE, *Arch. f. Ohrenheilkunde*, May, 1915.

The author considers the diseases collectively from the histological, anatomical, clinical and therapeutic points of view. The histological examination consisted principally of the mucous membrane of the maxillary antrum after the radical operation. The cases were usually of a chronic nature and Uffenorde distinguishes (1) a catarrhal-edematous form showing a regression of the cell infiltration and edema and later proliferation of connective tissue; (2) a pustular form in which there is more or less round-cell infiltration of the stroma and subsequently considerable fibrous thickening; (3) a mixed form in which the processes of the two preceding forms occur. The inflammation always takes place from the superficial to the deeper parts, the involvement of the bone being secondary and not primary. Metaplasia of epithelium was found only in the pustular form. Uffenorde argues against primary disease of the maxillary antrum and believes that it is always secondary to infection from the nose.

Anatomically the catarrhal-edematous inflammation is characterized, aside from the absence of pus, by severe edema of the mucous membrane often showing large polypi and numerous cysts. The mucous membrane may easily be dissected off also. In the pustular form the mucous membrane is more vascular and in early cases moderately swollen. The fibrous hyperplasia is considerable and hence the removal of the mucosa from the bone is more difficult. In the mixed form the processes taking place in the two previous forms are in evidence.

Clinically, in the acute forms there were present inflammatory hydrops with typical, yellowish secretion from the nose. In the chronic cases the catarrhal serous form is characterized by polypoid swelling of the mucous membrane with polyp formation in the middle nasal passages, more rarely with polypi occluding the entire nasal cavity. The affection is usually bilateral. In the pustular form there is an absence of polyp formation and instead of swelling in the antrum there is atrophy. Not rarely there may be secondary infection.

Treatment. In acute cases operative measures are disadvised and expectant treatment recommended. The same also in the dematous form with orbital complications. In chronic cases, Uffenorde recommends operation by the Ritter-Jansen or the Luc-Caldwell method. He believes that these methods offer the best prospects of avoiding a recurrence of the formation of polypi and of curing respiratory complications by getting rid of the irritating hypersecretion. In sixty such radical operations there was an entirely successful outcome in all. In the pustular form favorable results may be obtained in 40 per cent of cases by means of conservative methods. Uffenorde sees the key to the recurrent polyp question in the view that the edematous-catarrhal form is diffuse and involves all the accessory cavities requiring their radical exposure and cleaning out.

P. F.

606

Roentgenology of the Accessory Nasal Sinuses, with Special Reference to Sinusitis in Children. C. W. WALDRON, *Interstate Med. Jour.*, Oct., 1915.

Sinusitis is relatively frequent in the young and is often undiagnosed. Children may come complaining of long-standing nasal discharge uncomplicated by headaches or general symptoms. In others, headaches, more generally frontal, may be the prominent symptom with negative eye and gastro-intestinal symptoms. Or the child complains of cough and expectoration. Examinations of the chest and sputum are negative and another source of the trouble must be looked for.

The intranasal examination for sinusitis is difficult and frequently of little value in differentiating between ethmoidal and antrum infection. The author places no reliance on the transilluminating lamp. The roentgenogram is essential in the rational treatment of either maxillary ethmoidal or frontal sinus because it is necessary to determine if the maxillary sinus is the source of the chronic discharge and should be treated by lavage or drainage, or whether the frontal sinus are affected and the anterior ends of the middle turbinates should be removed to facilitate drainage or whether the ethmoidal cells are alone involved and inhalations, irrigations or local applications should be the form of treatment.

P. F.

632

A Study of 1,000 Cases of Stammering, with Special Reference to the Etiology and Treatment of the Affection. G. HUDSON MAKUEN, *Volta Review*, July, 1915.

Age. The affection usually begins during the first few years of the developmental speech period which is between the ages of two and four or five. It is rare for the affection to begin after puberty, for by that time the speech habits have become more or less fixed (one patient began to stammer after he was twenty-three years of age).

Sex. Seventy-seven per cent. were in males and twenty-three per cent. in females. The reason for the smaller percentage in females Makuen ascribes to the greater natural aptitude of females for all work requiring the finer co-ordination of muscles. Aphasia is also less common in females than in males and stammering is a species of this condition. Both stammering and aphasia are undoubtedly of cerebral origin, the results of either functional or organic disturbances of cerebral speech centers and these centers appear to be more unstable and variable in the male than in the female.

Temperament. Emotional, sensitive, hesitating. The fear of the stammerer usually originates in his embarrassment at not being able to talk. Fear is the most dominant characteristic of the stammerer and is largely the result of the stammering itself. Afterwards it may become a secondary causal factor.

Mentality. Dividing his cases into three classes Makuen places in Class 1 those of average mentality—(85 per cent. of the whole number); in Class 2, those slightly below the average (14 per cent) and in Class 3, those of distinctly feeble mentality (1 per cent.). The fact

that so large a percentage is of average mentality, Makuen considers of importance because these individuals having more to say, feel more the importance of saying it than those in the lower grades of mentality. Difficulties of speech always increase with the desire to speak and with the necessity for speech.

Heredity. This is the most important factor in the etiology of stammering, notwithstanding the fact that it is an acquired affection in the sense that speech itself is an acquired faculty. Thirty-nine per cent of the patients admitted having or having had relatives who stammered.

Consanguinity. Twenty-nine patients, or 2.8 per cent. were offspring of consanguineous marriages.

Association. Association with other stammerers and either voluntary or involuntary imitation are undoubtedly factors in the etiology of stammering.

Fright or Injury. Twenty-eight per cent. of the patients dated the origin of their affection from the instant of having received a nervous shock. A very important factor in the etiology of stammering. The cerebral speech mechanisms have been thrown out of gear and have never righted themselves.

Vision. Twenty-six per cent. of the patients had faulty vision owing chiefly to a defective action of the muscles.

Audition. Only 3 per cent. had subnormal hearing and this seemed in no respect to be related to the affection.

Nose and Throat Conditions. No less than 97 per cent. complained of some nose or throat trouble and in 36.8 per cent. operations were performed for their relief. Over 37 per cent had intranasal pressure due to various irregularities of the septum and about 11 per cent. of them were operated on. About 60 per cent. had diseased tonsils and adenoids and half of them were operated on.

Treatment. Psychological and physiological. The stammerer must learn to idealize speech. If he thinks stammering speech he will stammer. From past experience he sees trouble ahead and the fear of this trouble amounts in some instances to an obsession, paralyzing all effort. The patient must be helped to remove the mental confusion arising from this fear.

P. F.

633

The Psychology of Stammering. G. HUDSON MAKUEN, *N. Y. Med. Jour.*, p. 117, July 17, 1915.

The author agrees that stammering is a "transient auditory amnesia." The four language centers in the cerebral cortex are divided into two groups: the auditory and glosso-kinesthetic centers constitute the "primary couplet," as Bastian termed it, representing spoken language and the remaining couplet, the visual and chiro-kinesthetic centers, represent written language. The difficulty of the stammerer is more with phonation than with articulation. As the articulatory element of a word is represented for the most part kinesthetically and as the phonatory element depends upon auditory stimuli, it follows that the disability exists chiefly in the auditory region of the cerebral cortex. While the weakness or hyperesthetic condition of the auditory speech-center is often congenital, stammering speech is, in a way, an acquired

affection. So far as heredity is concerned, Makuen states that 39 per cent of his patients admitted having had relatives who stammered, but the stammering, like deaf-mutism, is not inherited, only the physical anomalies that give rise to the affection. Sometimes it may skip one, two or even three generations.

Fear and autosuggestion are only secondary factors in stammering and are themselves the result of the individual's experience as a stammerer and, if they do not become too well established, the stammering will cease.

Makuen is of the opinion that psychoanalysis, so far as treatment is concerned, is of little avail. Arousing and training the patient's auditory imagery through the proper use of the peripheral organs of speech or psychophysical therapeutics offers the better chances of improvement. The prognosis must, however, depend upon the degree of the underlying anomalous physical condition giving rise to the amnesia.

P. F.

684

Technic of Laryngeal Heliotherapy. DR. COLLET (LYON), *Revue de Laryngologie, D'Otologie et de Rhinologie*, Sept. 30, 1915.

Collet advocates heliotherapy in tuberculous laryngitis, and divides the methods into Direct and Indirect. He believes that the external method is ineffective as the ultra-violet rays, which have the greatest efficacy, are rapidly arrested and do not penetrate the external tissues.

In the "stomatodiale" method the direct sun rays are allowed to enter the larynx in a manner similar to that used in the practice of direct laryngoscopy. For the indirect method, Collet suggests an ingenious arrangement by which the patient himself projects the sun's rays into the larynx by means of a laryngeal mirror.

W. SCHEPPEGRELL.

698

Treatment of Laryngeal Tuberculosis in Sanatoria for Pulmonary Tuberculosis. T. BERGTRUP-HANSEN. *Ugeskrift for Læger*, Sept. 23, 1915.

During the last two years the author has observed at the Silkeborg Sanatorium, 63 cases of laryngeal tuberculosis, the diagnosis of which was based on ulceration, infiltration and granulation. But in only two of these cases was the laryngeal disease a more important factor than the pulmonary disease. In 55 cases the pulmonary disease was in the third and in 8 cases it was in the second stage. In 21 cases with severe laryngeal disease the patients were debilitated and febrile; hence the treatment could only be palliative. The only operative treatment was an occasional amputation of the epiglottis for dysphagia or an injection of alcohol into the laryngeal nerves. In 29 cases of slight laryngeal disease, the condition ran parallel with the pulmonary disease and the only treatment in these cases was the application of menthol and lactic acid in addition to the sanatorium treatment. The author concludes that radical operative treatment should be confined to few cases and should be undertaken when the condition of the lungs is not hopeless. Too active treatment of the larynx in several cases in his experience have proved detrimental to the lungs. Three advanced cases of laryngeal tuberculosis were greatly benefited by artificial pneumothorax, the ulcerations in two cases healing completely.

P. F.

747

Operative Treatment of Laryngeal Tuberculosis. TH. RUEDI, *Zeitschr. f. Ohrenheilkunde*, Bd. 73, H. 3, Nov., 1915.

From 1908 to 1914 the author examined about one thousand cases of tuberculosis of the larynx, 575 of which were operated upon laryngeally in a total of 1,548 sittings. Two-thirds of the patients were males, one-third females. In all cases the laryngeal tuberculosis was secondary to pulmonary tuberculosis. In many cases the clinical diagnosis was controlled and confirmed pathologic-anatomically. The 1,548 sittings were as follows: Curettement, 61 sittings; curettement and cauterization, 168 sittings; electrocautery alone, 1,319 sittings. Where there was a history or suspicion of syphilis a Wassermann test was made and if positive salvarsan and potassium iodid treatment instituted.

Without regard to the stage of the pulmonary tuberculosis, Rüedi divides his cases into three classes: (1) *Light cases*, in which there is a single, circumscribed focus without any tendency to rapid progress or rapid destruction; (2) *moderately severe cases*, with multiple, circumscribed foci without any tendency to rapid progress and rapid destruction and (3) *diffuse or severe cases*, (a) yielding to treatment and (b) characterized by fever, rapid destructive process and perichondritis.

There were 259 cases belonging to the first class, 265 to the second class and 51 to the third class.

The conclusions reached by the author are: (1) Laryngeal tuberculosis is curable. (2) In Davos (a tuberculosis *kur* place) spontaneous improvement and cure are often observed. Many cases not reacting to general treatment or conservative local treatment, notwithstanding the improvement in the pulmonary condition, may be permanently cured by operative treatment. (3) Operative treatment is indicated where there is no fever and where preferably the lung process is at a standstill. (4) The best operative procedure is the electrocautery, according to the Mermod-Siebenmann method, but in tuberculosis of the epiglottis, resection or amputation is more advisable. (5) In more than one-third of the cases there was a cure in at least three months after the last treatment. The best results were obtained in tuberculosis of the vocal cords with the electrocautery (52 per cent. cures). (6) In many cases operative treatment also induces a favorable influence on the lungs and the general condition. (7) The view that in laryngeal tuberculosis a high altitude *kur* is contraindicated, is false. P. F.

753

Laryngeal Papilloma Treated and Cured by the Roentgen Rays. I. G. SHALLCROSS and W. D. BAYLEY, *Jour. Ophthalmol., Otol. and Laryngol.*, Oct., 1915.

A woman of twenty-four years of age suffered from aphonia for about eight months, the condition first starting as a slight hoarseness and gradually progressing to complete aphonia. No dyspnea and no pain were complained of. Examination showed a papilloma about the size of a barley grain attached under the middle third of the right vocal cord near its free edge. Operation was performed on three different occasions but the tumor recurred and it refused to yield to any form of treatment,

including the galvanic cautery. The Roentgen rays were finally resorted to under which treatment, after several weeks, the tumor became smaller and lighter in color. After fifteen months' treatment the tumor finally had disappeared, leaving a small, thickened spot at the site of the original tumor. Twice weekly this was treated with argyrol. The patient regained her voice and all traces of the tumor disappeared. P. F.

754

Laryngeal Diverticula. GEORGE E. SHAMBAUGH and DEAN LEWIS, *Annals of Surgery*, Vol. LXI., No. 1, Jan., 1915.

Such diverticula are rare. Many cases in earlier literature were thyroglossal cysts, vascular struma, or a localized emphyema resulting from perforation of the cartilages of the larynx, by some inflammatory process, such as tuberculosis. Larrey was the first to give an accurate description of laryngeal diverticula. He saw them in Egypt in the blind who were employed by the priests to recite the Koran from the minarets. The authors summarize briefly the observations of other observers. They report a case of combined type of extra- and intralaryngeal diverticulum which they observed, and describe the operative treatment which they pursued. PACKARD.

758

The Conservative Treatment of Intrinsic Cancer of the Larynx by Thyroecricotomy or Thyrotomy. GEORGE DAVID STEWART, *Annals of Surgery*, Dec., 1915.

Operation for intrinsic cancer of the larynx offer better results with less crippling than possibly any other site in the body. This is due to several facts.

1. Early discomfort causes patient to seek examination.
2. Scanty lymphatic draining larynx and the restraining influences of the cartilage.
3. Intrinsic cancer of the larynx maintained by many or essentially a slow-growing neoplasm.

It is always advisable to remove section for examination. Difficult if growth is under cords, may have to do thyrotomy.

TECHNIQUE OF OPERATION.

1. Tracheotomy may be done at the time of removal of the growth or two days before.
2. Split thyroid cartilage in the angle between its protecting the anterior insertion of the vocal cords.
3. After thyroid is opened, paint larynx with 4 per cent. solution of cocaine to prevent coughing.
4. Tampon upper portion to prevent soiling from mouth.
5. Outline growth and remove.

If the arytenoid is involved, or the growth crosses the posterior commissure a partial laryngectomy must be done.

6. Wound can be closed without drainage.

After-treatment consists in keeping the head low until patient comes out of ether. Moist air in the room. Remove tube in 48 hours, but may have to replace on account of edema. Get patient up on the third day.

Feeding is difficult and may have to use nasal tube. Mortality small. Speech always possible, often satisfactory. This operation applicable in one-third of all cases of laryngeal cancer. Contraindicated if arytenoid cartilage and interarytenoid fold involved, also in other extrinsic cancer.

PACKARD.

761

Granuloma of the Vocal Cord. SIR ST. CLAIR THOMSON, *Jour. Laryngol., Rhinol. and Otol.*, Nov., 1915.

The patient, a male, aged forty, complained of hoarseness for the previous two months. Examination of the larynx showed a pedunculated, mobile tumor adherent to the free edge of the right vocal cord just in front of the processus vocalis. The growth was avoid, smooth, purplish and the attached extremity slightly yellowish and puckered. During inspiration it would fall below the glottis. Movement of the right cord was free. Some enlarged glands at the angle of the jaw. The growth was removed with a Mackenzie forceps by the indirect method and histological examination showed it to be a tumor consisting of fibrin and granulation tissue, covered by squamous-celled epithelium showing no evidence of malignancy.

The patient gave no specific history or a history of traumatism to account for the growth and it was situated too far back on the cord to be connected with nodular thickening produced by misuse of the voice. The author suggests that it probably originated in a submucous hemorrhage with incomplete absorption of the effused blood and irritated into a granuloma by misuse or overuse of the voice.

P. F.

772

Lung Gangrene from Primary Stone of Bronchus. BLECHER, *Mitteilungen d. Grenzgeb. d. Med. und Chir.*, Bd. 28, H. 4, 1915.

Blecher's patient, a man 23 years of age, had had an attack of what was diagnosed as pleurisy from which recovery was without incident. A month later the patient began to cough and complained of pain in the right side. The trouble proved to be a primary stone in the bronchus of the lower lobe, complicated by gangrene and pyopneumothorax from gas-producing suppuration. Outcome fatal from peritonitis.

P. F.

774

Foreign Body Pneumonias. A. CAILLE, *Arch. Pediatrics*, Dec., 1915.

Three cases are reported by Caille. In the first a large tack was lodged in the right lung for about five weeks. In the second case the roentgenogram showed a large nail in the left bronchus and partly in the trachea. In the third case a shawl pin was found in the left bronchus and lung, the point projecting at the bifurcation of the trachea. It had been in the lung for 43 days without inducing local inflammation. In the first two cases the foreign bodies were removed by bronchoscopy; in the third, the patient, during a spasmodic coughing spell, coughed up the pin, which was two and one-half inches long. All the symptoms cleared up after removal of the foreign bodies.

P. F.

824

Fistula of Esophagus and Bronchus. Report of Case, with Roentgenologic Findings. RAYMOND COLE BEELER, *Journal of American Medical Association*, Oct. 2, 1915.

Patient admitted to hospital for Roentgen examination of esophagus. Middle-aged man, history of syphilis some forty years previous; no history of swallowing escharotics. Had difficulty in swallowing that had gradually gotten worse over a period of six months. A week before had hemorrhages and suffered from violent coughing spells when he tried to eat or drink.

Given barium and the fluoroscope showed it had passed down the esophagus a short distance and then to go through and fill the bronchi. Further x-ray examination showed a mass of barium just above the bifurcation of the bronchi, probably a dilation at the region of the fistula.

Patient died one week later from pneumonia but no autopsy was obtained.

PACKARD.

839

Esophageal Stricture in Children. C. W. M. HOPE, *Proc. Roy. Soc. Med., Laryngological Section*, V. 8, No. 6, April, 1915.

Case 1. Boy, seven and a half years old, has vomited at least once a day ever since the age of two months. X-ray examination showed a well-defined stricture of the esophagus behind the pericardium. Examination with the esophagoscope showed enormous dilatation of the upper esophagus containing much undigested food. The mucosa above the stricture was denuded in parts, bled easily on swabbing and showed scars on the right side. Stricture admitted a 5 mm. bougie and could be dilated up to 10 mm. Since the dilatation was done the patient only vomited once and has gained about one pound every day. He is now able to take solid food, having lived up to now on liquid food.

Case 2. Eighteen-months-old baby with a history of daily vomiting since birth. Examination with the esophagoscope showed the whole of the thoracic esophagus greatly dilated but no food in dilated portion. Bougies and the second largest Killian tube could be passed into the stomach without difficulty. The mucosa of the sub-diaphragmatic portion was thrown into large transverse folds. After simple dilatation, the child can now eat bread and butter.

P. F.

874

Asthmatic Attacks and the Inhalation of Flour Dust. EMILE A. BERTUCCI, *New Orleans Medical and Surgical Journal*, Sept., 1915.

The causative relation of the flour in this case was demonstrated by dusting the air surrounding the patient with wheat flour, which was followed by a short attack of asthma.

SCHIEFFEGRELL.

926

Report of Intravenous Injection of Diphtheria Antitoxin. E. M. DUPAQUIER, *New Orleans Medical and Surgical Journal*, Sept., 1915.

Intravenous injection of diphtheria antitoxin, which is usually reserved for severe cases, is recommended as routine in all cases. It is less pain

ful and more swiftly acting as the neutralizing antitoxin is brought at once in contact with the toxins. He believes that striking serum reactions are not a contra-indication, and that when the technic is perfected, it will be the routine method of treating diphtheria. SCHEPPEGRIEL.

948

Schick Toxin Reaction for Immunity in Diphtheria. J. A. KOLMER and E. L. MOSHAGE, *American Jour. Dis. Children*, March, 1915.

The authors conclude that the toxin skin reaction is a valuable and reliable method for detecting susceptibility to diphtheria and that persons reacting negatively to this test usually contain at least one-twentieth unit of diphtheria antitoxin per cubic centimeter of serum, which amount of antitoxin is probably sufficient to protect against infection. Persons reacting positively to the test usually contain less than one-fortieth of a unit of antitoxin per cubic centimeter of serum and may be regarded as susceptible to infection; hence, in the event of exposure to infection, these persons should be passively immunized by an injection of antitoxin.

The studies of the authors show further that about 40 or 50 per cent. of children between 1 to 15 years of age react positively to the Shick test and that immunity conferred by a prophylactic injection of antitoxin begins to disappear after ten days and has generally passed away entirely after four weeks.

Another interesting point that the Shick test has shown is that scarlet fever patients have an increased susceptibility to diphtheria, 10 per cent. of them being susceptible within ten days, even after the injection of antitoxin. Another point is that the immunity conferred by an attack of diphtheria is of short duration or nil. P. F.

1011

A Case of Aberrant Thyroid. EDWIN L. DRAPER, *Albany Medical Annals*, Dec., 1915.

A married woman, 41 years old, mother of one child, complained of gradual swelling or enlargement of her abdomen. She had had a mastoid operation a year and one-half previously. In addition to the abdominal enlargement, the patient was short of breath on exertion.

Examination: slight facial paralysis (which had persisted since the mastoid operation). Heart and lungs negative. In the dorsal position both flanks were flat to percussion, tympanitic area 12x14 cm. in region of umbilicus shifting with change in position. Abdominal examination, because of the amount of fluid present, was otherwise negative. Blood and urine negative. Vaginal examination showed a distinct sense of resistance on right side.

Exploratory operation showed a mass on the right side with the broad ligament which proved to be a dermoid cyst. At the upper, inner and anterior corner of this mass were two walnut-sized masses closely connected with and protruding from the large mass, which, upon microscopic examination proved to be perfectly typical thyroid tissue of the colloid type. There was also a small cyst of left ovary which was also removed.

P. F.

1036

Comparison of Autoplastic and Homeoplastic Transplantation of Thyroid Tissue in Guinea Pigs. C. HESSELBERG, *Jour. Exper. Med.*, Feb., 1915.

A short time after operation no difference is noticeable in the behavior of the thyroid after autotransplantation and homeotransplantation. Later, the follicles in the homeografts begin to undergo destruction. This destruction is not caused by a direct primary disintegration of the follicles but is caused by the destructive activity of the lymphocytes and the connective tissue of the host tissue. The connective tissue grows into the homeografts in larger quantity than into the autografts. In the homeografts it becomes hyaline and fibrous, while in the autografts it remains cellular. Destruction by invasion of lymphocytes is direct. The rapidity and the degree of destruction in homeografts vary. In some, destruction by lymphocytes preponderates, in others, by connective tissue. The blood supply developed in the autografts is more abundant than in the homeografts.

P. F.

1050

The Isolation of the Compound Containing Iodin which Occurs in the Thyroid; Its Chemical Nature and Physiologic Activity. E. C. KENDALL, *Jour. A. M. A.*, June 19, 1915.

By alkaline alcoholic hydrolysis the author has been able to split up the thyroid proteins into two groups: group A are the acid insoluble compounds and group B are acid soluble. From group A a pure crystalline compound containing 60 per cent. of iodine and which appears to be di-iodo-di-hydroxy-indol, has been isolated. Group B contains iodine in an unknown form of combination. The physiologic action of the group A compound consists of an increase in the rate and vigor of the pulse, in metabolism and in nervous irritability. Group B compounds are more toxic except in cases of cretinism, myxedema and some cutaneous conditions.

P. F.

1160

Acute Otitis Media in Infancy and Early Childhood; Avoidable Mistakes in Diagnosis, Prevention and Treatment. W. R. P. EMERSON, *Boston Med. and Surg. Jour.*, Oct. 21, 1915.

Acute otitis media is the most frequently overlooked affection in infancy and childhood because the symptoms may not be at all distinctive. They may simulate pneumonia, meningitis, or just general symptoms with fever. The author reports five cases in which no earache was present and four of the cases had a double otitis media. In all cases, therefore, the ears should be examined as a routine measure whether the symptoms point to the ears as the source of the trouble or not, in the same way as the physician would examine the heart in rheumatism or the abdomen in typhoid. As to prevention, in every case of contagious disease or in affections of the respiratory tract in children, the nasopharynx should be kept clear by the use of a soothing spray to the mucous membrane and the instillation of a few drops of 10 per cent. solution of argyrol. Treatment of the onset of otitis media consists of the instillation of a few drops of adrenalin and one-half per cent. solution of cocaine followed by a few drops of 20 per cent. argyrol. If the symptoms do not subside, the drum should be incised.

P. F.

1161

A Case of Subacute Purulent Otitis Media Labyrinthitis and Purulent Leptomeningitis Due to a Capsulated Streptococcus; Spontaneous Recovery. J. S. FRASER and J. L. OWEN, *Edinburgh Med. Jour.*, Oct., 1915.

The patient, 32 years old, suffered a mild attack of otitis media without otorrhea in January, 1915. The trouble recurred in February and at this time appears to have been accompanied by labyrinthine or intracranial irritation as evidenced by headache and vomiting. No discharge from the ear was present at this time. Otorrhea only commenced in the beginning of March. The patient was confined to bed for three weeks but resumed work about March 20, but after he had been at work only a week he became very ill with symptoms of purulent labyrinthitis, rapidly passing on to leptomeningitis. The stage of delirium followed by coma lasted only forty-eight hours. The patient made a complete recovery as far as balancing is concerned, the healthy side compensating the loss of function of the other vestibular apparatus. The only inconvenience felt by the patient is the unilateral deafness. P. F.

1164

Prognosis and Treatment of Severe Acute Middle Ear Disease. F. KOBRACK, *Therapie der Genickart.*, Bd. 55, H. 6, June, 1915.

According to Kobrak blood cultures, which should be made in every case of acute otitis, if negative would indicate that conservative measures are justified for several days, even in the presence of high fever, provided there are no local or general signs of complications. Instead of a stormy onset from the first the otitis media may start insidiously and after a mild course apparently subside. But soon a serious complication. In such cases, the author has found, the streptococcus mucosus capsulatus is the responsible organism and he warns that this type of otitis should be kept under strict supervision even after an apparent cure. Continued sub-febrile temperature will warn that the process is progressing insidiously. P. F.

1185

Otitis Media in Relation to Infant Pathology. F. SCHERER and O. KUTVIRT, *Jahrbuch für Kinderheilk.*, Bd. 82, H. 2, Aug., 1915.

Fully in 22.3 per cent. of congenitally syphilitic infants and in 30.4 per cent. of tuberculous infants, the authors found otitis media; in both of these groups the ear disease developed insidiously. They found otitis media in 18.6 per cent. of nutritional disturbance and in as high as 25 per cent. of respiratory affection. They call attention to the gravity of the disease in infants and often convulsions of apparently obscure etiology are due to suppurative otitis media. P. F.

1187

Acute Otitis Media. Report of a Case in Seventeen Months Old Child. CLARENCE H. SMITH, *New York Medical Journal*, Nov. 20, 1915.

Author reports a case of bilateral acute otitis media developing in a child after an attack of influenza and bronchitis. There was no tender-

ness over the mastoid. Drums were incised and three days later an abscess of the elbow developed. An abscess of the great toe had healed previously. Diagnosis of pyemia was made, developing from septic sinus thrombosis. The mastoid on the left side was opened, but the mastoid cells were normal. Three days later the left sigmoid sinus was exposed and a large clot of blood removed and some pus liberated. On the next day the jugular was exposed and tied. About ten days later the right leg was greatly swollen and the inguinal glands enlarged. Child discharged with ear condition well, but still shows signs of septic arthritis.

PACKARD.

1188

Radium Therapy in Diseases of the Ear. D. J. SPINETTO, *Semana Medica*, Aug. 5, 1915.

After it was proved that radium was not detrimental to the sensorial apparatus of the ear and that it was capable of producing a retrogression of fibrous tissue, it was employed by Spinetto in chronic middle and internal ear affections showing a tendency to fibrous connective tissue proliferation. A glass tube containing ten mgm. of radium bromide was covered with black paper and rubber tissue and introduced into the ear so that it came in contact with the tympanic membrane. It was allowed to remain in contact for ten minutes. At intervals of eight to ten days, these periods were lengthened by five minutes. Improvement in hearing and beneficial changes in the tympanum were noted.

P. F.

1194

Tuberculosis of the Middle Ear Cleft in Children.—A Clinical and Pathological Study. A. L. TURNER and J. S. FRASER, *Jour. Laryngol., Rhinol. and Otol.*, June, 1915.

Cases of tubercular otitis media may be divided into two groups: (1) tubercular otitis in infants and young children who have been fed in whole or in part on unsterilized cows' milk containing tubercle bacilli. Tubercular otitis media as compared with other forms of otitis in infants and young children is exceedingly frequent. The authors' statistics show that 27 per cent of the cases of purulent otitis in children under two years of age are due to tuberculosis while the same is true in 50 per cent. of the cases in children under one year of age. On the other hand, tubercular otitis of all cases during all ages accounts for 2.8 per cent. There is no direct proof as yet that tubercular otitis in the infantile cases is caused by the bovine type of tubercle bacillus but there is good reason to presume so.

(2) The second type of tubercular otitis media occurs in the advanced stages of pulmonary tuberculosis. This form is not very frequent.

The route of infection in tubercular otitis media is still unsettled: (1) through the Eustachian tube and (2) by way of the blood stream. In the former route the infection may extend by tubercular infiltration of the mucous membrane of the tube or else the tympanic cavity may become infected by the insufflation of infectious particles from the naso-pharynx. In the latter route, hematogenous infection is a possibility if the mastoid

alone is diseased and the tube and tympanum are healthy. Direct extension through the Eustachian tube is perhaps the more probable route of infection.

The clinical types of tubercular otitis media may be classified as (a) lupoid, in cases of lupus from the nose and throat; (b) an infiltrating form which progresses rapidly and shows numerous tubercle bacilli but few giant cells; (c) a chronic form characterized by few bacilli and many tubercle follicles, the mucosa being greatly thickened and showing a tendency to polyp formation. This form also shows a tendency to encapsulation of the tubercle follicles and spontaneous cure of the condition. (d) A necrotic form which undergoes rapid caseation and destruction of the mucosa and of the bony capsule of the labyrinth. (e) A fibrinoid form in which a false membrane occurs on the tympanic mucosa and (f) as shown in the present paper by the authors, a fibro-ossifying type.

Involvement of the labyrinth in cases of tubercular otitis media is frequent, varying from 23 to 33 per cent. The statistics of the authors show that necrosis of the labyrinth was present in 22 per cent. of the cases operated on, while in a further 31 per cent. there was erosion of the labyrinthine wall. Clinically, tubercular labyrinthitis, like tubercular otitis media, appears to have an insidious onset, in marked contrast to the acute symptoms produced by acute purulent labyrinthitis.

The diagnosis of tubercular otitis is made by attention to the following points: (a) Clinical characteristics. (b) The operative findings. (c) The examination of the ear discharge for tubercle bacilli. (d) Microscopic examination of the "granulations." (e) Animal inoculation with the ear discharge. (f) Differentiation of the human from the bovine type of tubercle bacillus by making cultures on egg medium from the tubercular lesions in the guinea pig injected and the later injection into rabbits.

P. F.

1199

A New Treatment of Middle Ear Disease. C. E. WILLIAMS, *Jour. Ophthalmol. and Laryngol.*, June, 1915.

When treatment is begun the patient usually calls every other day for two weeks and then the treatment is continued at the same or less frequent intervals according to the results of the tests. With the Siegel otoscope the air in the external auditory canal is alternately rarified and condensed from ten to twenty times. The force exerted should be very mild at the start and gradually increased until the tympanic membrane shows greater mobility. Best results follow the gentler massage which is not followed by intense injection of the drum-head.

Following this, Dowling argyrol tampons are placed and left for a period of from ten to sixty minutes. After removal of the tampons the naso-pharynx and nasal fossae are douched thoroughly with a mild alkaline solution followed by an oily spray and the inhalation of an oily vapor. Where the naso-pharyngeal tissues show hypertrophy this should be treated by Lugol's or argyrol solution on an applicator. Adenoids should be removed surgically. The treatment is completed by massage from thirty to one hundred and twenty seconds over each ear. P. F.

1210

Sarcoma of the Cartilaginous Eustachian Tube. DR. JACOD (DE LYON), *Revue de Laryngologie, D'Otologie et de Rhinologie*, Aug. 31, 1915.

Jacod reports two cases, the first without and the second with operation, both of which terminated fatally. The first symptoms observed in sarcoma of this region are auricular, especially quickly developing unilateral deafness, and is usually accompanied by the sensation of a foreign body in the nasopharynx, and violent pain referred to the affected ear. The development of the sarcoma is unusually rapid and the prognosis very grave.

After discussing the various operations, he believes that the resection of the superior maxilla is the most practical route for enucleating the noplasm.

W. SCHEPPFEGRELL.

1236

The Origin of Labyrinthine Rest-tone. E. P. FOWLER, *Jour. A. M. A.*, Jan. 9, 1915.

Fowler reaches the following conclusions:

(1) Physiologic endolymph movements excite impulses from the end-organs which are interpreted as sensation complexes from all the ampullae. If the impulses from the two labyrinths approximately balance (or through practice are balanced by the aid of the co-ordinating apparatus), no sensation of movement is experienced. Binaural galvanic or caloric reactions with the head anywhere in the antero-posterior planes demonstrate this clearly, as does the rotation of bilaterally non-functionating labyrinths.

(2) The body normally receives a sensation-complex (and a tonus) in the antero-posterior vertical plane, approximately equal from its both sides, but in all other positions various complexes must be forthcoming, depending mainly on the position of the head.

(3) Balance tonus originates from all the labyrinth end-organs.

(4) Endolymph stress is sufficiently capable of maintaining static control in rest, as it does so in action.

(5) Positive endolymph stress, within the canal at rest, can occur only because of connection currents present therein.

Fowler regards convection currents as due (1) to a difference in temperature in the inner and outer labyrinth walls (this does not take place in health) and (2) to a constant but variable difference in temperature within the ampullae and the non-ampullated portions of the canals, because the blood-heat is transmitted mainly by conduction and radiation to the ampullae and as the tiny branches lie along the concavity of the canals, they receive less and less heat the further the blood courses from its source, for the reason that blood-vessels coming from within the cranium contain blood of a higher temperature than those from without and the internal auditory artery traverses the inside of the skull before distributing its blood-heat to the ampullae. The specific gravity of the endolymph adjacent to the ampullae is lowered as a result of the difference in temperature and within every canal not on the horizontal plane a current is impelled by the force of gravity, causing stress on the capulae of all the canals at all times.

P. F.

1239

War Deafness from Lesions of the Internal Ear. A. GOR, *Gazette hebdomadaire des sciences med.*, July, 1915.

Since the beginning of the war the author has had under observation about one hundred cases of deafness due wholly or in part to disease of the internal ear. All the cases showed cochlear hypo-excitability. Of seventeen deaf as a result of shell explosion, bombs, grenades, etc., four were hyper-excitability as regards the vestibule, eleven hypo-excitability, none were unexcitable, two were normal. Of thirteen deaf as the result of direct wounds of the head, five were hyper-excitability, four hypo-excitability, two unexcitable and two normal. From the etiological standpoint the cases (thirty-two in all) fall into two groups: one in which the injury resulted from explosions acting through the air as an intermediary, the other which resulted from projectiles inflicting direct injury to the head. In the former the hemorrhages are probably less severe than in the latter, but there is probably concussion of the organ of Corti with or without disintegration of the membrane. As regards prognosis, the author found that the cases improve fairly quickly, even with regard to the cochlear reaction, but the only cases in which he has seen a complete cure are the psychic cases. In the organic cases during improvement from the vestibular point of view, the vertigo, along with the nystagmus, is usually the first symptom to disappear and gradually equilibrium tends to become more normal. On the cochlear side, however, the symptoms tends to persist.

P. F.

1244

Syphilis of the Internal Ear (Hereditary). H. HASTINGS, *Jour. A. M. A.*, Aug. 14, 1915.

Patient, a girl twenty years old. Her parents had been infected with syphilis but considered themselves cured. When a child, the patient had had trouble with her eyes (interstitial keratitis) but otherwise her general health had been good. Her hearing was greatly impaired and she also had nystagmus and vertigo. Wassermann (blood) negative. Salvarsan being objected to, the patient was put on mercurial treatment, under which her hearing improved.

P. F.

1246

Isolated Vestibular Neuritis Following Typhoid Inoculation. C. HIRSCH, *Deutsche med. Woch.*, Aug. 19, 1915.

An army surgeon was twice inoculated with typhoid vaccine. The first inoculation was well tolerated but twenty-six hours after the second inoculation the patient became exceedingly giddy and developed severe vomiting on the slightest movement of the head. During the following night the pulse and respiration were abnormally slow, spasmodic contractions of the jaw muscles, tinnitus and numbness of the fingers. The eye grounds showed nothing abnormal. Diagnosis of hemorrhage into the left labyrinth was made. Four weeks after the appearance of the first symptoms the patient still felt giddy on turning his head or eyes, especially to the left. Marked ataxia on rapid movement; no spontaneous nystagmus or deafness. After three months the patient was able to some extent to resume his surgical work.

P. F.

1259

Galvanometric Studies of the Cerebellar Function. I. L. MEYERS, *Jour. A. M. A.*, Oct. 16, 1915.

The experiments conducted by Meyers were on two groups of cats. In one group the right lobe of the cerebellum was removed and in the second group the left lobe. The animals were kept under observation for from one to three weeks and of all the animals he selected seven, discarding all those that did not show marked unilateral ataxia and whose wounds did not heal promptly. He is of the opinion as a result of his experiments that the function of the cerebellum is that of control and inhibition, each half exhibiting its function on the opposite half of the cerebrum. Lesions of the cerebellum do not cause sensory disturbances or disturbances of the muscular sense. The phenomena following cerebellar lesions are motor. Forced movements, circus movements, nystagmus, conjugate deviation of the eyes and the characteristic attitude of the head in some cases of cerebellar disease in man or after unilateral ablation of the cerebellum in animals are not cerebellar in origin but vestibular in origin and due either to a lesion of the vestibular complex or its oculomotor tracts.

Meyers further believes that there is an association between the cerebral hemisphere on one side and the cerebellar on the other side, the cerebellum being subservient to the cerebrum. In short, the cerebellum has no direct effect on the periphery but acts primarily on the motor cortex, the paracerebellar nuclei and perhaps also on the basal ganglia and rubral tract. Its primary effects inhibition, control and regulation of the activity of these structures.

P. F.

1269

Translabyrinthine Removal of Acoustic Tumors. E. SCHMIEGELOW, *Ztschr. f. Ohrenheilk.*, Bd. 73, H. 1, July, 1915.

The author gives an historical review of the development of the paracerebellar method of Krause and the translabyrinthine method of Panse. On the basis of two successful cases of his own and the similar cases of Quix and Kümmel Schmiegelow is in favor of the translabyrinthine method. Both cases, which were fibrous acoustic tumors, are described. The advantages of the translabyrinthine method over the paracerebellar are that the route through the labyrinth is shorter, the entire operation is extradural and there is less danger of serious hemorrhage. Furthermore, of the four cases operated upon by the translabyrinthine method, not one patient died, whereas by the paracerebellar method the mortality is between seventy and eighty per cent.

P. F.

1280

The Differential Diagnosis of Lesions of the Labyrinth and of the Cerebellum. J. G. WILSON and F. H. PIKE, *Jour. A. M. A.*, Dec. 18, 1915.

One sharp distinction between cerebellar function and labyrinthine function lies in the fact that the cerebellum is an afferent station for the proprioceptive impulses and the labyrinth is a peripheral sense organ. The differentiation in lesions of the cerebellum and of the labyrinth lies essentially in the following symptoms: Nystagmus. When

the nystagmus is due to a labyrinthine lesion it consists of two phases, a slow lateral deviation, labyrinthine in origin, followed by a quick return movement cerebral in origin. The movements are synchronous in both eyes and vary indefinitely in direction with the labyrinth affected. The horizontal movement is of importance, the rotary not being of diagnostic value. The quick deviation is to the sound side. Cerebellar nystagmus is ataxic. It may be from or towards the sound side, according as we are dealing with an irritative or a destructive lesion, but with the head at rest it tends to be irregular in the plane of its direction.

Vertigo. In a labyrinthine lesion the plane of rotation of self is in the direction of the rapid phase of nystagmus. In regard to rotation of objects Wilson and Pike find it as a rule contrariwise to the preceding actual rotation. In the absence of nystagmus the apparent rotation of self is to the side of the lesion. In cerebellar vertigo the relationships are the same as in labyrinthine lesions.

Ataxia. In both cerebellar and labyrinthine lesions there is a deviation in walking to the side of the lesion if it is unilateral. But this deviation has nothing to do with ataxia. It is a disturbance of orientation in space which the patient corrects when his eyes are open; but with his eyes shut he overcorrects it and deviates to the sound side. As regards ataxia the cerebellum is the one, but not the sole, central organ which participates in the integration of impulses from the labyrinth.

Labyrinthine lesions differ from the cerebellar in (1) the presence of Romberg's sign; (2) variations in the attitude of the head influence the lack of equilibrium in labyrinthine and not in cerebellar lesions; (3) in labyrinthine disease movements of rotation or disorientation are not so readily perceived; (4) in labyrinthine disease the caloric and rotation tests are lost.

P. F.

1320

Prognosis of War Deafness. LANNOIS and CHAVANNE, *Bull. de l'Acad. de Med. de Paris*, Dec. 21, 1915.

The prognosis depends upon two factors: (1) The previous condition of the auditory apparatus; (2) the existence or otherwise of a direct injury to the skull. Under the first factor the authors found that men suffering from chronic suppurative otitis media, or well-marked sclerosis, after concussion of the labyrinth from shell explosion, showed a far greater proportion of deafness or impaired hearing than healthy persons. Of 189 men sent back from the front for chronic suppurative otitis media intensified by the concussion of shell explosions, only 43 per cent were returned to the front, while 56 per cent were relegated to the auxiliary services, and 1 per cent were discharged as unfit. Of 134 cases of sclerosis sent back because of defective hearing, 48 per cent were passed into the auxiliary services, and 52 per cent were returned to the front.

Under the second factor the authors found that in about 95 per cent of cases, deafness follows traumatic mastoiditis. In from 30 to 40 per cent of cases of traumatism to the branches of the facial nerves distributed to the ear, deafness was present. In both of these cases it was generally unilateral deafness that was produced. Without any direct traumatism concussion of the labyrinth alone rarely produced deafness.

So far as treatment is concerned in definite bilateral deafness recourse must be had to lip-reading as no treatment would be of avail. On the other hand, in simple deafness, associated or not associated with dumbness, the prognosis was very good the earlier the patient was placed under treatment.

P. F.

1324

A Case of Hysterical Deafness. A. M. MOTT, *Medical Jour. of South Africa*, Aug., 1915.

A school girl of eighteen years, passionately devoted to music, was depressed as a result of successive disappointments at school and during a prolonged emotional crisis a fellow-pupil told her the story of a young lady who, through deafness, had to forego the pleasures of music, the theater and all the other attractions of social intercourse. The patient, as a result of her hysteria, became totally deaf, in contrast to the more usual unilateral deafness, accompanied by hemianesthesia on one side, an attitude of marked indifference to her condition and absent-mindedness.

Hypnotic and psycho-therapeutic treatment effected a complete cure.

P. F.

1332

Typhoid Deafness. RHESE, *Medizinische Klinik*, Nov. 7, 1915.

Without including cases of transient deafness, Rhese states that of all the typhoid cases in his service (the exact number is not given) there was bilateral deafness in 14.9 per cent, the middle ear being intact. In 4 per cent there was otitis media with perforation, and in 1.1 per cent without perforation. The interesting point is that all the cases of deafness were bilateral. Not all the cases remained permanently deaf. Some did, while most regained their hearing in time. The onset of deafness was during the second, third or fourth week, as a rule.

The author states that vaccination against typhoid is of great importance in reducing the frequency and severity of ear complications. In the treatment of typhoid deafness, he obtained good results from mild courses of pilocarpin and also sodium iodid.

P. F.

1358

Syphilitic Lesions of the Ear. J. V. F. CLAY, *Jour. Ophthal., Otol. and Laryngol.*, Feb., 1915.

After considering the syphilitic manifestations of the external ear and the very indefinite symptoms of middle ear syphilis Clay takes up the question of syphilis of the internal ear and the auditory nerve. Involvement of the auditory nerve and internal ear usually occurs in the latter part of the second stage, so-called, or later, and forms one of the most frequent forms of primary disease of the aural perceptive apparatus.

Syphilis of the internal ear may be divided into three clinical types: (1) that occurring during the secondary or tertiary periods; (2) chronic syphilitic labyrinthitis; (3) labyrinthitis of acquired syphilis secondary to chronic suppurative middle ear disease.

The acute acquired form of syphilis of the internal ear, if recognized and treated early, offers the best prognosis. Chronic constitutional laby-

rinthitis offers a less favorable prognosis. The prognosis is least favorable in the congenital type of syphilitic labyrinthitis. The value of salvarsan in syphilis of the labyrinth is unsettled.

P. F.

1377

Treatment of Concussion of the Ear. O. HAMM, *Munch. med. Woch.*, No. 48, Nov. 30, 1915.

The author believes that treatment in disturbed function of the internal ear from concussion is promising. He has found otothermia of great service. His results have been such that the majority of the patients recovered their hearing and were returned to the front for further military service. It is only exceptionally that diathermia of the ear is valueless. He advises the treatments weekly. After each treatment the ear should be stopped up with cotton for 1 to 2 hours. Alcohol and especially tobacco should be absolutely prohibited. The duration of the treatment is four weeks. By the same method he has also obtained good results in chronic otitis media not yielding to any other treatment.

P. F.

1403

Lumbar Puncture in Aural and Nasal Cases; Pathology of the Fluid. W. WINGRAVE, *Jour. Laryngol., Rhinol. and Otol.*, V. XXX, p. 7, July, 1915.

The specific gravity is usually raised (1010). The alkaline reaction is always reduced in cases of infection and in some instances it may even be acid. The acidity is said to be due to lactic acid. Its presence may be demonstrated by testing with Uffelman's ????

Sugar is best tested for, mixing equal parts of a weak solution of methylene blue, liquor potassal and cerebrospinal fluid. The color disappears if the fluid is normal but remains unaffected if sugar is absent. Absence of sugar is very important evidence of bacterial infection and should be carefully looked for. The test is also applicable and useful for determining the presence of sugar in nasal discharge and in blood, but it is not applicable to urine. For spinal fluid it is preferable to Fehling's solution.

The bacteriology of the fluid should be studied by films and cultures. The former will differentiate between negative and positive groups. If positive the organisms probably belong to the "pyogen" group (streptococci, staphylococci, pneumococci [Frankel], tetracocci, tubercle, diphtheroid). If negative suspect, *B. capsulatus mucosus*, *B. coli*, *B. typhosus*, *diplo. catarrhalis*, *B. pyocyaneus*, *B. hastillo*, etc.

In chronic middle-ear cases the infection is almost always polymicrobial.

P. F.

1470

Occlusion of the Lateral Sinus as a Hemostatic Measure in Wounds Involving Large Vessels at the Base of the Brain. LANNOIS and PATEL, *Bull. de l'Acad. de Med. de Paris*, April 27, 1915.

In wounds involving large vessels at the base of the brain the danger from secondary hemorrhage is very great, resulting either in prompt death or in the formation of a diffuse hematoma. In two cases where

ligation of the carotid was done death resulted within a week through rupture of the internal jugular vein. In five other cases, however, occlusion of the lateral sinus was followed by recovery, the control of the hemorrhage being immediate. The packing introduced to compress the sinus was left in for twelve days.

P. F.

1521

Studies in the Localization of Cerebellar Tumors: Posterior New Growths Without Nystagmus. E. G. GREY, *JOINT. A. M. A.*, Oct. 16, 1915.

Nystagmus is recognized as a most important localizing sign in diseases of the posterior cranial fossa, especially lesions of the cerebellum, cerebellar peduncles, vestibular nerves, the neighboring nuclei and labyrinths. When nystagmus occurs in a patient showing symptoms of sub-tentorial tumor the presence of either an extracerebellar or an intracerebellar neoplasm is probable. Nystagmus may, however, be absent or it may be found in patients with tumors lying anterior to the cerebellum, which fact has somewhat restricted its usefulness as a sign of disease in the posterior fossa. The type of nystagmus in patients with tumor anterior to the cerebellum and in those with cerebellar or extracerebellar growths is occasionally identical.

Grey enters into a discussion of the mechanism of nystagmus. According to Bauer and Leidler, nystagmus is the result of lesions in the region of Deiters' nucleus. The horizontal jerks originate more from the ventrocaudal part, the vertical jerks from the oral part. Barany also assumes the existence of a nystagmus center in this region, the nystagmus being caused by increased tension in the posterior fossa. The presence of a nystagmus center is contested by Wilson and Pike, who believe that an agency somewhere produces a deviation of the eyes from the primary position of equilibrium. In labyrinthine nystagmus this agency lies in the labyrinth, but what agencies are responsible for the development of nystagmus from other portions of the brain we do not know, and just what relation the cerebellum bears to the development of rhythmic movements of the eyes is at present rather uncertain. Destruction of parts of the cerebellar hemisphere and vermis never causes spontaneous nystagmus, yet it causes excessive excitability of the vestibular nuclei. This has led to the view that in cerebellar disease the nystagmus is due to an alteration in function of the vestibular nuclei or their connections with the nuclei of the eye muscles through pressure.

Grey's discussion deals with 34 verified cases of intracerebellar tumor, in one-third of which there was no nystagmus, and with 40 verified cases of tumor situated anterior to cerebellum, in eight of which there was nystagmus previous to operation. In both groups, however, the same degree of intracranial pressure was found. All the cases in which nystagmus was absent contained intracerebellar new growths, which suggests that in the presence of a cerebellar tumor syndrome, but without nystagmus, the absence of this symptom points towards an intracerebellar localization of the lesion.

Caloric tests made in six of the 32, and seven of the 40 cases showing the absence of nystagmus resulted in characteristic nystagmus from either labyrinth.

P. F.

1524

Bilateral Lesion of the Auditory Centre. THOMAS GUTHRIE, *Jour. Laryngol., Rhinol. and Otol.*, Vol. XXX, No. 5, May, 1915.

The author considers the features of six previously reported cases (Mott, Benninghaus, Kahler and Pick, Mills, Wernicke and Friedländer, Pick) and places a seventh on record. This patient, male, twenty-two years of age, suffered complete deafness following an attack of apoplexy. Except for the deafness, which remained, he subsequently recovered completely. Lesion of the labyrinth or eighth nerve was excluded by the fact that while the cochlear functions were abolished the vestibular reactions were normal. The presumption that the deafness is of central origin is very strong, although autopsy findings are absent in this case to confirm this view. The special feature in this case, not present in the other cases on record, is the association of complete and permanent deafness with perfect recovery of all other functions. P. F.

1525

Tuberculoma of the Pons Varolii. L. GUTHRIE, *British Jour. Child. Dis.*, V. 12, No. 140, Aug., 1915.

The following symptoms were present: Hearing not affected. Tongue protruded straight; soft palate moved normally. When patient was sitting up the chin constantly deviated to the left and head inclined towards the left. Eyes negative. Complete sixth nerve paralysis; no outward movement beyond midline. Complete paralysis of left side of face. No weakness, hypotonia, tremors, ataxia or increase in reflexes in upper extremity. Same in regard to lower extremities; active knee-jerk on right side. No ankle clonus. Plantar reflexes sluggish. Slight vertical nystagmus. Death from convulsions. Section: tuberculous mass about one inch in diameter in center of the pons, much more diffused, histologically, than the gross appearance indicated. Several metastatic foci beneath cortex on frontal lobes and one on cribriform plate of ethmoid. P. F.

1528

A Rare Case of Cerebellar Abscess. B. HASELTINE, *Jour. Ophthal., Otol. and Laryngol.*, Aug., 1915.

The interest attached to this case lies in its being a bilateral cerebellar abscess, with recovery. A review of the history is as follows: Patient, a young man of 18 years. It began with an acute purulent rhinitis. The patient had previously never had any ear trouble or infections of any kind. Soon after the infection started he developed a middle ear infection with spontaneous rupture of the right drumhead and a profuse purulent discharge from the right ear. At this time the temperature was 98 degrees, pulse 92; the patient complained of vertigo, with a tendency to fall to the right side, which tendency remained throughout the illness, dilated pupils, with no reaction to light, occasional vomiting. About ten days later there developed paralysis of the right arm and leg, subnormal temperature and slow pulse (54-66). There was slight choked disc on each side. Hearing was normal. Examination of the spinal fluid showed fifteen lymphocytes per c.mm.; no red blood cells, no polymorphonuclears. No growth culturally. Nonné test (both phases), negative; Noguchi glob-

ulin test, negative. The mastoid had been entirely negative. The middle ear had become normal.

Simple mastoid operation performed. Cells were not broken down and no pus present. Cells were distinctly darkened and softened. Culture showed staphylococcus aureus. The labyrinth was exposed but not opened. The dura over the cerebellum in Trautmann's triangle was exposed and found normal in appearance. Mastoid cells were thoroughly cleaned out.

The immediate results of this operation were very good; the temperature returned to normal, the rotary nystagmus, which had been marked, disappeared in forty-eight hours; no vomiting; lessening of the pupillary dilatation.

On the seventh day after operation there was a recurrence of the cerebellar symptoms (interference with muscular co-ordination) with recurrent vomiting.

Mastoid wound re-opened and explored. Also the right temporo-sphenoidal lobe which, however, was normal. Trephined over the cerebellum and large cerebellar abscess uncovered. Immediate amelioration of the symptoms. Six days later, patient suddenly grew worse. X-ray taken, which showed extension of the abscess into the left cerebellar hemisphere. Left abscess was drained through the right opening. Culture showed staphylococcus aureus. Recovery.

P. F.

1547

Fracture of the Base of the Skull with Escape of Cerebrospinal fluid from the Ear. The Effect of Atropine and Epinephrin Upon the Secretion. J. WALKER MOORE, *American Journal of the Medical Sciences*, Vol. CXLIX, No. 4, page 380, April, 1915.

Reports a case, which resulted fatally. A careful autopsy was made and the cerebral findings, both gross and microscopic, are given. The evolution of the hypothesis that the cerebrospinal fluid is a true secretion is given, and the literature of the subject quite fully summarized. Then follows a series of tables, based upon the author's case, giving the quantity and composition of the cerebrospinal fluid at stated periods, showing the results from the administration of doses of atropine and epinephrin, and accompanied by remarks on the clinical condition of the child at the time of observation.

PACKARD.

1548

Diagnosis of Tumor in Cerebello-Pontine Angle. G. H. M. KROHN, *Norsk. Mag. for Lægevid.*, V. 76, No. 6, June, 1915.

The symptoms complained of by the patient, a woman of 25 years of age, were headache, occasional choked disc, vomiting, nystagmus and the cerebellar speech and gait. There were also paralysis of the fifth, sixth, seventh, eighth nerves on the same side as the tumor, hypotony, asynergy, adiadokokinesia, catalepsy, deviation and inability to draw parallel lines between two vertical lines (dysmetria). There were hemiparesis, hemianesthesia and positive Babinski on the opposite side.

P. F.

1550**Case of Temporosphenoidal Abscess Discovered by Exploration Through Multiple Incisions in the Dura; Remarks on Operative Technique.**

J. R. PAGE, *Surg., Gyn. and Obst.*, p. 718, June, 1915.

The author believes that the danger of infection from without as a result of small incisions in the dura, contiguous to the mastoid wound, is comparatively slight. The wound should, however, be thoroughly curetted and every crevice packed with gauze saturated with a strongly antiseptic solution before the dura is even uncovered. Freshly uncovered dura is sterile, but even if it be momentarily contaminated by the surgeon's gloves or instruments, the contamination does not have time to become established as an infection on its surface, and before an incision is made, gloves, instruments and sponges should be changed and the recently exposed dura is washed off with an antiseptic solution (alcohol) followed by saline.

Where granulations cover the dura from prolonged epidural infection experience has shown that the meninges beneath such an area are more resistant, because of the adhesion, to infection, even though antiseptics cannot reach all the bacteria infiltrated in the granulations. The advantage of small multiple incisions is that the whole wound can be left open to allow free access to the area for drainage without incurring the danger of hernia-formation or the necessity of closely suturing the wound for the same reason, which not only interferes with proper drainage but increases the danger, through exposure of a large area of cortex, to meningeal infection from within as well as from without.

P. F.

1555**Intracranial Telangiectasis: Symptomatology and Treatment, with Report of Two Cases.** ERNEST SACHS, *Amer. Jour. Med. Sc.*, Oct., 1915.

The clinical picture of intracranial telangiectasis consists of (1) attacks of Jacksonian epilepsy occurring in no-syphilitic persons at long intervals; (2) unconsciousness of long duration; (3) no evidence of increased intracranial pressure; (4) a very slow progression of symptoms; (5) telangiectases on the head or face.

Two typical cases, each in a boy ten years old, are reported by the author, who emphasizes that it is essential to distinguish between a true angioma—a neoplasm—and the telangiectasis observed in these cases which was congenital and not a neoplasm.

P. F.

1571**Diagnosis of Tumors in the Posterior Cranial Fossa.** T. H. WEISENBURG and P. WORK, *Jour. A. M. A.*, Oct. 16, 1915.

The authors believe that the chief function of the cerebellum is to synergize all movements of the body. All other symptoms in addition to the cerebellar must be taken into consideration to accurately diagnose a cerebellar lesion. They have often made a diagnosis of labyrinthine lesion only to find the cerebellum involved, and vice versa. According to the authors, lesions of the cerebellum itself cause more strictly limited symptoms than those invading the peduncles. Most cerebellar tumors are gliomatous and tend to involve the middle rather than the outer part

of the cerebellum; the vermis is practically always involved. Here are centered the synergic movements of the upper trunk and in the lower vermis the movements of the lower trunk. When the vermis is involved the staggering is mainly forward or backward; when a lateral lobe is involved the sway of the body is to the side of the lesion. When the lateral lobes alone are involved the asynergic movements are present only on the side of the lesion in the upper limb if the superior lobe is involved and in the lower limb if the inferior lobe is involved. If nystagmus is developed by voluntary movement the lesion is probably extracerebellar and the presence of cranial nerve symptoms also indicates that the lesion is extracerebellar. Vertigo associated with auditory disturbance is not a cerebellar symptom. It is supposed that the inferior and middle peduncles transmit impulses to the cerebellum and the superior peduncle transmits impulses from the cerebellum. It is probable, however, that the peduncles transmit impulses in both directions. In lesions of the middle cerebellar peduncle the symptoms consist of fifth or sixth nerve disturbance on the same side as the lesion with sensory and motor symptoms on the opposite side. In lesions of the inferior peduncle (the authors state that they have never seen a tumor limited to this peduncle, though they have seen extensions of growth into one or both), the associated symptoms, if the lesion extends into the medulla, show involvement of the vestibular tract and of the ninth, tenth and twelfth cranial nerves. If in addition to cranial nerve symptoms there is cerebellar asynergy in the trunk and limbs it is probable that the tumor grows either from the cerebellum or from the pons.

P. F.

1633

A New Tonsil-Holding Forceps. S. C. GLIDDEN, *Surg., Gyn. and Obstet.*, Nov., 1915.

The teeth in this instrument are similar to those in a vulsellum forceps. The opening between the teeth is $1\frac{1}{4}$ inches, allowing the tonsil to be firmly grasped either longitudinally or transversely. The handle is at an acute angle, with a lock at the end of it. The handle is small enough to allow a snare to be slipped over it without removing the forceps after the tonsil is grasped.

P. F.

1635

A New Tongue Depressor Designed to Facilitate the Siuder Method of Operation. JAMES, H. HEACOCK, *New York Medical Journal*, Dec. 18, 1915.

A new instrument designed to fit on the index finger and used as a tongue depressor, allowing more space to work.

PACKARD.

1644

A Pharyngeal Tube Combining Rebreathing and Insufflation. F. W. PINNES, *Jour. A. M. A.*, Oct. 9, 1915.

It is a one-piece metal blade with upturned sides to fit the contour of the tongue, having a channel for introduction of an oral catheter five inches long. It is curved from the lips to the pharynx, the mouth end being a flattened tube and adjustable for different sized patients. Its ad-

vantages are (1) it provides for automatic holding of the tongue forward without other means; (2) it has not the objections of a closed tube, being an open blade with channels easily cleaned; (3) it can be used in patients of different sizes; (4) it has a channel for introduction of an oral catheter for pharyngeal insufflation, the catheter not being part of the pharyngeal tube; (5) the anesthetic may be stopped while the blade remains in place; (6) it is efficient in holding the tongue forward; (7) it is efficient for rebreathing, keeping the teeth and lips apart and the tongue forward; (8) it does not get clogged with mucus or blood.

P. F.

1648

Accessory Sinus Angle Finder. F. M. LAW, *Amer. Jour. Roentgenol.*, Nov., 1915.

In order to obtain a satisfactory anteroposterior image of all the accessory sinuses at one exposure it is necessary that the rays pass through the head at a certain angle. The proper angle is one which will cause the shadow of the petrous to cut across the lower one-third of the orbit. To obtain this position the principal ray must be directed at an angle of 23 degrees from a line extending from the external auditory meatus to the glabella. The instrument described by the author is a means of finding this angle. He describes the technic of using it.

P. F.

1720

Combined Extracranial Paralysis of Cerebral Nerves. J. C. BECK and G. B. HASSIN, *Medical Record*, Aug. 21, 1915.

The patient was an 18-year-old girl and presented the following symptoms: (1) Pain in the neck, occiput and forehead; (2) marked hemiatrophy of the tongue; (3) a combination of tongue atrophy with paralysis of the ninth, tenth and eleventh cranial nerves; (4) nystagmus. The eighth nerve was also involved, the involvement being bilateral and the deafness was due to otosclerosis. The patient's mother and two sisters also suffer from deafness due to some pathological cause. The nystagmus was marked only during active movements of the eyeballs and latterly showed slow oscillatory movements to the left or right, according to the direction of the intentional movements of the eyes, without the quick return to the opposite, which is, according to Wilson and Pike, the only essential movement of a so-called labyrinthine nystagmus. The question arises if, in view of the presence of otosclerosis, this nystagmus in this patient was not of vestibular origin. The essential trouble of the patient was tuberculosis of the cervical glands with secondary involvement of the ninth, tenth, eleventh and twelfth nerves in their extracranial course. The authors believe that in this case the nystagmus and deafness are independent lesions. The diagnosis was verified by x-ray examination, successful operation and subsequent progressive improvement.

P. F.

1810

Calcium Factor in Hemophilia. O. F. HESS, *Bull. Johns Hopkins Hosp.*, Nov., 1915.

Typical hereditary hemophilia is not associated with calcium deficiency. The author's studies show that the addition of calcium to blood *in vitro*

delays rather than hastens coagulation, yet quantitative studies of the lime content of the patient's blood (a boy nine years old) showed that there was a deficiency compared with the normal, which became positive when lime salts were added to the diet. This particular case seemed to be a type of *hemophilia calcipriva*.

P. F.

1839

Use of Pituitary Extract as a Coagulant in Nose and Throat Surgery... H.

KAHN and L. E. GORDON, *Jour. A. M. A.*, Jan. 23, 1915.

To control hemorrhage in nose and throat surgery the dose is twelve minims in children, and fifteen minims in adults. This is administered hyperdermically not less than fifteen minutes before anesthesia. The authors have employed pituitary extract in this matter in over one hundred cases and they find that the coagulation time of the blood is considerably reduced as well as the hemorrhage, following turbinal operations especially. Systolic pressure was increased in 55.31 per cent of cases, reduced in 36 per cent and unchanged in 8.5 per cent. Diastolic pressure was increased in 35.5 per cent of the cases, reduced in 35.5 per cent, and no change in 29 per cent.

P. F.

1928

Some Ocular Complications of Mumps. F. RAMOND and G. GOUBERT,

Bull. et mem. Soc. Med. des Hop. de Paris, July 29, 1915.

The authors record some little known ocular complication of mumps met with by them in a military hospital. Out of 115 cases of mumps lacrimation was present in 45, affecting chiefly the bulbar conjunctiva. In one case dacryoadenitis was present. Of 16 cases in which the fundus was examined some showed a markedly increased rosy color of the optic disc; others showed a softness of the disc with disappearance of its contour; others showed swelling of the veins of the retina, and one case showed a congested iris. Another case showed papillary stasis. All the with the disappearance of the inflammation of the parotid gland. In a few instances they persisted for a month after recovery from the mumps.

P. F.

1945

Oto-laryngological Cases from a Military Hospital. ARCHER RYLAND, *Jour.*

Laryngol., Rhinol. and Otol., V. 30, No. 8, p. 304, Aug., 1915.

Case 1. Pan-sinusitis of long standing treated by radical operation on affected sinuses.

Complaint: nasal obstruction on both sides; foul discharge from both sides of nose and impairment of sense of smell. Left deviation of septum and basal spur on right. Large polypi in left nasal fossa, which were removed. The following operations were performed in the order named: (1) Caldwell-Luc on left maxillary antrum, which had been lavaged daily for three weeks previously without cessation of pus formation. (2) Ethmoid and fronto-ethmoidal curettage. (3) Luc's operation on left frontal sinus. (4) Exposure of sphenoid and removal of neighboring diseased mucosa.

The total treatment extended over a period of ten weeks. Accessory sinuses free from pus. Patient returned to military duty.

Case 2. Suppurative otitis media; acute mastoiditis and labyrinthitis.

Patient, 24 years old, had a left chronic otitis media. Acute mastoiditis supervened, followed a few days later by acute labyrinthitis. Operation decided upon by increase of mastoid tenderness, increase of vertigo and persistence of nystagmus and vomiting. Radical mastoid performed. No fistula present and no operative measure on labyrinth. Vertigo and vomiting ceased within forty-eight hours after operation. Uneventful recovery.

Case 3. Severe self-inflicted wound of the larynx; primary suture; tracheotomy; recovery.

Recovery without pulmonary complications, hemorrhage or interference with laryngeal movements.

P. F.

1975

Acute Actinomycosis of the Parotid Gland. E. D. TELFORD, *Brit. Med. Jour.*, Oct. 9, 1915.

Two cases are reported by Telford in which there was remarkable similarity in onset and symptoms. In each case the route of infection was by the parotid duct and from this point of entry the disease rapidly infiltrated the gland itself. In one case the infection was derived, it would seem, from the habit of chewing corn whilst engaged in feeding poultry. In the other case no exact source of infection could be discovered except that a week before the onset the patient had played in a ripe field of corn.

The features of acute actinomycosis would appear to be as follows: The fungus enters by the parotid duct and within a few days gives rise to an acute parotitis. The disease then bursts through the limits of the gland, followed by acute cellulitis involving the neck and even extending far over the scalp. Symptoms of constitutional sepsis are marked. Should the swollen parts be incised no pus will flow, but merely a bloody debris and the cut tissue will be found to be diffusely infiltrated and of a dirty gray color, flecked with yellow.

In treatment, iodine has appeared to yield good results in the more chronic cases and should be used freely in the acute cases. The internal administration of KI should be pushed freely. The incisions and sinuses should be irrigated with a weak mixture of tincture of iodine and water. In the author's second case 0.3 grams of neo-salvarsan was given as soon as the diagnosis was established on the twelfth day of the illness. Within twenty-four hours there was a marked increase in the amount of discharge and a rapid improvement in the local and general condition.

P. F.

2011

Oral Surgery. A Treatise on the Diseases, Injuries and Malformations of the Mouth and Associated Parts. DR. TRUMAN W. BROPHY, Philadelphia and London: P. Blakiston's Son & Co., 1915.

That this book by Dr. Brophy is an exhaustive work goes without saying. As can readily be understood the best chapters are those on hare-lip and cleft-palate. The latter comprises 162 pages and the detailed description of operative technic and the instructive plates and drawings elucidating the operative procedure makes this feature of the book a

masterpiece. Either chapter is well worth the price of the book, for we have here a presentation of the author's extensive experience in a field where he is pre-eminent. The honors are shared also by the splendid chapters on the embryology, development and disease of the tongue, fracture of the jaw and the bones of the face, maxillary and mandibular tumors, cysts, dentoalveolitis and infections of oral origin.

On perusal of the work, however, the impression becomes evident that it suffers from some duplication of subject-matter and that a good deal might have been omitted without deflecting from the value of the book as a whole. This applies especially to the chapters on blood examination, eugenics and infant feeding. The book does not contain a chapter on Roentgenography of the skull—which would be perfectly *apropos*—notwithstanding that numerous skiagraphs are scattered throughout the book. The chapter on eugenics merely points out the congenitality of hare-lip and cleft-palate, facts referred to under their respective heads, so that this chapter could have been omitted. Why the book should include discussions on wet-nurses, formulae for infant feeding, baby's stools and other encroachments on the province of pediatrics is not clear, especially as the author, Dr. F. W. Belknap, states at the outset "the various means of introducing food into the stomach of a child with a cleft-palate have been described in another chapter." Why not go a step further and consider the infectious diseases that a cleft-palate child, like any normal child, is likely to have? The chapter on general syphilis is quite unnecessary, too, because syphilis in the mouth and associated parts is taken up under the respective heads.

But these are faults which may be remedied and which will not prevent the book from becoming one of the standards of medical literature. Finally, the 39 colored plates and the 909 photographs and other illustrations are excellent products on which the publishers deserve to be complimented.

P. F.

2017

Anesthetics. Their Use and Administration. DR. DUDLEY WILMOT BUXTON, Philadelphia and London: P. Blakiston's Son & Co., 1915.

The scope of the book may best be comprehended by an outline of its chapters, of which there are twelve. Chapter I deals with the historical aspects of anesthetics. Chapter II, preparation for operation; selection of the anesthetic and method. Chapter III, nitrous oxid gas. Chapter IV, ether—hedonal. Chapter V, chloroform. Chapter VI, ethyl chlorid and its mixtures—ethyl bromid and the less commonly used anesthetics. Chapter VII, alkaloidal drugs with general anesthetics or with local analgesics; anesthetic mixtures, successions, and solution. Chapter VIII, anesthetics and analgesics in special surgery. Chapter IX, anesthetics in obstetric practice. Chapter X, the accidents and after effects of anesthesia and their treatment. Chapter XI, local analgesia—spinal analgesia. Chapter XII, medico-legal aspects of the administration of anesthetics.

It is thus seen that the consideration of the subject of anesthetics has been carried out from every angle. The views expressed by the author on the debatable problems in anesthesia will be found to be the result of mature judgment and wide experience. The book cannot be too highly recommended, especially to those who are primarily interested in the subject of anesthesia.

P. F.

2019

Diseases of the Nose and Throat. DR. ALGERNON COOLIDGE, Philadelphia and London: W. B. Saunders Co., 1915.

The evident purpose of the author was not to write a reference volume covering the diseases and surgery of the nose and throat, but a comprehensive introduction to the subject for the use of students. This purpose Dr. Coolidge has amply fulfilled for the characteristic feature of the book is its simple presentation of the subject, devoid of involved references and academic discussions so that the student just beginning the study of rhinology and laryngology is able to get a very comprehensive understanding of the subject, which he can amplify later if he should decide to go into these branches of medicine more extensively. The simplicity of the book is even noticeable in the illustrations, which are clear-cut drawings showing the relations of parts very distinctly and for the purpose in view are surely to be preferred to expensive plates and photographs of pathological specimens which at times utterly fail to give the reader a clear idea of what they are supposed to show. In view of the excellent presentation of the subject the book should have a wide circulation.

P. F.

2021

The Nose, Throat and Ear; Their Functions and Diseases. A Treatise Upon the Breath-road, Food-road and Accessory Organs. DR. BEN CLARK GILE, Philadelphia and London: P. Blakiston's Son & Co., 1915.

For teaching purposes as well as for the general practitioner Dr. Giles' book will be found very useful. Of the smaller works on oto-laryngology it has originality and is perhaps one of the best arranged. The subject-matter in each chapter is very completely considered. Operative measures are not neglected or dismissed in a short paragraph. The operation for tonsil enucleation by snare, the mastoid operation, the intranasal opening of the frontal sinus, etc., are all very fully described. The book also contains a short but comprehensive chapter on otosclerosis and another on intracranial complications from ear disease. The illustrations are good.

P. F.

2027

Diseases of the Throat, Nose and Ear. DR. WILLIAM H. KELSON, Oxford University Press, 1915.

For the purposes of the medical student this manual by Dr. Kelson will be found quite sufficient. The subjects are treated concisely but not to the sacrifice of clearness. The description of the tonsil operation might have been carried out with greater detail and the technic brought up to more modern standards. The operation as described deals more with tonsillotomy by the Mackenzie guillotine, the author mentioning his preference for enucleation by snare, in a *sotto voce* fashion only. The best chapter is that on the ear. The mastoid operations are referred to merely in their general indications. The functional tests for hearing are discussed quite sufficiently so that the student should be able to obtain a very comprehensive idea of their significance. The book also contains an appendix on general therapy including the use of electricity, x-rays, radium and vaccines in oto-laryngology.

P. F.

2035

Injuries to the Eyes, Nose, Throat and Ears. MAJOR ANDREW M. RAMSAY, MAJOR J. DUNDAS GRANT, CAPT. H. LAWSON WHALE and CAPT. CHARLES E. WEST, Oxford War Primers, New York: Oxford University Press, 1915.

The chapter on injuries of the eyes is treated by Dr. Andrew M. Ramsay; the chapter on nose and throat by Dr. J. Dundas Grant and Dr. H. Lawson Whale and the chapter on the ear by Dr. Charles E. West. The book covers only 156 pages, but it is replete with information dealing largely, of course, with wounds of the head which directly or indirectly involve the special sense organs. It is evident that the authors have had wide experience in the military aspects of the surgery of these parts and the little book is written exactly from that point of view and for medical men contemplating military service. A chapter consisting of thirty-eight illustrative cases is included, together with x-ray plates of the findings, a statement as to the operation performed and the results of same. The chapter on injuries of the ears might have been enlarged. One statement particularly by Dr. West is worthy of repetition: "It is remarkable with what completeness the possession of an antecedent middle-ear deafness seems to screen the labyrinth from damage by explosion." The value of this observation on the medical observation of recruits, as far as the ear is concerned, is quite evident.

P. F.

2045

Squint: Its Causes, Pathology and Treatment. DR. CLAUD WORTH, London: John Bale, Sons and Danielsson, 1915.

The fact that a fourth edition of this work has been found necessary speaks for itself. This edition does not differ materially from the preceding editions. What is especially characteristic of the book is the thoroughness with which the subject is considered, the easy flowing style of language, and the frequent references to the author's personal experiences and methods of treatment including operative technic. As a work of authority on the subject of squint, Dr. Worth's book will continue to hold first place.

P. F.

2046

What the Mother of a Deaf Child Ought to Know. PROF. JOHN DUTTON WRIGHT, New York: Frederick A. Stokes Company, 1915.

As an educator of a quarter of a century's experience, Prof. John Dutton Wright can speak authoritatively of the duties of the mother of the deaf child and as a man interested in the humanitarian aspects of the deaf child's education he puts his appeal for the co-operation of mothers in a most interesting and helpful style. The book will appeal especially to the average busy mother who wishes to know what hopes and possibilities can be counted on for her little deaf child and how she can best use the time at her disposal in laying the foundations for the child's oral education. She will learn in these chapters, in language devoid of all technicality, just what she is to do to help efface the handicap of deafness, and affliction which she sees will make her deaf child different from others unless it is offset. She will learn the value of beginning to talk

to the deaf child, to train his eye to take the place of his ear as far as possible, and how to encourage his efforts at imitating speech so that his vocal organs will not become stiffened through inaction.

Dr. Wright's book should have even a wider appeal to friends of deaf children, to social workers, physicians, teachers and others who come in contact with these unfortunates. The child whose mother has intelligently assimilated the contents of this little volume and acted upon its practical and valuable suggestions, and whose friends have adopted the habit of talking to the child and treating him like other children, will enter a regular oral school for the deaf, with the immeasurable advantage of having the ground already prepared for the work of the teacher. It is certainly to be hoped that "What the Mother of a Deaf Child Ought to Know" will find its way into every home where a deaf child is living.

ETHEL M. HILLIARD.

